

ECOLOGICAL SOCIETY OF AMERICA

Ecology/Ecological Monographs/Ecological Applications

PREPRINT

This preprint is a PDF of a manuscript that has been accepted for publication in an ESA journal. It is the final version that was uploaded and approved by the author(s). While the paper has been through the usual rigorous peer review process of ESA journals, it has not been copy-edited, nor have the graphics and tables been modified for final publication. Also note that the paper may refer to online Appendices and/or Supplements that are not yet available. We have posted this preliminary version of the manuscript online in the interest of making the scientific findings available for distribution and citation as quickly as possible following acceptance. However, readers should be aware that the final, published version will look different from this version and may also have some differences in content.

The doi for this manuscript and the correct format for citing the paper are given at the top of the online (html) abstract.

Once the final published version of this paper is posted online, it will replace the preliminary version at the specified doi.

1	Running head: Integrated approach to bycatch research
2	
3	Title: Bycatch mortality of endangered coho salmon: impacts, solutions,
4	and aboriginal perspectives
5	
6	Graham D. Raby ¹ *, Michael R. Donaldson ² , Vivian M. Nguyen ¹ , Mark K. Taylor ¹ , Natalie M.
7	Sopinka ² , Katrina V. Cook ¹ , David A. Patterson ³ , David Robichaud ⁴ , Scott G. Hinch ² , and
8	Steven J. Cooke ¹
9	
10	¹ Fish Ecology and Conservation Physiology Laboratory, Department of Biology and Institute of
11	Environmental Sciences, Carleton University, 1125 Colonel By Dr., Ottawa, ON, K1S 5B6,
12	Canada
13	² Pacific Salmon Ecology and Conservation Lab, Department of Forest and Conservation
14	Sciences, University of British Columbia, 2424 Main Mall, Vancouver, BC, V6T 1Z4, Canada
15	³ Fisheries and Oceans Canada, Science Branch, Pacific Region, Cooperative Resource
16	Management Institute, School of Resource and Environmental Management, Simon Fraser
17	University, 8888 University Dr., Burnaby, BC, V5A 1S6, Canada
18	⁴ LGL Limited Environmental Research Associates, 9768 Second St., Sydney, BC, V8L 3Y8,
19	Canada
20	
21	
22	
23	*E-mail: grahamraby@yahoo.com

Abstract. We used biotelemetry and human dimensions surveys to explore potential solutions to 24 migration mortality of an endangered population of coho salmon caught as bycatch in an 25 aboriginal beach seine fishery. From 2009 to 2011, 182 wild coho salmon caught as bycatch in 26 27 the lower Fraser River (Canada) were radio-tagged and tracked as they attempted to complete their migrations to natal spawning areas over 300 km upstream. Failure to survive to reach 28 terminal radio receiving stations averaged 39% over three years. This mortality estimate is low 29 compared to those obtained from telemetry studies on other salmon fisheries in the Fraser River. 30 However, this value is markedly higher than the mortality estimate currently used to manage the 31 fishery's impact. It is also in contrast to the perceptions of the majority of aboriginal fishers, who 32 did not think survival of coho salmon is affected by capture and release from their fishery. 33 Increased probability of survival was associated with lower reflex impairment which is 34 consistent with previous findings. Reflex impairment was positively correlated with 35 entanglement time, suggesting that greater efforts by the fishers to release bycatch from their 36 nets quickly would minimize post-release mortality. Survey responses by aboriginal fishers also 37 suggested that they are receptive to employing new bycatch handling methods if they are shown 38 to increase post-release survival. However, attempts to facilitate revival of a subset of captured 39 fish using cylindrical in-river recovery bags did not improve migration success. Fisheries 40 managers could use the new information from this study to better quantify impacts and evaluate 41 different harvest options. Since aboriginal fishers were receptive to using alternate handling 42 methods, efforts to improve knowledge on minimizing reflex impairment through reductions in 43 handling time could help increase bycatch survival. Such a direct integration of social science 44 and applied ecology is a novel approach to understanding conservation issues that can better 45 46 inform meaningful actions to promote species recovery.

Key words: By-catch, discards, fisheries management, biodiversity conservation, radio telemetry,
conservation social science, RAMP, integrative science, Pacific salmon.

49

50 INTRODUCTION

There are numerous examples in the literature of conservation scientists calling for an 51 integration of the social sciences into biological research, and for the need to bridge the gap 52 between science and conservation action (e.g., Campbell 2005, Fox et al. 2006, Lowe et al. 2009, 53 Sutherland et al. 2009, Margles et al. 2010). Yet, research papers that include and integrate both 54 sociological and biological data remain uncommon, despite their obvious potential to develop 55 more "actionable" science (Cook et al. 2013). There are some examples. Irvine et al. (2009) 56 interviewed managers of red deer Cervus elaphus populations, and combined their perspectives 57 with scientific data to build habitat use models for the species. Interview approaches alone have 58 been used to build models – such as for assessing threats to endangered sea turtles (Donlan et al. 59 2010). Donaldson et al. (2013) demonstrated that comparative physiology and radio telemetry 60 could be combined with human dimensions surveys to address revival strategies for angled and 61 released sockeye salmon Oncorhynchus nerka. Given that application of the best science can be 62 improved with stakeholder input, incorporating human dimensions data into conservation science 63 is a natural fit. 64

Pacific salmon *Oncorhynchus* spp. are among the most well studied wild animals. Their contribution to human culture, economies, and the functioning of ecosystems, coupled with the fact that many populations are in decline means that they receive enormous research attention from conservation scientists (Scarnecchia 1988, Gende et al. 2002). Their importance as a natural resource and the complexity of socio-political considerations in how they are managed

(Scarnecchia 1988, Lackey 1999) make this a particularly fruitful area for integrative research. In 70 the Fraser River, British Columbia, Canada, the fishery for Pacific salmon is complex, involving 71 different user groups (aboriginal, recreational, and commercial) which target several species 72 73 comprising hundreds of unique populations, many of which migrate upriver toward spawning grounds at the same time. Inherent in managing these fisheries is the objective that diversity be 74 maintained, both within and among species (DFO 2005). Interior Fraser River coho salmon (O. 75 kisutch that spawn in tributaries of the Fraser River upstream of Hell's Gate; Fig. 1) have 76 received particular attention in recent years owing to their listing as endangered by the 77 Committee on the Status of Endangered Wildlife in Canada (COSEWIC). This population of 78 coho salmon experienced a $\sim 60\%$ population decline during the 1990s, attributed mainly to 79 overfishing and habitat alteration (Bradford and Irvine 2000). Directed harvest of wild coho 80 salmon in British Columbia was closed in 1999 and the population has since stabilized at 20,000-81 30,000 returning adults, an incomplete recovery to pre-1990s abundance (R. Bailey, DFO, pers. 82 comm.). To ensure that bycatch mortality is not limiting recovery, in-river fisheries for other 83 species have been significantly curtailed at the time of year that interior Fraser coho begin 84 migrating through the river (IFCRT 2006). There have been some exceptions to this management 85 strategy including permitting aboriginal groups to conduct beach seine fisheries in the lower 86 Fraser River targeting pink O. gorbuscha, chum O. keta, or sockeye salmon. Beach seines are 87 thought to enable the live release of bycatch with higher subsequent survival than alternative 88 gear such as gill nets. Regulations for the beach seine fishery state that fishers must release all 89 wild coho salmon that are alive, with an inherent assumption that most of these fish will survive, 90 continue their migration, and ultimately spawn. The agency responsible for managing this fishery 91 92 (Fisheries and Oceans Canada; DFO) applies a 5% mortality rate to coho salmon by catch for the

purposes of accounting for fishing mortality and meeting spawning escapement targets (DFO
2011 - IFMP). However there is very little empirical evidence to inform estimates of beach seine
bycatch mortality. Factors affecting mortality remain poorly understood, limiting the ability of
fisheries to implement practices that can minimize mortality.

When a fish is captured in a fishery, it experiences a suite of physiological disturbances 97 (Farrell et al. 2001, Davis 2002, Clark et al. 2012) and typically, some degree of injury (Chopin 98 and Arimoto 1995). Although the proximate causes of post-release bycatch mortality are not well 99 understood in fishes (Wood et al. 1983), certain components of the capture experience are likely 100 more harmful than others. For example, crowding, hypoxia, air exposure, and exhaustive 101 exercise are all thought to negatively affect fish (Ferguson and Tufts 1992, Davis 2002, Marcalo 102 et al. 2006). Bycatch research can be used to identify specific components of a capture stressor 103 that most heavily influence mortality, information which can lead to improved capture or 104 handling practices (Davis 2002). The magnitude and duration of the capture stressor can 105 influence physiological recovery time (Donaldson et al. 2010), suggesting that minimizing 106 107 capture stress is important. In addition, there are techniques that can facilitate metabolic recovery of fish and, in some cases, improve post-release survival (Milligan et al. 2000, Farrell et al. 108 2001). For example, provision of a dark recovery environment that is free from predators and 109 provides high flow across the gills can promote re-oxygenation of tissues, re-invigorating fish 110 before release (Farrell et al. 2001). Recent work in a sockeye salmon recreational fishery has 111 shown that specially-designed fish "recovery bags" (Donaldson et al. 2013) could be used 112 increase survival in some contexts. 113

This three-year study aimed to quantify capture experience for individual fish and use
radio telemetry to assess survival. Combined with interviews of the aboriginal fishers taking part

in the fishery, this study used an all-encompassing approach to provide results relevant to 116 resource managers. There were three general objectives: 1) estimate immediate (i.e., at the time 117 of capture) and post-release bycatch mortality for wild coho salmon caught in beach seines, 2) 118 119 identify factors associated with mortality, and 3) assess potential ways to manage or reduce mortality. Given that a reflex impairment index can be used to predict delayed mortality in this 120 fishery (Raby et al. 2012), we also explored correlates of reflex impairment. For objectives 2 and 121 3, we hoped to identify specific aspects of the capture experience that were most associated with 122 mortality (e.g., crowding time) and evaluate whether cylindrical in-river fish "recovery bags" 123 could be used to promote survival. Semi-structured interviews were used to understand fisher 124 perspectives on issues relating to each of our objectives. 125

126

127 MATERIALS AND METHODS

128 *Study area and fish capture*

This study took place in the Fraser River watershed (British Columbia, Canada). Data 129 collection occurred over three years (2009-2011) during openings of the aboriginal beach seine 130 fisheries. The fishery was targeting pink salmon in 2009 and 2011, and sockeye salmon in 2010. 131 In each case, Fraser River coho salmon caught as by catch were en-route upstream towards their 132 natal streams to spawn. We attended the fishery to collect data on every day it was open in each 133 year; Sept. 21-23 in 2009, Sept. 15 - 17 in 2010, and Sept. 17 - 19, 22, and 24 in 2011. The 134 mean daily river temperature, measured at a temperature monitoring station upstream of Hope 135 (Fig. 1), was 15.88 °C (range: 15.76-16.08 °C) on the fishing days in 2009, 15.12 °C in 2010 136 (15.07-15.18 °C), and 15.32 °C in 2011 (14.85-15.59 °C). 137

We worked with multiple fishing crews at five different locations (some fishing sites had two or more fishing crews operating adjacent to each other). Almost all of the data collection occurred at three sites: Peg Leg, Seabird, and Peters (Fig. 1), which were 108, 129, and 131 river kilometers (rkm), respectively, upstream of the river mouth. Those sites and crews were chosen based on accessibility, the number of crews, and the size of catches. The two other sites were Mountain Bar (just downstream of Peg Leg – 8 fish tagged) and a location halfway between Peg Leg and Seabird (2 fish tagged; Fig. 1).

The fishing crews used beach seines that were 90 m long x 9 m deep x 5 cm diamond 145 stretch mesh. Nets were pulled out from the riverbank using powerboats with one end anchored 146 on shore and were allowed to drift downstream for $\sim 30-60$ s then pulled by the boat in an arc to 147 a downstream point on shore (total time ~ 5 min), before being pulled in by hand or with a truck 148 until fish were crowded into shallow water (< 0.5 m deep) for sorting (Fig. 2). Attempts at 149 locating and releasing by catch varied among crews, but in most cases crews spent 1 - 3 min 150 searching for bycatch immediately after the net was in shallow water prior to collecting their 151 target species. When coho salmon were found by the fishing crew they were given to us for 152 tagging and biopsy (details below) rather than released directly to the river. Entanglement time 153 was calculated for each fish as the time between the net being pulled to shore and the fish being 154 released from the net. For each set, we also obtained the total number of adult salmon caught 155 from a member of the fishing crew whose duty was to enumerate the catch. For some sets only 156 the number of females was recorded (only females were being retained); for such sets we 157 estimated the total catch by assuming a 50-50 sex ratio. 158

159

160 *Fish sampling and tagging*

161	Upon removal from the seine, coho salmon were immediately placed in individual fish
162	holding bags for tagging, biopsy, and measurements. The bags were cylindrical (1 m long x 30
163	cm diameter) and made of Hypalon (thick synthetic rubber) with fine mesh ends that allowed
164	river water to flow through. Radio transmitters were gastrically inserted; a rapid (< 10 s) tagging
165	method with no anesthesia that has been validated and used extensively in Fraser River salmon
166	(e.g., Cooke et al. 2005, English et al. 2005, Martins et al. 2011, Wilson et al. 2013). We used
167	two models of individually-coded radio transmitters that were functionally identical with the
168	same transmission frequencies, dimensions (16 mm diameter, 46 mm long, 460 mm long
169	antenna), and weight (17 g in air, 7 g in water; Pisces 5, Sigma-Eight Inc., Newmarket, Ontario,
170	Canada; and MCFT-3A-3 V, Lotek Wireless Inc., Newmarket, ON).
171	Once tagged, each fish was measured for fork length (FL, nearest cm) and any apparent
172	injuries were noted. A ~ 0.5 g piece of adipose fin tissue was removed from each fish using a
173	hole punch, and stored in 95% ethanol for DNA analyses. Finally, we conducted a rapid (< 20 s)
174	assessment of the presence or absence of five simple animal reflexes (identical to Donaldson et
175	al. 2012, Raby et al. 2012, Raby et al. 2013, Brownscombe et al. 2013, Cooke et al. 2014). This
176	method, called RAMP ('Reflex Action Mortality Predictions'), was validated and summarized
177	using data from 2009 (Raby et al. 2012). The first reflex assessed was tail grab, considered
178	impaired if the salmon failed to exhibit a burst swim response after having its tail grabbed by the
179	handler (up to 3 attempts, or grabs). For tail grab, the fish was underwater in the bag and
180	otherwise not held or impeded from attempting to swim. Next, the fish was held around the
181	middle of its body using two hands (forming a ring with the index fingers and thumbs) and lifted
182	into the air, just above the water's surface: this reflex (body flex) was assessed as impaired if the
183	fish failed to vigorously struggle free of the handler within 3 s. Simultaneously, we assessed

'head complex' – whether the fish exhibited a regular pattern of ventilation. Vestibular-ocular 184 response (VOR) was also assessed - whether the fish's eye rolled to track the handler and 185 maintain level pitch when the handler rolled the fish along its lengthwise axis (a.k.a. ocular 186 counter-rolling). Finally, upon release, orientation was assessed by releasing the fish upside-187 down in the water column just below the surface - this reflex was recorded as impaired if the 188 salmon failed to right itself within 3 s. The reflex responses were cumulated into a RAMP score 189 that represented a proportion of reflexes that were impaired, with a higher score thus indicating a 190 fish in poorer condition that is less likely to survive (Davis 2010, Raby et al. 2012). Following 191 tagging, measurements, and the reflex assessment, coho salmon were released to resume their 192 migration towards spawning areas (except for fish exposed to the recovery bag treatment – see 193 below). 194

Beginning in 2010, tagged coho salmon were chosen at random for an experimental 195 treatment to evaluate whether a cylindrical recovery bag could be used to increase post-release 196 survival. The bags that were used for facilitated recovery (Fig. 3) were narrower than those used 197 for tagging (20 cm diameter x 1 m length) and had much larger mesh (4 cm diameter rigid 198 diamond mesh) to maximize the flow of river water through the bag along its longitudinal axis. 199 Once each fish was transferred into the recovery bag facing into the current, the lengthwise 200 zipper along the top was closed and the bag was attached to a 1.5 m long reinforcing bar that had 201 been driven into the riverbed (Fig. 3B). The duration of the treatment was 30 min, after which 202 time the zipper was opened with the bag fully submerged and the fish encouraged to swim out of 203 the top of the bag. Since fish were captured and tagged at different sites, the water speed passing 204 through the recovery bag varied among fish. However, the recovery bag treatment occurred only 205 at three tagging sites (Peg Leg, Seabird, Peters), and we consistently used the same location at 206

each site. The mean water speeds where the recovery bags were placed were ~ $0.5 \text{ m} \cdot \text{s}^{-1}$ at Peg Leg, ~ $0.2 \text{ m} \cdot \text{s}^{-1}$ at Seabird, and ~ $0.7 \text{ m} \cdot \text{s}^{-1}$ at Peters.

Cumulatively, we radio-tagged and tracked 182 coho salmon: 50 in 2009, 53 in 2010, and 209 210 79 in 2011. In each year, all coho salmon were tagged until our supply of radio transmitters was exhausted - in total we handled 295 coho salmon. Fish that did not receive transmitters were 211 otherwise processed in the same way, with the primary goal being to collect additional RAMP 212 and entanglement time data. Of the 182 that were tagged, 64 were exposed to the recovery bag 213 treatment – roughly half of the 132 fish we tagged over 2010 and 2011 (no recovery bag 214 treatment in 2009). Most of the data collection in 2010 and 2011 occurred at Peg Leg and thus 215 most of the recovery bag treatments (43 of 64). In 2011, we collected additional data on bycatch 216 condition (e.g., RAMP, FL, entanglement time) from 53 individuals that were not tagged. Also in 217 2011, we opportunistically used a handheld meter (CellOx 325, WTW Inc., Weilheim, Germany) 218 to monitor oxygen depletion over time in beached seines. This was done to estimate the extent to 219 which localized DO depletion can occur in crowded seines during sorting. 220

221

222 *Population identification*

There are five genetically unique and geographically distinct populations of coho salmon within in the interior Fraser River population complex (Beacham et al. 2011; Fig. 1). Adipose fin tissue collected from each individual was used for stream-origin identification via analysis of variation of 17 microsatellite loci (Beacham et al. 2011). Fish were identified to individual spawning streams, but spawning stream fidelity within subpopulations can be low in some years if low flows necessitate finding an alternate spawning site (R. Bailey, DFO, pers. comm.). Thus, for our analyses fish were simply grouped into their populations: Fraser canyon, Fraser middle

drainage, lower Thompson River, North Thompson River, and South Thompson River (Fig. 1,
Table 1). Coho salmon tagged in this study also belonged to three additional populations that
were not from the interior Fraser River watershed: Birkenhead River, Chilliwack River, and the
lower Fraser drainage (Table 1). Identification of population-origin facilitated accurate
determination of whether individuals successfully migrated towards their natal subwatersheds,
beyond terminal radio receivers (see below).

236

237 *Telemetry tracking*

Radio-tagged coho salmon were tracked using an array of radio receiver stations installed 238 at strategic points in the watershed (Fig. 1); methods previously used to assess survival of Fraser 239 River salmon (e.g., English et al. 2005, Cooke et al. 2006, Donaldson et al. 2011, Donaldson et 240 al. 2013, Nguyen et al. 2014). The receiver array differed in each year but key receivers were 241 present at the same locales each year (e.g., Hope, Hell's Gate, Thompson-Fraser confluence). We 242 considered coho salmon as migratory 'survivors' if they were detected at the upstream-most 243 receiver stations en-route to spawning areas. In eight instances (among all 182 fish) there was 244 clear evidence of straying whereby fish migrated to terminal spawning areas to which they were 245 not DNA-identified (e.g., into the South Thompson watershed rather than the North Thompson), 246 in some cases undergoing lengthier upstream migrations than if they had migrated to their natal 247 stream. Those fish were assessed as survivors. Distances from terminal receivers to actual 248 spawning sites varied depending on spawning location ($\sim 5 - 300 + \text{rkm}$; Fig. 1). We also 249 quantified shorter-term, two day survival. Detection efficiency was variable among receivers and 250 years (range = 24 - 100%), but was generally high, particularly at the key receivers (i.e. 251 Thompson-Fraser rivers confluence) and at many of those used as "terminal" points for assigning 252

survivorship. For example, detection efficiency at the Thompson-Fraser confluence was 100% in 253 2009, 96.7% in 2010, and 85.4 % in 2011. Detection efficiency was 100% at Hell's Gate in 2009 254 and 2010, and 56.8% in 2011. Detection efficiencies were assessed by diving the number of fish 255 detected on a receiver by the total number known to have passed that receiver - confirmed either 256 by detection at subsequent receivers, by mobile tracking efforts, or by tag recaptures that were 257 reported. Detection efficiency estimates were therefore inherently less robust for the upstream-258 most receivers and in some cases, unknown for a given terminal receiver in a given year. In 259 general, very little mortality was apparent in upper watersheds (beyond the Thompson-Fraser 260 confluence) such that non-detection at terminal receivers was uncommon for fish that had 261 reached the next most upstream receiver. Among all 182 fish, there were six individuals assigned 262 as mortalities based on non-detection at a terminal receiver whose efficiency was unknown in 263 that year (two in 2010, four 2011), meaning there was some uncertainty associated with final 264 migratory success of those fish. Nevertheless, receivers in upper watersheds typically have good 265 detection efficiency because fish are generally moving through narrower and shallower waters as 266 they approach spawning areas, thus reducing the likelihood of fish using deep water locations 267 that prevent them from being detected. 268

269

270 Data analysis and statistics

To determine what factors affected post-release survival (0 = died, 1 = survived), we used a forced-entry binary logistic regression with six predictor variables: entanglement time (s), total catch size (number of fish), air temperature, FL, use of revival bag treatment (0 = not used, 1 =used), and RAMP score. Air temperature was used in place of river temperature because it varied widely among study days, possibly affecting water temperatures in shallow streamside areas

where entanglement occurred, whereas deeper river temperatures exhibited minimal variation 276 where temperature monitoring stations were located. Differences in survival among populations, 277 years, and capture locales were assessed for significance using Pearson's chi-square test. To look 278 279 for associations between RAMP scores, handling time, and total catch, we used Spearman rankorder correlations. Confidence intervals for mortality rates were calculated to provide a measure 280 of sampling error and took the form of binomial confidence intervals, based simply on the 281 number of mortalities and the total number of fish (using the Clopper-Pearson exact method, 282 similar to Stokesbury et al. 2011). All statistical tests were conducted using R (v. 3.0.0). Tests 283 were assessed as significant at $\alpha = 0.05$. 284

285

286 Human dimensions surveys

In 2011, we conducted face-to-face interviews with aboriginal fishers and members 287 involved with the aboriginal fishing process, which included crew monitors, fish buyers, and 288 buying employees (N = 111). We included the latter groups (< 10 % of interviewees) because 289 they were also aboriginal and most had directly participated in this harvest fishery in the past as 290 fishing crew members or had extensive exposure to beach seining through their role in the 291 fishery. Moreover, buyers and monitors are highly involved with the fishing process, are 292 knowledgeable about fishing methods and locations, and can potentially influence fisheries 293 policy. Participants were chosen opportunistically due to logistical limitations (i.e., access to 294 fishing sites), and timing of when fishing crews were on breaks from fishing. We aimed to 295 interview at least 50% of the members of each crew, including the crew chief, and to interview 296 all crews present at a given site. We interviewed members of $\sim 30\%$ of the fishing crews who 297 298 participated in the beach seine fishery in 2011 (Karen Burnett, DFO, pers. comm.). Beach seine

crews consisted of 11 individuals on average (range = 8 - 13). Fewer than five percent of those 299 approached declined to take part in the interview. Seventy-two (72) of 111 interviews were 300 conducted streamside during the aboriginal economic opportunity beach seine fishery in the 301 302 lower Fraser River. The remaining 39 interviews were opportunistically conducted during an aboriginal economic opportunity gillnet fishery (targeting sockeye salmon) that occurred a 303 month earlier (on 24/08/11 and 25/08/11) in the lower Fraser River watershed. Because questions 304 #1-4 (see below) only pertained to the beach seine fishery, respondents in the gillnet fishery were 305 only asked those questions if they stated that they also take part in the beach seine fishery, or 306 have done so in the past (28 of 39). 307

The interviews were semi-structured, meaning that aboriginal fishing members could 308 identify any additional topics, concerns, or ideas relating to coho bycatch during the interview. 309 The interviews were used to address a number of research objectives, some of which are beyond 310 the scope of this paper (see Nguyen et al. 2012). We used two separate interviews: one shorter 311 questionnaire and a long-form questionnaire. The short questionnaires were a subset of the long 312 questionnaire, and included the same closed-ended questions (data not presented here) while 313 open-ended questions alternated among the following themes: participant perspectives on coho 314 by by catch, perspectives about biotelemetry science, and thoughts on the fish recovery bag. The 315 short questionnaire was intended to be for individuals that only had 5-10 minutes to spare, while 316 effort was made to ensure all crew leaders participated in the long questionnaire. For individual 317 questions, the number of responses was somewhat lower than the total number of interviewees 318 (111, see Fig. 5, 6, 7) because alternating versions of the shorter questionnaire contained only a 319 subset of the questions, or because a question was not applicable to the interviewee (e.g., if they 320 321 were a fish buyer). We asked the following questions:

322	1. Do you think that beach seine capture has any effect on a coho's chance of successful						
323	migration or spawning?						
324	2. What do you think causes the greatest stress for coho released from beach seines?						
325	3. What suggestions do you have to increase the survival of coho bycatch?						
326	4. If you knew that leaving the seine in knee-high water for sorting would increase the						
327	chance of survival for released fish, would you voluntarily do it?						
328	The interviewees were then shown a fish recovery bag (those tested in this study), accompanied						
329	by an explanation of its purpose and use. They were then asked the following questions:						
330	5. What are your thoughts on the recovery bag?						
331	6. If research data show the recovery bag improves post-release survival for salmon, how						
332	likely would you be to use one on a voluntary basis?						
333	7. Do you think the recovery methods and gear should be mandatory for reviving coho						
334	bycatch from seine capture?						
335	All survey responses were coded according to emergent themes following standard						
336	qualitative protocol (Strauss 1987, Creswell 2009). We assessed for consistencies among codes						
337	(similar meanings or pointing to a basic idea), and revealed themes and categories reflecting						
338	fisher responses to the particular questions (Fig. 5, 6, and 7). For example, clear "yes" or "no"						
339	answers were often provided, but responses were in some cases classified as "no-conditional"						
340	when responses were "no" with conditions or dependent on certain situations (e.g., coho salmon						
341	are not affected by capture <i>if</i> they are handled carefully). For further information on how						
342	responses were categorized and details of their meaning, see the online supplementary						
343	information associated with this article. We did not conduct any statistical tests on resultant						

survey responses, noting the paper by Drury et al. (2011) that recommends the use of qualitativeapproaches to using social science data in conservation research.

346

347 RESULTS

Collectively, we handled 295 wild coho salmon, bycatch that was spread across 98 net 348 sets and three study years. In those net sets, collectively, an estimated 54,790 adult salmon were 349 caught, meaning that the coho salmon by catch for those sets was $\sim 0.5\%$ of total catch. The real 350 bycatch rate is somewhat lower, owing to the fact that we did not record any data for sets in 351 which no bycatch occurred ($\sim 20-40\%$ of all sets). The median total catch (number of adult 352 salmon) for net sets from which coho salmon were tagged was estimated as 500 (range: 70 -353 2000). All but eight of the coho salmon we tagged and tracked were DNA-identified as 354 belonging to the interior Fraser River population complex (Table 1; note that no fish were 355 excluded from the mortality estimate on the basis of population-origin). 356 Among all the coho salmon we handled from 2009-2011, the mean fish size (FL) was 357 61.6 cm (median: 62 cm, range: 37 - 82 cm). Median entanglement in the beached seine was 3 358 min 20 s (mean: 6 min 32 s, range: 5 s - 55 min 58 s). The most frequently observed RAMP 359 score was 0.4, an impairment level that was characterized 97% of the time by loss of the tail grab 360 and body flex reflexes. There was an apparent immediate effect of the recovery bag on fish 361 vitality: after 30 min in the recovery bag treatment, all fish were highly vigorous (i.e., RAMP 362 score of zero). 363

364

365 Bycatch mortality

Of the 182 coho salmon we tagged, 65 (36%) failed to migrate past terminal radio 366 receivers en route to spawning areas, and were thus classified as mortalities (Table 2). Adjusting 367 our sampling effort (N = 182) to a binomial distribution provides a 95% confidence interval 368 369 (C.I.) for post-release mortality of 29-43% (Stokesbury et al. 2011). Nine (9) of the 295 coho salmon (3%) caught during the study were dead upon being pulled from the net (Table 2). The 370 cumulative mortality rate was 37.3% (95% C.I. = 32 - 43% mortality). Using the probability of 371 mortality associated with each RAMP score to predict migration success of the non-tagged fish 372 that were released alive (rather than applying a blanket post-release mortality rate to those fish 373 whose fates were unknown), the extrapolated total mortality estimate was 39.3% (Table 2; 95% 374 C.I. = 34 - 45% mortality). There were no significant differences in post-release mortality among 375 the four main capture locations ($\chi^2_{3,181} = 1.71$, P = 0.63) or among years ($\chi^2_{2,181} = 1.29$, P =376 0.53). Estimated population-specific differences in mortality were not significant (Table 1; Chi-377 square test: $\chi^2 = 1.39$, df = 5, P = 0.99). Among all the variables we tested as predictors of 378 mortality, none were significant except for RAMP score, whereby fish with higher RAMP scores 379 (more impaired) were less likely to be successful migrants (Table 3, Fig. 4). Use of a revival bag 380 did not influence post-release mortality (P > 0.05, Table 3). 381

Since some in-river mortality is natural, there is a need to attempt to differentiate mortality caused by the capture itself. To do so, RAMP scores can be used whereby coho salmon released with little or no reflex impairment (vigorous) are assumed to experience no post-release bycatch mortality. Using that conservative assumption, the post-release mortality rate for those fish can then be used as a baseline within the dataset. Additional mortality above that baseline that occurs at higher levels of reflex impairment can then be assigned to the fishery (see Fig. 4). The 'baseline' post-release mortality rate at zero reflex impairment was 23% (Fig. 4). Applying a

net mortality rate above that baseline for higher RAMP scores to the total number of fish at those 389 RAMP scores yields a new estimated proportion of fish that died as a result of the capture stress 390 -13.6% overall of those released alive (Fig. 4). Combined with immediate mortality, this 391 conservative approach accrues a bycatch mortality rate of 16.6% (95% C.I. of 13 - 21%). 392 Entanglement time and total size of the catch were not predictive of survival, but they 393 were associated with reflex impairment (i.e., RAMP score - itself predictive of survival). There 394 was a small but significant positive correlation between entanglement time and RAMP score (rs 395 = 0.18, P = 0.03). Total catch was not significantly correlated with RAMP score (r_s = 0.15, P =396 0.06), but total catch and entanglement time were strongly correlated (Pearson's R = 0.64, P < 0.06) 397 0.001). 398

399

400 Human dimensions surveys

The majority of the participants of the beach seine fishery we surveyed stated that they 401 thought beach seine capture does not or is not likely to affect a coho salmon's subsequent 402 migration or spawning success (no effect -41%, no effect under most conditions -21%; Fig. 403 5A; note – total numbers of respondents provided in Figure). As such, only 12% of participants' 404 responses were codified as a "yes", with a further 12% as a "conditional yes" (e.g., yes - stressful 405 enough to reduce survival under certain scenarios). In a following question, many respondents 406 identified crowding (23%), handling (19%), and air exposure (9%) as particularly stressful 407 components of the capture experience (Fig. 5B). Fishers were then asked if they had any 408 suggestions about how to increase the survival of coho salmon caught in their nets, and the most 409 common response (26%) was to release fish quickly (Fig. 6A). Cumulatively, however, more 410 411 respondents (35%) stated that nothing could be done (21%) or did not have an approach to

412 propose (14%; Fig. 6A). When asked whether they would be likely to voluntarily leave the seine 413 in knee-deep water for sorting (if it were shown to increase survival), 51% responded in the 414 affirmative, while a further 26% stated that they already use this practice (Fig. 6B). Nine percent 415 expressed concern that leaving the net in deeper water could compromise the safety of their 416 crew.

We received positive feedback to the concept of the fish recovery bag. Forty-seven (47) 417 percent of respondents (44 of 94) gave fully positive responses when briefed about its design and 418 purpose and asked for their thoughts, with a further 17% providing conditionally positive 419 responses (e.g., it would have to be shown to improve survival; Fig. 7A). Many fishers thought 420 the use of a revival bag should be mandatory, without the qualifier that evidence would support 421 its use (44%; Fig. 7B). However, responses were more positive to a proposal that recovery bag 422 use be voluntary rather than mandatory, given scientific data to support use of recovery bags 423 (Fig. 7C). A number of the conditionally positive responses arose from uncertainty about the 424 monetary cost of the bags and whether they would be provided or subsidized by management. 425

426

427 DISCUSSION

We have provided an estimate of bycatch mortality for an endangered population of coho salmon captured in an aboriginal beach seine fishery, based on three years of tracking fish released from the fishery. In addition, we have evaluated whether altered capture and handling techniques (Fig. 2) or the use of a revival bag (Fig. 3) could reduce bycatch mortality. The inclusion of human dimensions data provides insight into the perspectives of the fishers on the impacts of, and solutions to, coho salmon bycatch in their fishery. This combined approach allows us to make recommendations to management that are informed by the resource users, who

themselves can directly affect successful implementation of management strategies (e.g., rapidrelease of bycatch).

437

438 *Bycatch mortality*

The total observed mortality of 39%, inclusive of natural mortality, is relatively low 439 when compared against similar studies on post-release survival for Fraser River salmon. For 440 example, sockeve salmon caught by anglers experienced ~65-70% total mortality from release in 441 the lower river to reach terminal radio receivers, while those released in parallel from a beach 442 seine had a mortality rate of ~43-48% (Donaldson et al. 2011, 2013). Gillnets typically have 443 higher post-release mortality – up to 70% within just 24-48 h based on net pen holding studies 444 (in the marine environment; Buchanan et al. 2002); a 60% mortality rate is applied to gillnet 445 bycatch by management to account for bycatch mortality in attempts to meet spawning 446 abundance targets (DFO 2011). Thus, there appears to be consistent evidence that beach seine 447 capture results in the lowest post-release mortality for Fraser River salmon, particularly when 448 using the more holistic approach of examining long-term mortality. 449

Not all observed mortality (39%) can be attributed to beach seine capture as there is a 450 component of natural en-route mortality which we have no rigorous way of distinguishing. To 451 account for impacts on numbers of salmon reaching spawning areas (termed escapement), DFO 452 typically apply mortality rates to bycatch that are based on 24 or 48 h post-capture net pen 453 holding studies. Net pen studies can underestimate mortality because of possible synergies 454 between capture stress and predation risk (Rogers et al. 2014), but do have the advantage that 455 very little of any observed mortality is likely to be natural because of the small time window. 456 457 Immediate and short-term (48 h) mortality (combined) was estimated to be 18.6% in the present

study (95% C.I. of 14 - 24%). An alternate approach to calculating a bycatch mortality rate that 458 attempts to distinguish bycatch from natural mortality, is to use RAMP scores and their mortality 459 rates at each level of impairment, and assume negligible bycatch mortality for the fish that were 460 least impacted (vigorous at release). That approach (Fig. 4) yields a bycatch mortality rate of 461 17% (rounded from 16.6%). The remaining mortality (39% - 17% = 22%) could in such a case 462 partly be attributed to a combination of natural migration failure, unreported fisheries removals, 463 and the unknown long-term effects of the transmitter. Although we have no true control for the 464 mortality we observed, this approach (Fig. 4) uses low RAMP score fish within our dataset as 465 comparative controls. The 'control' mortality rate within the dataset was 23%, from release to 466 upper watersheds. For coho salmon, high uncertainty exists about natural in-river mortality, but 467 available data for Thompson River sockeye salmon stocks with similar migration timing show 468 that natural mortality ranges from 8-20% from Mission (Fig. 1) to spawning areas (English et al. 469 2005; Martins et al. 2011). 470

True control mortality rates are often unknown in post-release mortality studies; every 471 method of estimating mortality has limitations. Short-term captivity is the most common method 472 (e.g., using a net pen) of estimating post-release mortality (Rogers et al. 2014), but this method 473 fails to expose fish to the challenges of upstream migration and imposes severe chronic 474 confinement stress (cortisol elevation) in wild Pacific salmon (Donaldson et al. 2011). The best 475 alternative to captivity is to use biotelemetry (Donaldson et al. 2008), but most biotelemetry 476 applications involve intracoelomic implantation via surgery or external attachment of a 477 transmitter, with the latter typically requiring piercing the dorsal musculature to attach a tag that 478 creates external drag (e.g., 'backpack' attachment – see Colotelo et al. 2012). The gastric tagging 479 480 method we used is among the most rapid and least injurious tagging methods available in

fisheries science: it involved a < 1% tag burden (by weight), a rapid < 10 s procedure with no 481 injury or anesthetic, and created minimal hydrodynamic drag. This tagging procedure has been 482 widely used in Fraser River salmon (Cooke et al. 2008; Martins et al. 2011) and, to some extent, 483 validated as benign (Cooke et al. 2005; English et al. 2005; Wilson et al. 2013). The other 484 components of our handling procedure were similarly rapid - the length measurement, DNA 485 biopsy, and RAMP assessment each required < 20 s to complete and instances of air exposure 486 (e.g., for assessing certain reflexes) were very brief (~ 5 s). Although blood biopsy has been 487 commonly included in past biotelemetry studies on the Fraser River as a means of assessing 488 stress, exhaustion, and identifying sex (e.g., Cooke et al. 2005), we avoided using it here to allay 489 concerns of managers that it could affect mortality, despite evidence to the contrary (Cooke et al. 490 2005; G.D. Raby, unpublished data). Nevertheless, it would be near-impossible to empirically 491 test the independent or synergistic effects of different components of researcher handling on 492 delayed mortality in the context of a real fishery. Although the length of research handling time 493 was no different between fish that died and those that migrated successfully (Raby et al. 2012), 494 the potential for our handling and tagging to add to mortality remains an area of some 495 uncertainty. It is for that reason that we offer a conservative approach for assigning mortality to 496 the fishery for management purposes (Fig. 4). 497

As with any mortality estimate, uncontrolled factors undoubtedly affect its accuracy and it is impossible to quantify this error. Reporting rates for radio-tag recaptures in the Fraser River are quite good in the recreational angling community, but the First Nations fishery has limited knowledge of the tag return-and-reward program in spite of its existence for ~ 10 years (Nguyen et al. 2012). In some cases, there were multiple beach seine crews directly upstream of where we were releasing tagged coho salmon, increasing the likelihood of recapture. Our interactions with

many crews during the fishery will likely have increased awareness, but there may nevertheless 504 have been some incidents of fish being recaptured and released after removal of the radio 505 transmitter and this would have mainly had the effect of increasing our 48 h mortality rate. Some 506 507 fish may have died because of capture despite being vigorous at release (e.g., at RAMP scores of 0 or 0.2 – see Fig. 4), resulting in an imperfect relationship between RAMP and mortality. 508 Assuming that is the case, the baseline mortality rate we used (23%) in calculating our 'alternate' 509 mortality rate of 17% would be an overestimate, with the true mortality caused by the fishery 510 thus being higher than 17%. In addition, our receiver array did not extend all the way to 511 spawning grounds and it was not possible to verify spawning success. Latent capture-mediated 512 disease-induced mortality likely occurs in fish (Lupes et al. 2006), though little is known about 513 the potential for this to occur in any real fishery. In sockeye salmon, gillnet injuries can cause 514 latent spawning failure for fish that reach spawning grounds because of their effect on 515 reproductive physiology (Baker and Schindler 2009, Baker et al. 2013). Most importantly, 516 observer effects, a well-known limitation to bycatch research (Benoit and Allard 2009), may 517 have been a factor. All crews we worked with were clearly aware of our desire to radio-tag coho 518 salmon, and spent a concerted 1-3 min attempting to locate bycatch upon landing of the net, prior 519 to focusing on sorting their target catch. If such efforts are not typically made, entanglement 520 times would increase resulting in higher reflex impairment and thus higher mortality. 521

522

523 *Possible solutions to mortality*

Among all the variables we measured at the time of capture, the only one that was statistically linked to post-release mortality by our analyses was RAMP score – our multi-year dataset supports a previous single-year study (Raby et al. 2012). The finding that variables other

than reflex impairment were not associated with survival was surprising. This discrepancy may 527 be largely explained by some of the mortality occurring as a result of factors unrelated to capture 528 (e.g., natural mortality or tagging effects). It also suggests that the severity of capture stress 529 530 experienced by an individual fish was influenced by aspects of capture that were not measured. Interestingly, while entanglement time did not show a direct statistical association with post-531 release mortality, it was correlated with RAMP scores, which have previously been shown to 532 predict survival in Fraser River coho salmon (Raby et al. 2012). Although this is not a strong 533 statistical relationship, we can advise that any capture or handling techniques that result in lower 534 RAMP scores and more vigorous fish (e.g., reducing entanglement time) would be beneficial to 535 minimizing post-release mortality. 536

The full suite of factors that led to reflex impairment may have not been identified by our 537 study because a fish's RAMP score is likely to be reflective of its unique capture experience, 538 such as its location within a net or the techniques used by the crew that catches it. We did not 539 attempt to quantify such variables and it would be very difficult to do so in an active fishery. We 540 handled bycatch from 12 separate fishing crews during the study, and there were variations in 541 how their nets were pulled into shore, how fish were crowded and sorted (Fig. 2), and slight gear 542 differences (net material and mesh size). Some of the differences likely depended on variation in 543 the local riverbank morphology (slope and substrate), the strength of the river current, the 544 training and coordination of the crew, whether the crew used a truck to pull in their net, and the 545 size of the catch in a given set. Very large hauls (> 1000 fish) would almost always necessitate 546 leaving the net in deeper water for sorting, thus suggesting that bycatch at least remained 547 submerged and supplied with oxygen until it was located and released (Fig. 2B). However, 548 overcrowding resulted in dissolved oxygen (DO) depletion to the extent that DO saturation 549

decreased to 56-60% within 10 minutes of the start of sorting in one large set we measured (Fig. 550 2B). Approximately 30 minutes into the sorting process for that set, DO had further decreased to 551 50% saturation at the upstream end of the net, and to 36-40% at the downstream end. The effect 552 553 of those changes in dissolved oxygen were visibly evident: the target species (pink salmon) at the upstream end of the net were alive, upright, and facing upstream, while at the downstream end 554 all the catch appeared to be dead or moribund. In smaller more typical net sets (Fig. 2A), the 555 catch was brought into a similar water depth but more easily spread out along the shoreline 556 resulting in less crowding, although the fish were often in water shallow enough that half their 557 gills were air exposed. In those sets it was easier to quickly identify and release coho salmon. In 558 two such sets we monitored, oxygen only descended to \sim 75-80% saturation during the < 20 min 559 required to complete sorting, while in a third net set (e.g., Fig. 2A), DO saturation decreased to 560 42% 25 min into the sorting process (by which time all bycatch had been released). There were 561 instances of crews pulling their entire catch completely onto the beach for sorting, air exposing 562 both the target catch and bycatch (Fig. 2C). Though resulting in air exposure, this method always 563 ensured coho salmon were identified and released very quickly, typically after 1-2 min of air 564 exposure. Pulling the entire net onto shore was only physically possible with smaller sets of \sim 565 100-300 salmon, and often necessitated by a steep riverbed and strong current that made leaving 566 the net in deeper water for sorting unsafe for the fishers (though this was not the case in Fig. 2C). 567 We found no evidence that the fish revival bag tested in this study benefitted post-release 568 survival. Coho salmon provided with a 30-min recovery period in the flow-through fish bag (Fig. 569 3B) prior to release were no more likely to migrate successfully than fish immediately released 570 after tagging. While fish were clearly invigorated by receiving 30-min of ram ventilation (i.e., 571 forced water flow over the gills), the bag seemed to offer no added survival benefit for fish 572

beyond that provided by a free-swimming recovery in the river. Physiological evidence has 573 shown that a free-swimming recovery is the optimal way for salmonids to return to homeostasis 574 following exhaustive exercise or fisheries capture (Milligan et al. 2000, Farrell et al. 2001). 575 576 Thus, we suspect that facilitating revival using recovery bags only benefits those fish that are severely impaired (i.e., unable to swim), whereas it likely represents added chronic confinement 577 stress for fish otherwise able to maintain equilibrium during the recovery period. Indeed, in the 578 present study there was a small (non-significant) increase in survival associated with the 579 recovery bag for fish with a negative orientation (equilibrium) reflex (47% survival with 580 recovery bag treatment, 35% without, N = 17 fish per group). In general, facilitated revival 581 techniques may be most useful in contexts where post-release predation occurs (Brownscombe et 582 al. 2013, Raby et al. in press); in the lower Fraser River where water turbidity is high there is 583 little evidence that this is a major issue, as compared with the marine environment where revival 584 techniques are currently in use for coho salmon (Farrell et al. 2001). It is possible that a shorter 585 revival duration would have been more beneficial (e.g., 10-15 min) by allowing enough time to 586 regain basic vitality without extended confinement stress. Other research has shown that various 587 means of facilitating post-capture revival (including recovery bags) can be effective but that 588 attempts at revival can also be misguided in some contexts (Farrell et al. 2001, Brownscombe et 589 al. 2013, Donaldson et al. 2013, Robinson et al. 2013, Nguyen et al. 2014). Future research on 590 facilitating revival of fish bycatch should take the form of well-controlled experiments that 591 examine what recovery durations and techniques are optimal and at what level of impairment 592 fish benefit from revival. 593

594

595 *Perspectives of the fishery participants*

The most notable pattern in the results of the survey was the general willingness to 596 engage in strategies to mitigate bycatch mortality. For example, the majority of aboriginal fishers 597 were receptive to using the fish recovery bags (Fig. 7). The recreational angling community in 598 599 the lower Fraser River were asked similar questions regarding fish recovery bags (Donaldson et al. 2013) and were less receptive (> 40% negative responses) when asked, "What do you think of 600 the idea of a revival bag?". Our biotelemetry data suggest that recovery bags may have limited 601 potential for improving survival in the beach seine fishery. Nonetheless, responses on the 602 recovery bag questions suggest there is a general likelihood that members of the fishery would 603 embrace strategies that benefit coho salmon they catch. Though a plurality of respondents stated 604 that beach seine capture does not affect survival of coho salmon, which conflicts somewhat with 605 our findings, many were able to identify causes of stress and suggest ways to improve survival 606 (Fig. 5 and 6). Even more encouragingly, there was a clear willingness to alter handling practices 607 to improve the condition of bycatch, such as by leaving the net in deeper water for sorting to 608 ensure adequate oxygen supply to bycatch until it is found. Collectively, the human dimensions 609 610 data painted a picture of a fishing community that would be receptive to advice about how to best handle coho salmon to maximize their survival if the advice is supported with scientific 611 evidence. 612

613

614

14 Integration – management recommendations

In the introductory text of DFO's selective fishing policy (DFO 2001), Thompson River coho salmon were cited as the lead example of a fish population whose restoration was being slowed because of bycatch in fisheries targeting other more abundant Pacific salmon. Since the shift towards a selective fishing policy began in Canada's Pacific fisheries (DFO 2001),

managers have applied a 5% mortality rate to coho salmon caught in the aboriginal beach seine 619 fishery for the purpose of ensuring that overall bycatch mortality (all fisheries combined) is 620 limited to a level that does not significantly compromise numbers of spawning adults. The 621 622 bycatch mortality estimate in this paper is higher than the 5% currently applied. Based on our data, we recommend fisheries management consider the following options for updating the 623 bycatch mortality rate used to manage this fishery: a minimum of 17% bycatch mortality (Fig. 624 4); a value based on 48 h mortality (18.6%), or an estimate more reflective of total mortality 625 (39%; Table 2). More work is needed to articulate the benefits and disadvantages of each 626 approach. Unfortunately, comparable multi-year studies have not been done on all the other 627 fisheries that bycatch interior Fraser coho salmon. Of the little work that has been done, beach 628 seines have consistently resulted in the highest post-release survival relative to the other Fraser 629 River fisheries that have been evaluated (i.e., higher than angling or gillnets; Buchanan et al. 630 2002, Donaldson et al. 2011, Donaldson et al. 2012, Donaldson et al. 2013, Nguyen et al. 2014), 631 which suits the management shift towards a selective fishing policy (DFO 2001). In fact, 632 management and stakeholders can use this information to contemplate the possibility of an even 633 greater shift towards gear types that minimize release mortality. 634

Beyond simply documenting the extent of the problem, we have attempted to identify solutions to potential causes of mortality. At this time we are unable to recommend use of fish recovery bags in the beach seine fishery. Encouragingly though, many of the participants of the fishery were enthusiastic about the idea. Those responses suggest that fishers would similarly be enthusiastic about other means of improving survival of bycatch, perhaps including reducing entanglement time. Alternatively, fish bags could be used on a voluntary basis with a precaution that they only be used with severely impaired coho salmon and whereby the fish only remains in

the bag until reflex actions are regained. The expanded validation of the RAMP approach in the present study provides confirmation that this simple technique is ready for use in this fishery if needed (Raby et al. 2012). The observers in the fishery could easily be taught how to conduct RAMP assessments to monitor the condition of bycatch in real time, provide advice to their crews on how to improve fish condition, and make decisions about whether individual fish should be revived using recovery bags.

The key variable that was significantly associated with reflex impairment was 648 entanglement time. Entanglement time was also positively correlated with total catch, with 649 by catch often being difficult to locate in larger sets, resulting in a wider range of entanglement 650 times for those sets. To minimize reflex impairment and maximize survival, the fishers should be 651 encouraged to: a) make all efforts to release coho salmon to the river within 5 min of pulling the 652 net into shore, and b) particularly if the catch is large (e.g., > 500 fish), minimize crowding by 653 leaving the net in deeper water during sorting so that bycatch are adequately supplied with 654 oxygen until they can be located. In our surveys, we learned that many of the fishers understand 655 the benefits of such practices, are willing to use them, and that some already do so. However, 656 there is clear potential for a) an increase in fisher awareness of best bycatch handling practices 657 and b) reductions in reflex impairment. Most of the participants in the fishery have intuition 658 based on a lifetime of experience capturing and handling salmon (i.e., traditional ecological 659 knowledge; Huntington 2000) that could enable them to develop their own techniques that 660 reduce reflex impairment. With observer-based collection of RAMP data, fisheries management 661 would have the option to develop incentives that motivate the fishers to minimize RAMP scores 662 (and thus mortality; a general approach used with success in tuna fisheries; Hall et al. 2000). 663

664

665 *Synthesis*

This paper demonstrates that fisheries science, biotelemetry, and human dimensions 666 surveys can be combined to evaluate a conservation problem for an endangered population of 667 salmon and inform resource managers and users. We consider this a model approach for 668 conservation research, because it can help address the persistent challenge of generating science 669 that "bridges the knowledge-action boundary" (Cook et al. 2013). A well-known barrier to 670 transitioning from scientific knowledge to conservation action is the scientific structure that 671 values publications and grant income but not engagement with stakeholders (Cook et al. 2013). 672 As part of broader stakeholder meetings regularly held to present findings from several projects 673 by our research group (see Cooke et al. 2012), we disseminated the findings of this study 674 throughout the course of data collection and received input for following phases of research. 675 676 Collaboration with stakeholders was done to increase the likelihood that findings are subsequently adapted. In fact, this study was initiated by conversations with fisheries managers 677 that followed a 2009 stakeholder meeting and was only possible with various forms of support 678 from the resource users, government, and environmental NGOs (among others). Based on the 679 experience with this study, adding a human dimensions component to conservation research 680 represents good value considering the modest time investment required (e.g., conducting the 681 surveys required ~ 8 days for two researchers). Previous research has shown the utility of such 682 an approach for Fraser River sockeye salmon recreational fisheries (Donaldson et al. 2013). 683 Other papers have shown the value of combining sociology and natural sciences to address 684 animal conservation issues (African elephants - Loxodonta africana; Guerbois et al. 2012; red 685 deer - Irvine et al. 2009). In the conservation of tropical forests subject to logging, a conservation 686 687 context somewhat analogous to harvest fisheries, this approach has been applied numerous times

(Lele and Kurien 2011 and references therein). Fisheries bycatch is recognized widely as a leading global threat to biodiversity (Gray 1997, Kappel 2005, Davies et al. 2009), towards which a great deal of ecological knowledge and research have been applied (Hall et al. 2000, Soykan et al. 2008). Our effort is among the first integrative research papers in the realm of fisheries bycatch, and certainly the first for a freshwater bycatch issue (Raby et al. 2011). In addition to helping address a conservation issue for an endangered population of salmon, we hope this paper can have value for future interdisciplinary research in aquatic conservation.

695

696

697 ACKNOWLEDGMENTS

All fish handling and tagging protocols were approved by the Carleton University Animal 698 699 Care committee and the University of British Columbia Animal Care committee. The human dimensions survey protocol was approved by the Ethics Committee at Carleton University. We 700 are grateful to the many First Nations fishing crews of the lower Fraser River that allowed us to 701 702 work with them – especially the fishing crew led by Lester Mussel, and the crews from the Peters band and the Seabird Island band. Marla Maxwell, Rob Tadey, Sheldon Evers, Jason Mahoney, 703 Karen Burnett, and Matthew Parslow from DFO facilitated access to the fishery. Funding for this 704 project was provided through the Natural Sciences and Engineering Research Council of Canada 705 (NSERC) via a Strategic Grant (to S.J.C. and S.G.H.) and by the Canadian Wildlife Federation 706 Endangered Species Research Program. Further support was provided by Carleton University, 707 the University of British Columbia, the Canada Foundation for Innovation, the Canada Research 708 Chairs Program (S.J.C.) and Ocean Tracking Network Canada. Lisa Thompson from the DFO 709 710 Environmental Watch program provided river temperature data. Richard Bailey from DFO stock

711	assessment provided helpful information on the biology and stock assessment of interior Fraser					
712	coho salmon populations. John Candy at the DFO Pacific Biological Station conducted DNA					
713	analyses for stock identification. Samantha Wilson, Nolan Bett, Marion Dupeux, Sarah					
714	Larocque, David Roscoe, Andrew Lotto, S. Matthew Drenner, Sarah McConnachie, and					
715	Charlotte Whitney provided assistance in the field. Sean Tyerman of LGL Ltd. installed and					
716	maintained the telemetry array. Anonymous referees provided helpful input that improved the					
717	manuscript. G.D.R., M.R.D., V.M.N., and N.M.S. were supported by NSERC scholarships.					
718						
719	LITERATURE CITED					
720	Beacham, T. D., M. Wetklo, L. Deng, and C. MacConnachie. 2011. Coho salmon population					
721	structure in North America determined from microsatellites. Transactions of the American					
722	Fisheries Society 140:253-270.					
723	Benoit, H. P., and J. Allard. 2009. Can the data from at-sea observer surveys be used to make					
724	general inferences about catch composition and discards? Canadian Journal of Fisheries and					
725	Aquatic Sciences 66:2025-2039.					
726	Bradford, M. J., and J. R. Irvine. 2000. Land use, fishing, climate change, and the decline of					
727	Thompson River, British Columbia, coho salmon. Canadian Journal of Fisheries and Aquatic					
728	Sciences 57:13-16.					
729	Brownscombe, J. W., J. D. Thiem, C. Hatry, F. Cull, C. R. Haak, A. J. Danylchuk, and S. J.					
730	Cooke. 2013. Recovery bags reduce post-release impairments in locomotory activity and					
731	behaviour of bonefish (Albula vulpes) following exposure to angling-related stressors.					
732	Journal of Experimental Marine Biology and Ecology 440:207-215.					

- Buchanan, S., A. P. Farrell, J. Fraser, P. Gallaugher, R. Joy, and R. Routledge. 2002. Reducing
- 734 gill-net mortality of incidentally caught coho salmon. North American Journal of Fisheries

735 Management 22:1270-1275.

- Campbell, L. M. 2005. Overcoming obstacles to interdisciplinary research. Conservation biology19:574-577.
- 738 Cook, C. N., M. B. Mascia, M. W. Schwartz, H. P. Possingham, and R. A. Fuller. 2013.
- Achieving conservation science that bridges the knowledge-action boundary. ConservationBiology, in press.
- 741 Cooke, S. J., G. T. Crossin, D. A. Patterson, K. K. English, S. G. Hinch, J. L. Young, R.
- Alexander, M. C. Healey, G. van der Kraak, and A. P. Farrell. 2005. Coupling non-invasive
- 743 physiological and energetic assessments with telemetry to understand inter-individual
- variation in behaviour and survivorship of sockeye salmon: development and validation of a
- technique. Journal of Fish Biology 67:1342-1358.
- Cooke, S. J., S. G. Hinch, G. T. Crossin, D. A. Patterson, K. K. English, M. C. Healey, J. M.
- 747 Shrimpton, G. Van Der Kraak, and A. P. Farrell. 2006. Mechanistic basis of individual
- mortality in Pacific salmon during spawning migrations. Ecology 87:1575-1586.
- 749 Cooke, S. J., et al. 2008. Developing a mechanistic understanding of fish migrations by linking
- telemetry with physiology, behaviour, genomics and experimental biology: an
- interdisciplinary case study on adult Fraser River sockeye salmon. Fisheries 33:321-338.
- 752 Cooke, S. J., S. G. Hinch, M. R. Donaldson, T. D. Clark, E. J. Eliason, G. T. Crossin, G. D.
- Raby, K. M. Jeffries, M. Lapointe, K. Miller, D. A. Patterson, and A. P. Farrell. 2012.
- Conservation physiology in practice: how physiological knowledge has improved our ability

to sustainably manage Pacific salmon during up-river migration. Philosophical Transactions

756 of the Royal Society B 367:1757-1769.

- 757 Cooke, S. J., V. Messmer, A. J. Tobin, M. S. Pratchett, and T. D. Clark. 2014. Refuge-seeking
- impairments mirror metabolic recovery following fisheries-related stressors in the Spanish
- flag snapper (*Lutjanus carponotatus*) on the Great Barrier Reef. Physiological and
- Biochemical Zoology, 87:136-147.
- 761 Creswell, J. W. 2009. Research Design: Qualitative, Quantitative, and Mixed Methods

762 *Approaches*. 3rd ed. SAGE Publications Inc., Thousand Oaks, CA.

- 763 DFO (Fisheries and Oceans Canada) 2001. A policy for selective fishing in Canada's Pacific
- fisheries. Available online at: http://www.dfo-mpo.gc.ca/Library/252358.pdf.
- 765 DFO (Fisheries and Oceans Canada) 2005. Canada's policy for the conservation of wild salmon.
- Available online at: http://www.pac.dfo-mpo.gc.ca/publications/pdfs/wsp-eng.pdf.
- 767 DFO (Fisheries and Oceans Canada) 2011. Pacific region integrated fisheries management plan:
- salmon, southern B.C, June 1, 2011 to May 31, 2012. Available online at: http://www.dfo-
- 769 mpo.gc.ca/Library/343942.pdf.
- Davies, R. W. D., S. J. Cripps, A. Nickson, and G. Portier. 2009. Defining and estimating global
 marine fisheries bycatch. Marine Policy 33:661–672.
- Davis, M. W. 2010. Fish stress and mortality can predicted using reflex impairment. Fish and
 Fisheries 11:1-11.
- Donaldson, M. R., T. D. Clark, S. G. Hinch, S. J. Cooke, D. A. Patterson, M. K. Gale, and A. P.
- Farrell. 2010. Physiological responses of free-swimming adult coho salmon to simulated
- predator and fisheries encounters. Physiological and Biochemical Zoology 83:973-983.

777	Donaldson, M. R., S. G. Hinch, D. A. Patterson, J. Hills, J. O. Thomas, S. J. Cooke, G. D. Raby,						
778	L. A. Thompson, D. Robichaud, K. K. English, and A. P. Farrell. 2011. The consequences of						
779	angling, beach seining, and confinement on the physiology, post-release behaviour and						
780	survival of adult sockeye salmon during upriver migration. Fisheries Research 108:133-141.						
781	Donaldson, M. R., S. G. Hinch, G. D. Raby, D. A. Patterson, A. P. Farrell, and S. J. Cooke.						
782	2012. Population-specific consequences of fisheries-related stressors on adult sockeye						
783	salmon. Physiological and Biochemical Zoology 85:729-739.						
784	Donaldson, M. R., et al. 2013. Evaluation of a simple technique for recovering fish from capture						
785	stress: integrating physiology, biotelemetry, and social science to solve a conservation						
786	problem. Canadian Journal of Fisheries and Aquatic Sciences 70:90-100.						
787	Drury, R., K. Homewood, and S. Randall. 2011. Less is more: the potential of qualitative						
788	approaches in conservation research. Animal Conservation 14:18-24.						
789	English, K. K., W. R. Koski, C. Sliwinski, A. Blakley, A. Cass, and J. C. Woodey. 2005.						
790	Migration timing and river survival of late-run Fraser River sockeye salmon estimated using						
791	radiotelemetry techniques. Transactions of the American Fisheries Society 134:1342-1365.						
792	Farrell, A. P., P. E. Gallaugher, and R. Routledge. 2001. Rapid recovery of exhausted adult coho						
793	salmon after commercial capture by troll fishing. Canadian Journal of Fisheries and Aquatic						
794	Sciences 58:2319-2324.						
795	Fox, H. E., C. Christian, J. C. Nordby, O. R. W. Pergams, G. D. Peterson, and C. R. Pyke. 2006.						
796	Perceived barriers to integrating social science and conservation. Conservation Biology						
797	20:1817-1820.						

Gende, S. M., R. T. Edwards, M. F. Wilson, and M. S. Wipfli. 2002. Pacific salmon in aquatic
and terrestrial ecosystems. Bioscience 52:917-928.

- 800 Guerbois, C., E. Chapanda, and H. Fritz. 2012. Combining multi-scale socio-ecological
- approaches to understand the susceptibility of subsistence farmers to elephant crop raiding on

the edge of a protected area. Journal of Applied Ecology 49:1149-1158.

- 803 Gray, J. S. 1997. Marine biodiversity: patterns, threats and conservation needs. Biodiversity
- 804 Conservation 6:153–175.
- Hall, M. A., D. L. Alverson, and K. I. Metuzals. 2000. By-catch: problems and solutions. Marine
 Pollution Bulletin 41:204–219.
- 807 Huntington, H. P. 2000. Using traditional ecological knowledge in science: methods and
- applications. Ecological Applications 10:1270-1274
- 809 IFCRT (Interior Fraser Coho Recovery Team). 2006. Conservation strategy for coho salmon
- 810 (Oncorhynchus kisutch), interior Fraser River populations. Fisheries and Oceans Canada, 132
- 811 pp.
- 812 Irvine, R. J., S. Fiorini, S. Yearley, J. E. McLeod, A. Turner, H. Armstrong, P. C. L. White, and
- 813 R. van der Wal. 2009. Can managers inform models? Integrating local knowledge into
- models of red deer habitat use. Journal of Applied Ecology 46:344-352.
- 815 Kappel, C. V. 2005. Losing pieces of the puzzle: threats to marine, estuarine, and diadromous
- species. Frontiers in Ecology and the Environment 3:275-282.
- Lackey, R, L. 1999. Salmon policy: science, society, restoration, and reality. Renewable
- 818 Resources Journal 17:6-16.
- Lele, S., and A. Kurien. 2011. Interdisciplinary analysis of the environment: insights from
- tropical forest research. Environmental Conservation 38:211-233.
- Lowe, P., G. Whitman, and J. Phillipson. 2009. Ecology and the social sciences. Journal of
- 822 Applied Ecology 46:297-305.

- 823 Lupes, S. C., M. W. Davis, B. L. Olla, and C. B. Schreck. 2006. Capture-related stressors impair
- 824 immune system function in sablefish. Transactions of the American Fisheries Society

825 135:129-138.

- 826 Margles, S. W., R. B. Peterson, J. Ervin, and B. A. Kaplin. 2010. Conservation without borders:
- building communication and action across disciplinary boundaries for effective conservation.
 Environmental Management 45:1-4.
- 829 Martins, E. G., S. G. Hinch, D. A. Patterson, M. J. Hague, S. J. Cooke, K. M. Miller, M. F.
- Lapointe, K. K. English, and A. P. Farrell. 2011. Effects of river temperature and climate
- 831 warming on stock-specific survival of adult migrating Fraser River sockeye salmon
- 832 (*Oncorhynchus nerka*). Global Change Biology 17:99-114.
- 833 Milligan, C. L., G. B. Hooke, and C. Johnson. 2000. Sustained swimming at low velocity
- following a bout of exhaustive exercise enhances metabolic recovery in rainbow trout. The
- Journal of Experimental Biology 203:921-926.
- Nguyen, V. M., G. D. Raby, S. G. Hinch, and S. J. Cooke. 2012. Aboriginal fisher perspective on
- use of telemetry technology to study adult Pacific salmon. Knowledge and Management of
- Aquatic Ecosystems 406:08.
- 839 Nguyen, V. M., E. G. Martins, D. Robichaud, G. D. Raby, M. R. Donaldson, A. G. Lotto, W. G.
- 840 Willmore, D. A. Patterson, A. P. Farrell, S. G. Hinch, and S. J. Cooke. 2014. Disentangling
- the roles of air exposure, gillnet injury, and facilitated recovery on the post-capture and
- release mortality and behavior of adult migratory sockeye salmon (Oncorhynchus nerka) in
- freshwater. Physiological and Biochemical Zoology 87:125-135.
- 844 Raby, G. D., A. C. Colotelo, G. Blouin-Demers, and S. J. Cooke. 2011. Freshwater commercial
- bycatch: an understated conservation problem. BioScience 61:271-280.

846	Raby, G. D., M. R. Donaldson, S. G. Hinch, D. A. Patterson, A. G. Lotto, D. Robichaud, K. K.					
847	English, W. G. Willmore, A. P. Farrell, M. W. Davis, and S. J. Cooke. 2012. Validation of					
848	reflex indicators for measuring vitality and predicting the delayed mortality of wild coho					
849	salmon bycatch released from fishing gears. Journal of Applied Ecology 49:90-98.					
850	Raby, G. D., S. J. Cooke, K. V. Cook, S. H. McConnachie, M. R. Donaldson, S. G. Hinch, C. K					
851	Whitney, S. M. Drenner, D. A. Patterson, T. D. Clark, and A. P. Farrell. 2013. Resilience of					
852	pink salmon and chum salmon to simulated fisheries capture stress incurred upon arrival at					
853	spawning grounds. Transactions of the American Fisheries Society 142:524-539.					
854	Raby, G. D., J. R. Packer, A. J. Danylchuk, and S. J. Cooke. The understudied and					
855	underappreciated role of predation in the mortality of fish released from fishing gears. Fish					
856	and Fisheries, in press, DOI: 10.1111/faf.12049.					
857	Rogers, M. W., A. B. Barbour, and K. L. Wilson. 2014 Trade-offs in experimental designs for					
858	estimating post-release mortality in containment studies. Fisheries Research, 151:130-135.					
859	Scarnecchia, D. L. 1988. Salmon management and the search for values. Canadian Journal of					
860	Fisheries and Aquatic Sciences 45:2042-2050.					
861	Soykan, C. U., J. E. Moore, R. Zydelis, L. B. Crowder, C. Safina, R. L. Lewison. 2008. Why					
862	study bycatch? An introduction to the theme section on fisheries bycatch. Endangered					
863	Species Research 5:91–102.					
864	Strauss, A. L. 1987. Qualitative Analysis for Social Scientists. Cambridge University Press,					
865	Cambridge, UK.					

Sutherland, W. J., et al. 2009. One hundred questions of importance to the conservation of global
biological diversity. Conservation Biology 23:557-567.

868	Wilson, S. M., S. G. Hinch, E. J. Eliason, A. P. Farrell, and S. J. Cooke. 2013. Calibrating
869	acoustic acceleration transmitters for estimating energy use by wild adult Pacific salmon.
870	Comparative Biochemistry and Physiology Part A 164:491-498.
871	
872	
873	
874	Ecological Archives Material:
875	- Additional methods detail on coding human dimensions survey responses
876	
877	
878	
879	
880	
881	
882	
883	
884	
885	
886	
887	
888	
889	
890	

891 Tables

Table 1. Population (stock) proportions and individual stream IDs of coho salmon captured in the

- study from 2009-2011, determined via analysis of microsatellite variation in DNA (N = 223; see
- Beacham et al. 2011). These samples include fish tagged and released with transmitters and
- additional fish sampled for DNA (mostly blood-sampled fish from 2009; see Raby et al. 2012).
- All of these groups are part of the interior Fraser coho group except for the bottom three
- 897 populations, which split off the Fraser River main stem downstream of Hell's Gate. "Actual
- 898 percentage of overall population group" refers to the true percentage of the entire interior Fraser
- 899 River coho salmon group represented by each population, based on DFO spawning ground stock
- assessment data for the years 2009-2011 (R. Bailey, DFO, pers. comm. 03/10/12). Post-release
- survival was not significantly different among populations ($\gamma^2 = 1.39$, df = 5, P = 0.99).

Population	Actual	Mean Fork	Post-release	Stream origin
(% of interior	percentage of	Length	survival	(N)
Fraser coho)	overall	(± S.E.)	estimate (%)	
	population group			
North Thompson		61 ± 0.7 cm	67%	Pig Channel (37)
River				Lemieux Creek (20)
N = 86	35%			Louis Creek (8)
(41%)				Birch Island (7)
				Fennell River (6)
				Reg Christie Creek (5)
				Dunn Creek (2)
				Mann Creek (1)

Lower Thompson		$61 \pm 0.9 \text{ cm}$	73.6%	Coldwater River (28)
River				Spius Creek (12)
N = 41	23%			Bonaparte River (1)
(20%)				
South Thompson		62 ± 0.8 cm	58%	Eagle River (20)
River				Harbour Creek (7)
N = 38	21%			McMomee Creek (6)
(18%)				Bessette Creek (3)
				Salmon River (3)
				Wap Creek (2)
Fraser River		61 ± 0.9 cm	52%	Bridge River (22)
middle drainage				McKinley Creek (13)
N = 37	13%			Gates Creek (2)
(18%)			<u>, </u>	
Fraser River		67 ± 2.4 cm	71%	Nahatlatch River (7)
canyon				
N = 7	8%			
(3%)				
Birkenhead River		67 ± 3.8 cm	67%	Birkenhead River (6)
N = 6				
Fraser River		71 ± 2.2 cm	75%	Chehalis River (4)
lower drainage				
N = 4				



Table 2. Observed mortality rates of coho salmon bycatch for each of the three study years and 923 estimated cumulative mortality rates. Immediate mortality numbers were generated by direct 924 observation of whether individual fish were alive when released from the net, whereas mortality 925 926 estimates to 48 h and to natal sub-watersheds were made using radio telemetry. The overall estimates are weighted and cumulative such that the post-release mortality estimates were 927 applied to fish that were released alive but not tracked. Differences in post-release mortality 928 among years were not statistically significant (Pearson's chi-square test: $\chi^2 = 1.29$, df = 2, P = 929 0.53). The binomial probability 95% confidence interval for the final overall mortality rate 930 (39.3%) was 34 – 45%. For 48 h mortality (18.6%) it was 14-24%. 931

	Post-release mortality		mortality
	Immediate		To upper
Year (N tagged)	mortality	48 h	watersheds
	5%	6%	26%
2009* (50)	(5 of 104)	(3 of 50)	(13 of 50)
	2%	15%	43%
2010 (53)	(1 of 55)	(8 of 53)	(23 of 53)
	2%	23%	37%
2011 (79)	(3 of 135)	(18 of 79)	(29 of 79)
Overall	3%	18.6%	37.3%
estimates	9 of 295	55 of 295	110 of 295
	Final mortality	y rate estimate if	
	RAMP scores	used to predict	39.3%
fates of non-ta		gged fish	116 of 295

932 933	*From Raby et al. 2012
934	
935	
936	
937	
938	
939	
940	
941	
942	
943	
944	
945	
946	
947	
948	
949	
950	
951	
952	
953	
954	
955	

Table 3. Regression coefficients (*B*, with standard error) and odds ratios (with 95% CIs) for each

957 predictor variable included in the logistic regression model with post-release survival as the binary

958 response variable (0 = unsuccessful migrant, 1 = successful migrant). Odds ratios of below 1 indicate a

- 959 negative relationship with the response variable whereas the opposite is true of odds ratios above 1.
- 960 RAMP score was the only significant predictor of survival (P < 0.001).

		95% CI for odds		
			ratio	
Predictor variable	B (SE)	Lower	Odds ratio	Upper
Constant	1.19			
	(2.60)			
RAMP score	- 3.56**	0.004	0.03	0.16
	(0.93)			
Use of revival bag	-0.33	0.31	0.72	1.65
	(0.42)			
Entanglement time	4.5×10^{-4}	1.00	1.00	1.00
	(5.1 × 10 ⁻⁴)			
Total catch	-6.8×10^{-4}	1.00	1.00	1.00
	(5.3 × 10 ⁻⁴)			
Air temp.	0.05	0.94	1.05	1.17
	(0.06)			
Fork length (cm)	-1.1×10^{-3}	0.94	1.00	1.07
	(0.03)			

962 Note: Overall model $R^2 = 0.127$ (Hosmer-Lemeshow), 0.154 (Cox-Snell), 0.21 (Nagelkerke). Model $\chi^2(6)$ 963 = 24.45, P < 0.001. * P < 0.05, ** P < 0.001.

964

966 Figure captions

968	FIG. 1. Fraser River watershed map with locations of radio receivers used in different years and
969	the locations of the three main capture/release sites (Peg Leg, Seabird, and Peters). The spawning
970	areas of the four main interior Fraser coho populations are indicated and circled by grey hashed
971	lines (not circled: the Fraser canyon population, which comprises only one DNA-identified
972	spawning tributary - the Nahatlatch River ~ 30 rkm upstream of Hell's Gate).
973	
974	FIG. 2. Photos showing different ways in which the aboriginal fishing crews pulled their nets
975	into shore for sorting, during which "entanglement time" was recorded for individual coho
976	salmon caught as bycatch. Photos show a typical set (A), a very large set with high crowding that
977	required hours to sort through (B), and a smaller set that was pulled entirely onto shore for
978	sorting (C).
979	
980	FIG. 3. Images showing the fish recovery bag open at the water's surface (A) prior to having a
981	fish inserted for (B) a 30-min revival period in the river current with the bag attached to a
982	reinforcing bar driven into the riverbed.
983	
984	FIG. 4. Observed post-release mortality rates (whole bars) at each level of reflex impairment
985	(RAMP score). The black sections of each bar represent a proposed 'baseline' mortality rate
986	within the dataset, whereby only mortality that occurred above that level was assigned to the
987	fishers. Develop this selection, the much are static bettern of each her much selection of
	isnery. Based on this calculation, the numbers at the bottom of each bar represent the number of
988	fish assigned as survivors of the capture event at each RAMP score, while the numbers in the

black bars are the numbers of fish assigned as mortalities at each RAMP score by applying a net 989 mortality rate (total mortality minus 23%) to the total number of fish with each RAMP score. For 990 example, the net mortality rate used for the fish with RAMP scores of 0.6 was 28% (51% - 23% 991 baseline), which was then applied to the total number of fish tagged at that RAMP score (N =992 37). The two numbers within each bar add to the actual total number of fish released with 993 transmitters at that RAMP score, but note that the total N in this figure is 176 – although 182 994 coho salmon were tagged and tracked, RAMP scores were not recorded for 6 individuals. Thus, 995 this alternate post-release mortality rate of 13.6% comes from dividing the cumulative fishery 996 997 mortalities indicated (24) by the total (176). 998 FIG. 5. Responses of aboriginal fishers in our human dimensions surveys to questions relating to 999 1000 coho salmon capture stress and mortality. For A there were 58 responses and for B, N = 70. Responses of individual fishers were coded into the response categories shown in the figure, and 1001 according to further detail available in the online supplementary material for this paper. 1002 1003 FIG. 6. Responses of aboriginal fishers in our human dimensions surveys to questions relating to 1004 ways to improve the survival of coho salmon bycatch in the beach seine fishery. Responses of 1005 1006 individual fishers for both questions were coded into the response categories shown in the figure, and according to details available in the online supplementary material for this paper. There were 1007 63 responses for question A. For B, N = 54. 1008

1009

FIG. 7. Responses of aboriginal fishers in our human dimensions surveys to questions relating tothe use of fish recovery bags for coho salmon bycatch in the beach seine fishery. Prior to asking

- 1012 questions A (N = 94 responses), B (N = 96), and C (N = 86), the interviewees were shown a
- 1013 recovery bag and its purpose and use explained to them. Responses of individual fishers were
- 1014 coded into the response categories shown in the figure, and according to details available in the
- 1015 online supplementary material for this paper.
- 1016







ECOLOGICAL SOCIETY OF AMERICA

What do you think causes the greatest stress for coho released from beach seines?

If you knew that leaving the seine in knee-high water for sorting would increase the chance of survival for released fish, would you voluntarily do it?

Should the recovery bag method be mandatary for reviving coho salmon after beach seine capture?

What are your thoughts on the recovery bag? (after description provided)