

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/353210435>

Twenty-five essential research questions to inform the protection and restoration of freshwater biodiversity

Article in *Aquatic Conservation Marine and Freshwater Ecosystems* · July 2021

DOI: 10.1002/aqc.3634

CITATIONS

4

READS

1,066

36 authors, including:



Meagan Harper

4 PUBLICATIONS 17 CITATIONS

[SEE PROFILE](#)



Hebah Mejbel

University of Ottawa

6 PUBLICATIONS 12 CITATIONS

[SEE PROFILE](#)



Robin Abell

The Nature Conservancy

37 PUBLICATIONS 6,675 CITATIONS

[SEE PROFILE](#)



Joseph R Bennett

Carleton University

134 PUBLICATIONS 2,085 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



LOFRESH: Understanding the ecological relevance of lotic environmental DNA [View project](#)



Species description *Romanogobio* sp.nov. [View project](#)

VIEWPOINT

Twenty-five essential research questions to inform the protection and restoration of freshwater biodiversity

Meagan Harper¹  | Hebah S. Mejbil²  | Dylan Longert² | Robin Abell³  | T. Douglas Beard⁴  | Joseph R. Bennett¹  | Stephanie M. Carlson⁵  | William Darwall⁶  | Anthony Dell⁷ | Sami Domisch⁸  | David Dudgeon⁹  | Jörg Freyhof¹⁰  | Ian Harrison¹¹  | Kathy A. Hughes¹² | Sonja C. Jähnig^{8,13}  | Jonathan M. Jeschke^{8,14,15} | Richard Lansdown¹⁶ | Mark Lintermans¹⁷ | Abigail J. Lynch¹⁸ | Helen M. R. Meredith¹⁹ | Sanjay Molur²⁰ | Julian D. Olden²¹ | Steve J. Ormerod^{22,23} | Harmony Patricio²⁴  | Andrea J. Reid²⁵  | Astrid Schmidt-Kloiber²⁶  | Michele Thieme²⁷  | David Tickner¹² | Eren Turak²⁸  | Olaf L. F. Weyl^{29†} | Steven J. Cooke¹ 

¹Department of Biology, Carleton University, Ottawa, Ontario, Canada

²Department of Biology, University of Ottawa, Ottawa, Ontario, Canada

³Center for Sustainable Lands and Waters, Conservation International, Arlington, Virginia, USA

⁴National Climate Adaptation Science Center, US Geological Survey, Madison, Wisconsin, USA

⁵Environmental Science, Policy, and Management, University of California, Berkeley, California, USA

⁶Freshwater Biodiversity Unit, IUCN Global Species Programme, Cambridge, UK

⁷National Great Rivers Research and Education Center, East Alton, Illinois and Washington University in St Louis, St Louis, Missouri, USA

⁸Department of Ecosystem Research, Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin, Germany

⁹Division of Ecology and Biodiversity, School of Biological Sciences, The University of Hong Kong, Hong Kong

¹⁰Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science, Berlin, Germany

¹¹Moore Centre for Science, Conservation International, Arlington, Virginia, USA

¹²Science & Conservation Department, Living Planet Centre, Woking, WWF-UK, UK

¹³Geography Department, Humboldt-Universität zu Berlin, Berlin, Germany

¹⁴Institute of Biology, Freie Universität Berlin, Berlin, Germany

¹⁵Berlin-Brandenburg Institute of Advanced Biodiversity Research, Berlin, Germany

¹⁶IUCN SSC Freshwater Plant Specialist Group, Gloucestershire, UK

¹⁷Centre for Applied Water Science, Institute for Applied Ecology, University of Canberra, Canberra, Australian Capital Territory, Australia

¹⁸US Geological Survey, National Climate Adaptation Science Center, Reston, Virginia, USA

¹⁹Amphibian Survival Alliance, % Synchronicity Earth, London, UK

²⁰Systematics, Ecology, and Conservation Laboratory, Zoo Outreach Organization, Coimbatore, Tamil Nadu, India

²¹School of Aquatic and Fishery Sciences, University of Washington, Seattle, Washington, USA

²²School of Biosciences, Cardiff University, Cardiff, UK

²³Freshwater Biological Association, Ambleside, Cumbria, UK

²⁴Commission on Protected Areas Freshwater Specialist Group, IUCN World, Telluride, Colorado, USA

²⁵Institute for the Oceans and Fisheries, The University of British Columbia, Vancouver, Canada

²⁶Institute of Hydrobiology and Aquatic Ecosystem Management, University of Natural Resources and Life Sciences, Vienna, Vienna, Austria

Meagan Harper, Hebah S. Mejbil and Dylan Longert contributed equally.

† Deceased.

²⁷Freshwater, World Wildlife Fund-US, Washington, DC, USA

²⁸NSW Department of Planning, Industry and the Environment, Parramatta, New South Wales, Australia

²⁹Centre for Invasion Biology, South African Institute for Aquatic Biodiversity, Makhanda, South Africa

Correspondence

Meagan Harper, Department of Biology,
Carleton University, 1125 Colonel By Drive,
Ottawa, Ontario K1S 5B6, Canada.
Email: meaganharper@cmail.carleton.ca

Funding information

National Sciences and Engineering Research
Council of Canada

Abstract

1. Freshwater biodiversity is declining at an unprecedented rate. Freshwater conservationists and environmental managers have enough evidence to demonstrate that action must not be delayed but have insufficient evidence to identify those actions that will be most effective in reversing the current trend.
2. Here, the focus is on identifying essential research topics that, if addressed, will contribute directly to restoring freshwater biodiversity through supporting ‘bending the curve’ actions (i.e. those actions leading to the recovery of freshwater biodiversity, not simply deceleration of the current downward trend).
3. The global freshwater research and management community was asked to identify unanswered research questions that could address knowledge gaps and barriers associated with ‘bending the curve’ actions. The resulting list was refined into six themes and 25 questions.
4. Although context-dependent and potentially limited in global reach, six overarching themes were identified: (i) learning from successes and failures; (ii) improving current practices; (iii) balancing resource needs; (iv) rethinking built environments; (v) reforming policy and investments; and (vi) enabling transformative change.
5. Bold, efficient, science-based actions are necessary to reverse biodiversity loss. We believe that conservation actions will be most effective when supported by sound evidence, and that research and action must complement one another. These questions are intended to guide global freshwater researchers and conservation practitioners, identify key projects and signal research needs to funders and governments. Our questions can act as springboards for multidisciplinary and multisectoral collaborations that will improve the management and restoration of freshwater biodiversity.

KEYWORDS

‘bending the curve’, freshwater conservation, horizon scanning, priority setting, research questions

1 | INTRODUCTION

Freshwater biodiversity faces unprecedented threats from human activities (Dudgeon et al., 2006; Reid et al., 2019). Many of these threats have been increasing in severity in recent decades (e.g. invasive species, fragmentation of rivers by dams, habitat loss), but there are also emerging threats (e.g. novel pollutants and pathogens, climate change), as well as interactions and cumulative effects (Birk et al., 2020) that further threaten freshwater biodiversity (Reid et al., 2019). Given how catchments function, everything that occurs in upland areas has the potential to affect freshwater ecosystems downstream. Even activities that happen well beyond the floodplain and riparian areas can have severe effects on freshwater biodiversity

(Hynes, 1975; Weijters et al., 2009). Recent estimates have shown that, on average, the abundance of monitored freshwater vertebrate populations in the Freshwater Living Planet Index has declined by an average of 84% over the past five decades (World Wildlife Fund, 2020), double the rate of decline seen in marine and terrestrial realms. This has led to the recognition of the current global freshwater biodiversity emergency (Tickner et al., 2020). In addition, approximately 30% of the freshwater species assessed by the International Union for Conservation of Nature (IUCN) are threatened (i.e. Critically Endangered, Endangered or Vulnerable to global extinction; IUCN, 2012) in the Americas, more than 20% are threatened in Africa, and in Europe and Central Asia 37% of freshwater fish, 45% of freshwater snails and 23% of amphibians are threatened (Watson

et al., 2018). To facilitate management interventions that can effectively curtail or even reverse the decline in freshwater biota (i.e. 'bending the curve' of biodiversity loss to enable the recovery of freshwater biodiversity), research and conservation practices must continue to be coordinated to address key knowledge gaps that, at present, impede progress (Mace et al., 2018; van Rees et al., 2020; Tickner et al., 2020).

Often, current research remains focused on improving understanding of natural history and the present status of freshwater biodiversity, and identifying the effects of various human threats. This research is critical, but it is also essential to ensure that there is dedicated research on actions that will directly alter and reverse the current downward trajectory of biodiversity loss. We define 'bending the curve' actions in freshwater biodiversity conservation as those that will lead to the recovery of freshwater biodiversity (*sensu* Tickner et al., 2020) as opposed to the deceleration or stabilization of the current downward trend. Actions for bending the curve aim to guide restoration and conservation, engage with the public and decision-makers, and target investments in tools, research and policy. Those actions will reverse the impacts on freshwater biodiversity loss of direct threats (e.g. point source pollution) and indirect drivers (e.g. climate change). Research on the status of freshwater life and on new threats to its existence is an essential part of conservation but knowledge gaps in these areas are already well recognized (Reid et al., 2019). Instead, inspired by recent calls to motivate change (van Rees et al., 2020; Tickner et al., 2020), we focus on identifying essential research areas in the natural and social sciences that will support efforts towards the recovery of freshwater biodiversity.

In contrast to marine science (Parsons et al., 2014), which is better represented in conservation science in general (Boon & Baxter, 2016), there have been few research agendas in freshwater science focused directly on biodiversity. Current freshwater biodiversity research agendas include one focused on migratory fishes (Lennox et al., 2019), a broader European agenda focused on overall biodiversity loss with freshwater content (European Commission, 2011), a preliminary unpublished freshwater research agenda (BioFresh, 2011) and various national agendas (Jähnig et al., 2019). None of these explicitly focus on research that will help in bending the curve. Despite the recent development of frameworks detailing the major causes of freshwater biodiversity loss (Strayer & Dudgeon, 2010; Garcia-Moreno et al., 2014; Flitcroft et al., 2019) and efforts to support post-2020 policy agendas (van Rees et al., 2020), the issue of targeting research to facilitate freshwater biodiversity recovery remains challenging.

To address this challenge, a broad sample of the global freshwater research and management community was solicited to identify unanswered research questions in freshwater biodiversity conservation. Through this outreach, six overarching themes were identified that encompass important areas for future research. Within these themes, both foundational and cross-cutting issues and specific strategies and challenges inherent to freshwater biodiversity conservation are presented together to ensure that future research efforts are built on robust foundations and provide useable outcomes. Broad questions

within these themes were identified, as were examples of possible research questions (ranging from narrow to broad) that would aid the freshwater community in effectively reversing freshwater biodiversity loss. These themes and questions are intended to serve as a guide for freshwater scientists, conservation practitioners, research funders and policymakers by pointing to possible future projects and identifying pressing research topics and priorities related to bending the curve of freshwater biodiversity loss. We acknowledge that there are other broader conservation science questions that extend across realms (e.g. marine, terrestrial, freshwater) especially related to social science (Bennett et al., 2017b), as well as critical social justice issues pertaining to freshwater health (Mascarenhas, 2007). The questions presented here are those specifically related to freshwater biodiversity conservation.

2 | DERIVING QUESTIONS AND IDENTIFYING THEMES

The best practices identified in Sutherland et al. (2011) were adopted to guide this exercise. Original questions were solicited through an online questionnaire (surveyplanet.com) and requests for participation were distributed by the authors through targeted emails, list-serves and social media between 23 September and 1 November 2019. The call for questions was shared as broadly as possible by the authors and their network contacts with no limits on outreach (i.e. a 'snowball approach' or 'chain-referral sampling'). It was not possible, therefore, to quantify the full extent of the reach of the call for questions, which is typical of the approach by Sutherland et al. (2011) for these exercises. It is not known how many individuals or nations received a request to participate (or were aware of the survey) and chose not to respond. Those who did respond were asked to provide questions that would help address the knowledge gaps and barriers associated with bending the curve of freshwater biodiversity loss, as well as to provide information on their sector, role and geographical location. To obtain as many questions as possible and to allow participants to contribute fully, there were no limits to the number of times an individual could participate.

The call for questions achieved global reach with participants active in 45 countries (Table 1; Figure S2); however, it is important to note that 27 of these 45 countries (60%) had a single respondent. The top three participating countries were Canada ($n = 25$ participants),

TABLE 1 Participants by geographical region

Region	Number of participants
North America	48
Central and South America	4
Asia-Pacific	35
Europe	49
Africa	16

the United States ($n = 23$), and Australia ($n = 18$). Participants represented all sectors: Industry ($n = 2$; 1.2%), Government ($n = 30$; 18.5%), Not-for-profit ($n = 48$; 29.6%) and Academic ($n = 61$; 37.7%), and an additional 21 participants (13%) who self-identified as Other (Figure 1a). Several participants ($n = 11$) selected more than one sector. The most common primary role was Researcher ($n = 74$; 43.3%), followed by Practitioner ($n = 35$; 20.5%), Decision-maker ($n = 20$; 11.7%), Other ($n = 25$; 14.6%) and Student/post-doc ($n = 17$; 9.9%); the only unrepresented primary role was Funder (Figure 1b). As with sector, participants often selected more than one primary role; a total of 21 participants selected two or more.

An initial list of 424 questions, submitted by 144 participants, was screened by the review team (MH, HSM, DL, and SJC). Questions that were deemed less applicable to the aim of bending the curve were removed. Questions removed included those that were: (i) highly region specific, (ii) extremely taxonomically specific (e.g. regarding life history of a single species), (iii) focused on threat identification (e.g. the impact of X on Y) and (iv) based on natural history (e.g. where does X species spawn?). Questions aimed at guiding restoration and conservation, educating the public and decision-makers, and targeting investments in tools, research and policy to lead to the recovery of freshwater biodiversity were retained (see Supporting Information for more detailed methodology and expanded results, and Table S2 for the complete list of submitted questions). After the initial screening by the review team, a short list of questions was evaluated by all authors to group or split specific questions, suggest re-wording for clarity and assess the likelihood of a question leading to research that would advance actions for bending the curve. In addition, all authors had the opportunity to advocate for questions that had been initially removed or to suggest their own. The final list of questions was selected through an iterative process and edited by all authors, including the review team, and were then condensed to six major themes (Figure 2) using the methods described in Sutherland et al. (2011).

Six major themes are presented, each including several broad 'essential questions' (25 questions in total) that represent knowledge gaps and areas of concern identified by the respondents to our call for questions and by our author team. Whereas Tickner et al. (2020) presented six curve-bending actions for freshwater biodiversity (representing one framework for assigning questions to themes), the essential questions (and research needs) presented here transcend and cut across those actions, so are grouped into slightly different themes (Table S3). The 25 essential questions are presented in no particular order, as priorities inevitably depend on the context and vary by geographical region and in response to socio-economic and political realities. These questions could be arranged under a variety of overlapping and cross-boundary themes, while themes and questions can interact in the development of specific hypotheses. This selection was further expanded with a limited subset of 75 possible research questions (Table 2) ranging from those that are narrowly focused to others that are broadly applicable. These additional example questions reflect some of the diversity of interests and the stage of development globally of freshwater biodiversity research. Such lists could be virtually endless, so these 75 further examples are just that – examples of specific questions which, if answered, could help further bend the curve of freshwater biodiversity loss.

3 | SIX THEMES AND 25 QUESTIONS

3.1 | Theme 1: Learning from successes and failures

This theme considers what can be learned from previous successes and failures in biodiversity conservation and how that knowledge can be applied to present and future initiatives. Understanding the strategies and tactics that are most effective and efficient in




FIGURE 1 (a) Frequency (%) of participants from different sectors involved in freshwater biodiversity research and protection including industry, government, not-for-profit organizations and other sectors. (b) Frequency (%) of participants with different primary roles including students/post-docs, decision-makers, practitioners, researchers and other primary roles. No funders participated in our call for questions

FIGURE 2 Six major themes for ‘bending the curve’ of freshwater biodiversity loss. ‘Learning from successes and failures’ and ‘Improving current practices’ focus on improving conservation and protection of freshwater biodiversity; ‘Balancing resource needs’ and ‘Rethinking built environments’ consider balancing human and freshwater biodiversity needs; ‘Reforming policy and investment’ and ‘Enabling transformative change’ emphasize the need to improve funding, knowledge exchange and public engagement in freshwater biodiversity research and conservation



producing lasting conservation impact, at scale, in the face of complex and increasingly dynamic socio-economic, political, cultural and governance challenges is an essential component of learning from successes and failures. Questions included in this theme assess the characteristics of protected areas for freshwater organisms, consider the spatial scale of conservation initiatives, the effectiveness of flagship and umbrella species in freshwater biodiversity restoration, and the benefits of effective monitoring. The identification of successful conservation initiatives, when scaled up (Bennett et al., 2016), can lead to improvements in freshwater biodiversity.

3.1.1 | Question 1: Opportunities for learning:
Where and why have past conservation efforts been successful or failed, and how can we learn from these outcomes?

In disciplines such as business, it is common practice to engage in extensive, formal reflective processes to learn from success and failure (Lant & Montgomery, 1987). Only recently has this idea been fully embraced by the conservation science community (Knight, 2006), but often successes are celebrated and failures forgotten. Also troubling is

the fact that many current efforts in freshwater biodiversity conservation appear to be ineffective in the face of an increasing number of persistent, emerging and synergistic or additive stressors (Craig et al., 2017). Efforts to understand the enabling factors for success can be illuminating, and further research on factors that extend beyond the ecological realm (including economic, institutional, social and cultural factors) can contribute to determining the ultimate success of conservation initiatives. Learning from success and failure, with a focus on identifying enabling factors, provides opportunities to support evidence-based conservation for long-term freshwater conservation outcomes.

3.1.2 | Question 2: Optimizing scale: At what spatial and temporal scales are management interventions best applied to benefit freshwater biodiversity?

To improve management of freshwater biodiversity, the spatial and temporal scales of conservation initiatives must be considered. The scales at which conservation efforts are implemented is a primary factor in how freshwater biodiversity is enhanced and which species

TABLE 2 Example research questions for each of the 25 essential questions

Theme	Essential question	Example research questions
1. Learning from successes and failures	1. Where and why have past conservation efforts been successful or failed, and how can we learn from these outcomes?	<ol style="list-style-type: none"> 1. What lessons stand to be gained from successful efforts for expanding the application of freshwater conservation policies? 2. How can conservation success stories be translated into increased resilience and resistance to perturbation for freshwater species' populations? 3. What are the different contributing factors and elements of success for different types of freshwater ecosystems?
	2. At what spatial scale and temporal scales are management interventions best applied to benefit freshwater biodiversity?	<ol style="list-style-type: none"> 1. How can we develop a better understanding of the interconnectedness of terrestrial and aquatic ecosystems for improved freshwater restoration? 2. How can catchment approaches be delivered on a sufficiently broad scale to reverse freshwater biodiversity decline? 3. To what extent can local-scale management interventions (e.g. property scale) reduce threats to freshwater biodiversity and what are the cost/benefit implications of making changes at different scales?
	3. What are the characteristics of current protected areas and networks, as well as lands and waters stewarded and managed by Indigenous people, that lead to improved status of freshwater biodiversity and how can these be employed in future conservation efforts?	<ol style="list-style-type: none"> 1. What spatial gaps in protected areas need to be addressed to ensure successful management strategies? 2. How and where should freshwater protected areas be established? 3. How can protected-area networks incorporate connectivity between terrestrial, freshwater, and marine systems to successfully protect freshwater ecosystems?
	4. How can flagship or umbrella species be effectively used to both increase restoration and protection of freshwater biodiversity and increase public involvement in freshwater biodiversity restoration initiatives?	<ol style="list-style-type: none"> 1. Which threatened taxa are umbrella species candidates for freshwater conservation? 2. How can the often-overlooked components of freshwater biodiversity (plants, invertebrates, amphibians, etc.) be prioritized for flagship and/or umbrella species? 3. What is the umbrella potential of freshwater mega-fauna?
	5. How can we improve monitoring metrics and resources to guide restoration, conservation and sustainable management of freshwater biodiversity?	<ol style="list-style-type: none"> 1. Is freshwater biodiversity conservation improved by concentrating efforts in a single location or spreading efforts over multiple locations? 2. How can we improve freshwater biodiversity monitoring in historically under-represented regions and habitat types? 3. What are the key elements in a successful global freshwater biodiversity monitoring programme and how can they be implemented in the most cost-effective manner?
2. Improving current practices	6. What are the Key Biodiversity Areas that need to be prioritized for conservation of freshwater biodiversity?	<ol style="list-style-type: none"> 1. How can we prioritize key sites that, if restored, would provide the greatest improvements to the condition of freshwater ecosystems and freshwater biodiversity? 2. How should we select areas from which future human activities should be barred through strict conservation initiatives? 3. How can the protection of freshwater Key Biodiversity Areas be improved, both through legal, and physical means (i.e. barriers)?
	7. What approaches to pollution reduction and remediation efforts will most benefit freshwater biodiversity?	<ol style="list-style-type: none"> 1. How can we effectively communicate to industrial and commercial entities the dangers of dumping waste (physical and chemical) into freshwater systems and provide cost-effective solutions to the creation and safe disposal of waste? 2. To what extent are nature-based solutions applicable to point and non-point source pollution control in freshwater ecosystems? 3. How can the effects of newly emerging contaminants such as pharmaceuticals, microplastics etc. in freshwater systems be detected and mitigated more effectively?
	8. What research innovations are needed to help restore freshwater biodiversity?	<ol style="list-style-type: none"> 1. How can established management tools, such as repatriation of local biota, field assessments and stocking in freshwater biodiversity conservation, be improved? 2. What novel techniques (e.g. drones, eDNA, community science) could be applied to develop knowledge for improved freshwater biodiversity monitoring, conservation, and restoration activities?

TABLE 2 (Continued)

Theme	Essential question	Example research questions
	9. How do we incorporate climate change adaptation into freshwater biodiversity conservation?	<ol style="list-style-type: none"> 3. How can resilience assessments inform decision-making for freshwater biodiversity conservation? 1. Are current, conventional measures and metrics adequate to evaluate climate change effects (e.g. securing fish passage, water quality) and, if not, how can we improve them? 2. How can restoration projects incorporate resilience to a variety of climate impacts? 3. How should the climate change impacts on water resources best be mitigated to maintain optimal ecosystem function and services?
	10. What are the best ways to manage freshwater invasive species and diseases to ensure proactive and meaningful improvements to freshwater biodiversity?	<ol style="list-style-type: none"> 1. What are emerging pathways of new species introductions and how can they be managed to prevent harmful invasions from occurring in the future? 2. How can we improve measures to control or slow the spread of invasive species, including using techniques such as integrated risk assessments, biotechnology and community science? 3. How can proactive invasive species risk management, rather than reactive management (i.e. eradication), be integrated with current practices?
	11. What are the optimal riparian management actions that contribute to the protection of freshwater biodiversity?	<ol style="list-style-type: none"> 1. How do riparian zone setbacks modulate impacts of land-use change? 2. How can lateral continuity be better maintained in riparian zones? 3. What evidence will convince developers and planning authorities that human activities in riparian zones have dramatic effects on freshwater biodiversity and should be avoided?
	12. How can we develop conservation and restoration measures that most effectively and efficiently address synergistic threats to freshwater biodiversity?	<ol style="list-style-type: none"> 1. How can field-based experiments be improved in terms of scale and scope to identify management strategies that effectively decrease the negative effects of synergistic and additive stressors? 2. What management approaches used for individual threats could be utilized for effective management of multiple threats? 3. What measures could be applied to mitigate the confounding effects of climate change and warming-induced weather events (e.g. wildfires, hurricanes) on freshwater biodiversity?
3. Balancing resource needs	13. What are the joint priorities for sustainable food production and freshwater biodiversity conservation?	<ol style="list-style-type: none"> 1. How can we move away from traditional/industrialized inland fisheries management towards sustainable harvesting and improved conservation practices? 2. How can land-based agricultural practices (e.g. ranching or irrigation) be reformed to integrate freshwater biodiversity? 3. What steps can aquaculture take to ensure freshwater biodiversity is protected from escapees, disease and genetic alterations?
	14. How can the need for dams and associated infrastructure be balanced with connectivity, health and flow requirements of freshwater ecosystems and biodiversity?	<ol style="list-style-type: none"> 1. How can we enhance and operate existing dams to reduce impacts on freshwater species, and achieve energy production and conservation objectives? 2. How can site selection for new large and small hydropower projects be improved to reduce impacts on freshwater biodiversity? 3. What are the alternatives to traditional hydropower (i.e. dams) and how can these non-traditional options be adopted?
	15. How can we better balance conflicting interests between human demand for natural resources and freshwater biodiversity?	<ol style="list-style-type: none"> 1. How can we regulate human activities and resource use to better accommodate the needs of natural systems? 2. How can water abstraction (i.e. groundwater or surface water extraction) be mitigated to reduce the impacts on freshwater ecosystems and habitats? 3. What types of innovative technological efficiencies can decrease the impacts of, and demand for, resource extraction (e.g. sand alternatives) and benefit freshwater biodiversity?
4. Rethinking built environments	16. What policies, programmes and activities can we implement to turn the risks associated with urbanization into benefits/opportunities for freshwater biodiversity enhancement?	<ol style="list-style-type: none"> 1. Which urban restoration and rehabilitation actions provide the most effective results for enhancing freshwater biodiversity? 2. How can the distribution of people in cities be optimized to avoid destruction or degradation of wetlands and floodplains?

(Continues)

TABLE 2 (Continued)

Theme	Essential question	Example research questions
	17. How can freshwater biodiversity conservation be better integrated into economic infrastructure planning, implementation and operation?	<ol style="list-style-type: none"> 1. How can water allocation systems be redesigned to ensure sufficient water for freshwater ecosystems? 2. How can wastewater infrastructure be adapted to contribute to freshwater habitat development? 3. What changes to transportation infrastructure could decrease fragmentation and reinstate movement of freshwater species through enhanced freshwater connectivity?
	18. What is the role of novel and designed ecosystems in conservation, and how can these systems be managed to benefit freshwater biodiversity?	<ol style="list-style-type: none"> 1. How do we recognize ecosystems that cannot be returned to pre-disturbance conditions and how do we intervene to restore new biodiversity value, despite the changes experienced? 2. How can ecosystems, such as retention ponds and similar human-made features, be designed to provide sanctuaries for threatened species? 3. What management approaches are most applicable in novel and designed ecosystems to support native freshwater biodiversity?
5. Reforming policy and investment	19. What public policy measures can most effectively promote conservation and restoration of freshwater biodiversity?	<ol style="list-style-type: none"> 1. How can we aid decision-makers in improving their understanding of the state of freshwater biodiversity to gain additional political support in complementary legislation? 2. What policy strategies can be used to improve long-term funding stability for freshwater conservation management projects? 3. How can government strategies be improved to integrate freshwater biodiversity into policy to avoid contradictory regulatory objectives?
	20. How can we scale up and optimize financial investments from government, private sector and other sources such that there is a step change in funding for global freshwater conservation and restoration efforts?	<ol style="list-style-type: none"> 1. Would quantification and communication of the economic consequences of freshwater biodiversity loss be an effective method to convince stakeholders to increase investment? 2. How can data portals and knowledge platforms be used to help decrease conservation costs and to optimize the reallocation of funds? 3. What valuation methods should we use to embed freshwater biodiversity in freshwater ecosystem services, to make protection and restoration more adoptable?
	21. What are the social and natural science investments needed to develop and implement environmental flows that benefit freshwater biodiversity?	<ol style="list-style-type: none"> 1. What methods can we use to better link the components of artificially altered hydrology to biodiversity in perennial and non-perennial streams? 2. How can we mainstream and implement the principles of environmental flows within national legislation? 3. What scale of environmental flow implementation leads to improved freshwater biodiversity outcomes?
	22. What type of investments in <i>ex situ</i> conservation (e.g. captive breeding, reintroduction, managed relocation) are most effective for imperilled freshwater biodiversity?	<ol style="list-style-type: none"> 1. At what thresholds or trends of population abundance or decline does it make sense to invest in <i>ex situ</i> initiatives for different taxa? 2. Under what conditions do the benefits outweigh the risks/costs for <i>ex situ</i> conservation of threatened freshwater species? 3. What policies could be implemented to avoid genetic homogenization in <i>ex situ</i> conservation initiatives?
6. Enabling transformative change	23. How do we develop management frameworks and evidence bases that gain greater traction with stakeholders and managers?	<ol style="list-style-type: none"> 1. How can disparate evidence-bases (e.g. academic, corporate, Indigenous) be integrated to support improved conservation outcomes? 2. How can prioritization frameworks be adapted to improve inclusion of stakeholders in conservation and restoration? 3. Can specific freshwater biodiversity frameworks be developed to improve conservation outcomes and returns at national and international levels?
	24. What steps should be taken to better communicate and share evidence and knowledge about the science of freshwater biodiversity among stakeholders?	<ol style="list-style-type: none"> 1. How can we do a better job of translating scientific findings into actions for on-the-ground practitioners? 2. How do we improve communication and exchange of scientific findings with underrepresented regions, especially where language or restricted dissemination of research creates barriers? 3. How can Findable Accessible Interoperable Reusable (FAIR) data principles be best implemented into freshwater biodiversity science for

TABLE 2 (Continued)

Theme	Essential question	Example research questions
	25. How can we increase the level of public engagement to change mindsets and build social licence and political will to 'bend the curve' of freshwater biodiversity loss?	<p>the longevity of research findings (e.g. systematic publishing processes for data)?</p> <ol style="list-style-type: none"> 1. What innovative new techniques can be developed for more effectively engaging the general public and fostering greater understanding of (and care for) our freshwater biodiversity and ecosystems? 2. What is needed to shift mindsets and inspire the next generation to be excellent ambassadors and custodians of freshwater biodiversity? 3. How can we broaden the current models and orthodoxies at the science-policy interface to integrate worldviews from Indigenous and multicultural understandings?

Note: the inclusion of a specific example research question does not imply that it has any particular importance or priority over others. The examples were selected to emphasize the diversity of ways in which the essential question can be addressed, from very localized, perhaps taxon-specific research, to broader, multiregional or even global research that spans taxa and systems.

and populations benefit (Lintermans, 2013). Implementing freshwater conservation at effective scales often involves trade-offs of terrestrial or aquatic resource exploitation with downstream consequences. It is necessary to assess the effectiveness and interactions of strategies at different scales to mitigate, restore or avoid adverse impacts (Feld et al., 2018). A key determinant of success in conserving freshwater biodiversity is the development of integrative assessments of appropriate catchment scales required for effective results, recognizing that conservation efforts must adapt through time. For example, increasing habitat connectivity at different scales can promote species diversity (Shao et al., 2019) and enhance population resilience to climate change (Jaeger, Olden & Pelland, 2014), if done responsibly to avoid unintended consequences, such as species invasions.

3.1.3 | Question 3: Protected areas: What are the characteristics of current protected areas and networks, as well as lands and waters stewarded and managed by Indigenous people, that lead to improved status of freshwater biodiversity and how can these be employed in future conservation efforts?

The use of protected areas in freshwater ecosystems often lags behind those for marine or terrestrial ecosystems (Hermoso et al., 2016; Loury et al., 2018). Resource use in protected areas recognized by IUCN varies widely and ecosystem protection is inconsistent as a consequence. The responses of freshwater organisms to protected areas remains variable, but there is a growing body of evidence suggesting that protected areas can be a useful tool for freshwater biodiversity conservation provided that their design and management is robust (see Acreman et al., 2019 for a systematic review specifically related to the impacts of freshwater protected areas). Indigenous lands may function similarly, although less is known about aquatic systems on these lands (but see Schuster et al., 2019, for a terrestrial example). Although catchment-scale protected areas are highly desirable (Saunders, Meeuwig &

Vincent, 2002), protected areas are often more limited in size. Research is needed to understand how to enable broader implementation and management of protected areas for both groundwater and surface water, and the optimal configurations and management approaches when full catchment-scale protection is not possible (for a fuller discussion on systematic conservation planning, see Question 15). This will require consideration of alternatives to traditional top-down approaches to protected areas; for an example, consider the community-level fish sanctuaries used in Thailand that have benefited both fish biodiversity and community members who depend on these fisheries (Koning et al., 2020).

3.1.4 | Question 4: Flagship/umbrella species: How can flagship or umbrella species be used effectively both to increase restoration and protection of freshwater biodiversity and increase public involvement in freshwater biodiversity restoration initiatives?

The concepts of flagship and umbrella species have been applied successfully in terrestrial systems (e.g. giant pandas serving as both; Li & Pimm, 2016) and could be similarly successful in freshwater environments (e.g. freshwater turtles; Kalinkat et al., 2017). Flagship species act as ambassadors for conservation and are used to raise conservation funding and to attract public attention. Umbrella species are expected to benefit a wide range of co-occurring species. Questions remain regarding which species to select and whether they should be endemic or threatened, megafauna or from often overlooked groups (e.g. benthic invertebrates, Ormerod et al., 2010, or macrophytes), or if they truly function as intended. Similarly, it is not certain whether more general, systematic techniques for choosing flagship species (Verissimo et al., 2014; McGowan et al., 2020) are applicable to aquatic ecosystems. Working across disciplines with marketing and communications professionals to select species that resonate with the public and meet ecological goals may increase the success of these initiatives (Kalinkat et al., 2017).

3.1.5 | Question 5: Monitoring: How can we improve monitoring metrics and resources to guide restoration, conservation and sustainable management of freshwater biodiversity?

Some freshwater ecosystems are subject to comprehensive and long-term monitoring, yet it often remains unclear how those data feed into decision making (Dixon & Chiswell, 1996). In other instances, monitoring is haphazard or non-existent and it is likely that some freshwater species will be imperilled, or even extinct, before their existence is known (Burkhead, 2012). Major investments in different interventions (such as restoration) often occur with little or no monitoring of effectiveness (Cooke et al., 2018). Well-designed and -executed monitoring plans should feed directly into current and future management planning cycles. Many of the metrics currently used in conservation (e.g. habitat quality, species richness, species abundance) are inadequate to quantify biodiversity losses in freshwater habitats (Turak et al., 2017) and research is needed to improve monitoring metrics. Community science (also known as 'citizen science') can make a huge contribution to biodiversity monitoring (Chandler et al., 2017), but more work is needed to determine how this capacity can be enhanced for freshwater biodiversity and how different forms of knowledge (e.g. conventional science or traditional knowledge) can be blended in ways that are more comprehensive and strategically focused in relation to the aims and objectives of conservation and restoration efforts.

3.2 | Theme 2: Improving current practices

Questions in this theme identify gaps in current knowledge of measures to protect and restore freshwater biodiversity and ecosystems successfully. This includes the identification of high-priority biodiversity conservation areas, improvement of pollution control and remediation measures, identification of methods that proactively manage the effects of global change (e.g. species invasions) and the discovery of solutions to mitigate the effects of synergistic threats. The identification and application of these measures can enhance future action to bend the curve of freshwater biodiversity loss.

3.2.1 | Question 6: Key Biodiversity Areas: What are the Key Biodiversity Areas that need to be prioritized for conservation of freshwater biodiversity?

Key Biodiversity Areas are sites that contribute significantly to the global persistence of biodiversity (IUCN, 2016). Although recent research has contributed to the identification of Key Biodiversity Areas in the freshwater realm (Carrizo et al., 2017), more work is necessary to identify which attributes of these areas ensure the conservation of freshwater biodiversity. For example, catchments are recognized as useful planning and management units, but efforts to manage at catchment scales have often failed to prevent biodiversity

loss (Hermoso et al., 2016). In addition, determining which locations and species should be given conservation priority remains challenging (Whitehead et al., 2014), but should not be a barrier to conservation. Improving identification and protection of these areas is essential for biodiversity conservation.

3.2.2 | Question 7: Pollution: What approaches to pollution reduction and remediation efforts will most benefit freshwater biodiversity?

Point source and non-point source pollution continues to threaten freshwater ecosystem functions and biodiversity (Reid et al., 2019), necessitating better management and mitigation techniques both for groundwaters and surface waters. Stopping pollution at the source with better licensing and harm-reduction policies is essential, but it is equally important to find strategies for water resource management practitioners to meet their obligations and objectives once a pollutant is present. Reduction and remediation measures have been effectively applied to some freshwater systems (Søndergaard et al., 2007), but finding measures that will ensure long-term success continues to be a challenge for some pollutants, especially from non-point sources. With the identification of new pollutants (e.g. microplastics, pharmaceuticals), further research into improving existing techniques for pollution reduction and remediation is necessary. In addition, researching and adopting new measures (such as the use of 'nature-based solutions'; Liqueste et al., 2016) that are developed specifically for freshwater ecosystems could benefit freshwater biodiversity.

3.2.3 | Question 8: Tool development: What research innovations are needed to help restore freshwater biodiversity?

Understanding of freshwater ecosystem integrity and functioning has increased greatly over the past few decades. However, many threats to freshwater biodiversity are increasing in severity and frequency, while new threats continue to emerge (Reid et al., 2019). Leveraging new research techniques such as big data analytics, knowledge synthesis, community science or novel field techniques could advance conservation efforts (Cheruvilil & Soranno, 2018). Further developing techniques that allow decreased field work intensity (i.e. remote offload; Lennox et al., 2017) and approaches that do not require lethal sampling (e.g. environmental DNA, camera traps, remote sensing) is essential. Improving existing methods through facilitating longer-term field research (Mirtl et al., 2018) and study reproducibility (Fidler et al., 2017) or co-developing decision-support tools with conservation managers (Kuehne, Strecker & Olden, 2020) and community scientists could lead to the development of more effective conservation tools and initiatives. To be clear, this is not research for the acquisition of knowledge *per se*, but rather exploiting innovations in research for the meaningful advance of freshwater conservation.

3.2.4 | Question 9: Climate change: How do we proactively incorporate climate change adaptation into freshwater biodiversity conservation?

The effects of climate change continue to have severe impacts on freshwater ecosystems despite considerable research into the topic (e.g. the Fish Climate Change Database <https://fici.shinyapps.io/database/>; Krabbenhoft et al., 2020). It is essential that measures are used that enhance the resilience of freshwater systems to the effects of climate change (Huang et al., 2019). Understanding how to mitigate the impacts of climate change requires more evidence of the effectiveness of conservation strategies to support the functioning of freshwater ecosystems. For instance, some researchers advocate strategies that consider species vulnerability, exposure and adaptive capacity (Dawson et al., 2011) to improve the protection of freshwater habitats and species. Novel approaches could harness synergistic interactions where biodiversity gain arises from mitigation (e.g. carbon sequestration, reduced emissions), adaptation (e.g. restored riparian forest) and nature-based solutions (e.g. flood-risk management), but more evidence on their effectiveness is needed (Thomas, Griffiths & Ormerod, 2016).

3.2.5 | Question 10: Invasive species: What are the best ways to manage freshwater invasive species and diseases to ensure proactive and meaningful improvements to freshwater biodiversity?

The introduction and proliferation of invasive species and diseases in freshwater ecosystems can cause serious economic and conservation losses (Johnson & Paull, 2011; Pyšek et al., 2020). Unfortunately, these impacts are expected to become more extensive through new pathways such as easy access to invasive species through e-commerce (Peres et al., 2018) and a changing climate (Rahel & Olden, 2008). Although increasingly studied, knowledge of how to prevent and manage invasive species is often limited by insufficient information (Rytwinski et al., 2018). Strategies for better managing intentional introductions (e.g. fisheries enhancements for economic opportunities or vegetation control) that result in adverse impacts (Ellender et al., 2014) are needed to meet conservation goals. Although improving current control and prevention methods will be challenging, better understanding and communication of the impacts and management of invasive species will facilitate meaningful advances.

3.2.6 | Question 11: Riparian zones: What are the optimal riparian management actions that contribute to the protection of freshwater biodiversity?

Riparian areas, including floodplains, have long been regarded as important for freshwater ecosystems and a variety of management actions are used by practitioners to protect riparian areas and

adjacent fresh waters (Naiman, Decamps & McClain, 2010). Many questions remain regarding the importance of maintaining longitudinal continuity of riparian zones and lateral connectivity to floodplains, and the role of groundwater–riparian zone interactions on freshwater biodiversity. Riparian buffers and setbacks are common tools that have been shown to reduce flooding, limit erosion and protect aquatic and terrestrial habitats. Benefits could also arise for pollution reduction, thermal damping, enhanced energetic subsidies and habitat provision (Feld et al., 2018). Current guidelines on setback requirements and design criteria in some regions need further development and evaluation (Olugunorisa, 2009; Haley et al., 2016). Although setback widths are often defined by the size of the drainage area (National Research Council, 2000) and fixed-width buffers are standard practice (Richardson, Naiman & Bisson, 2012), more research is needed to determine the influence of landscape types on setback effectiveness. Defining best management practices and providing recommendations for riparian area and floodplain management could help protect freshwater biodiversity and freshwater ecosystem functioning.

3.2.7 | Question 12: Synergistic threats: How can we develop conservation and restoration measures that most effectively and efficiently address synergistic threats to freshwater biodiversity?

Multiple threats can lead to combined effects being greater than (synergism), less than (antagonism) or equal to the sum of (additive) their individual effects or can manifest in the opposite direction to independent effects (reversals). This can lead to unanticipated ecological responses (e.g. warming can reverse the trend of increasing phytoplankton biomass observed under cold acidification conditions; Christensen et al., 2006). A recent synthesis indicated that the net effects of paired alterations to freshwater ecosystems were more frequently antagonistic (41%) than synergistic (28%), additive (16%) or reversed (15%) (Jackson et al., 2016). Moreover, conservation projects targeting single threats often fail to address synergistic and additive effects (Craig et al., 2017). Given multiple and sometimes synergistic stressors, it is necessary to target limited resources so that the most significant stressors or threats are addressed and the most restorative blend of actions is identified.

3.3 | Theme 3: Balancing resource needs

There is a constant tension between human development and freshwater biodiversity conservation, especially in ecosystems where the high economic benefits gained by some groups through exploiting ecosystem resources is juxtaposed with the ecosystem management necessary to maintain biodiversity. Conventional approaches to economic development often focus on a narrow set of priorities at the cost of wider biodiversity (Flitcroft et al., 2019). This theme is focused on generating solutions that lead to positive outcomes for

freshwater biodiversity and for humans. Questions related to this theme include balancing resource extraction, sustainable food production and energy generation with the needs of freshwater biodiversity. Raising the priority of freshwater biodiversity and considering trade-offs in resource use and development will help in bending the curve and supporting wider sustainability in development outcomes.

3.3.1 | Question 13: Sustainable food: What are the joint priorities for sustainable food production and freshwater biodiversity conservation?

Demands from aquatic and terrestrial food production put pressure on freshwater ecosystems (e.g. through land-use conversion, over-exploitation, nutrient enrichment, pollution and water abstraction; Cottrell et al., 2018). Although efforts have been made to integrate terrestrial biodiversity within sustainable food systems (The Food and Land Use Coalition, 2019), less work has focused specifically on freshwater biodiversity. Freshwater conservation initiatives require integration with agriculture, aquaculture and inland fishery practices to minimize the adverse impacts of these pressures while providing food sustainability (Phang et al., 2019). Protecting freshwater biodiversity through the development and uptake of new methods in the food sector, such as alternative water sources (Intriago et al., 2018) or intensified production (Tanentzap et al., 2015), is challenging and sometimes controversial (e.g. balanced harvest; Zhou et al., 2019). These methods will be heavily influenced by geographical region and socio-economic context, so must be tailored to specific situations. Questions remain regarding the implementation of new techniques and harmonization of conservation and food-sustainability goals.

3.3.2 | Question 14: Dams and associated infrastructure: How can the need for dams and associated infrastructure be balanced with connectivity, health, and flow requirements of freshwater ecosystems and biodiversity?

Dams and associated infrastructure enable water storage, flood control and energy production, but are increasingly recognized as threats to freshwater ecosystems and biodiversity. Even small barriers and small hydropower plants can have damaging impacts on aquatic ecosystems (Couto & Olden, 2018; Lange et al., 2018; Belletti et al., 2020). There are growing calls to transform the use of dams to balance their benefits and costs and to address associated impacts and externalities more effectively during all phases of planning and design (Moran et al., 2018). Expanding energy portfolios to further develop alternative energy sources beyond hydropower will also lead to improved freshwater biodiversity outcomes. While there are some recent examples (Opperman et al., 2019; Hurford et al., 2020), there is a need for further research on how to assess trade-offs across social, environmental and economic variables (e.g. fisheries, agriculture and

hydropower; Pittock, Dumaresq & Orr, 2017). Additional research on the improvement of regulatory enforcement and site selection is necessary. Ensuring connectivity, improving operational flow regimes and incorporating freshwater biodiversity into policies affecting dam design and operation remain challenging but necessary (Poff & Olden, 2017).

3.3.3 | Question 15: Conflicting needs: How can we better balance conflicting interests between human demand for natural resources and freshwater biodiversity?

Conflicts between natural resource demands (e.g. groundwater and surface water abstraction for agriculture, industry, sanitation and domestic consumption, forestry, extraction of aggregates) and freshwater biodiversity will continue as human populations grow and per capita consumption increases (Motesharrei et al., 2016). Efficient consumption of resources that explicitly considers the protection of freshwater biodiversity and ecological limits is essential. Systematic approaches for freshwater conservation planning (Linke, Turak & Nel, 2011) and frameworks to improve decision-making in resource use (Huysman et al., 2015) could help balance these goals. However, shifts in economic practices (Martin, Maris & Simberloff, 2016), improved legislation and policy (Bringezu et al., 2016) and the development of new technologies (Czech, 2008) will probably be necessary to avoid many of the trade-offs to conservation gains. Promoting research on multidisciplinary solutions and applying limits in areas of current demand are important efforts to reduce risks to freshwater biodiversity.

3.4 | Theme 4: Rethinking built environments

This theme is representative of the increasing need to consider new avenues for freshwater biodiversity conservation such as in urban and suburban areas previously considered to be biodiversity poor. Questions relating to this concept aim at improving the recognition of opportunities and facilitating the development of programmes, policies and infrastructure that actively seek to incorporate freshwater biodiversity conservation to help expand understanding of valuable freshwater spaces. Considering indirect effects from infrastructure development (e.g. river aggregate extraction; Koehnken et al., 2020) and working to rethink and explicitly design infrastructure for freshwater conservation will facilitate bending the curve of freshwater biodiversity loss.

3.4.1 | Question 16: Urbanization: What policies, programmes and activities can we implement to turn the risks associated with urbanization into benefits/opportunities for freshwater biodiversity enhancement?

Frameworks for including biodiversity in urban development can mitigate the effects of urban growth and intensification

(e.g. Biodiversity Sensitive Urban Design; Garrard et al., 2018), but freshwater biodiversity has rarely been considered. Focusing on evaluating the persistence of freshwater species and ecosystems in development initiatives and capitalizing on opportunities realized during the development process can lead to improved outcomes (e.g. wetlands used for stormwater management in China's Sponge Cities; Chan et al., 2018). Influencing the distribution of people in cities to maximize species diversity is one possible strategy (Geschke et al., 2018). However, identifying ways to enable co-existence of humans and freshwater biodiversity through urban planning (Nel et al., 2008) and stormwater management (Hassall & Anderson, 2015) may be even more effective. These opportunities require the rethinking of targets and indicators (e.g. freshwater reptiles; Turak et al., 2020) in efforts to protect and improve urban biodiversity.

3.4.2 | Question 17: Infrastructure: How can freshwater biodiversity conservation be better integrated into infrastructure planning, implementation and operation?

Infrastructure development, including transportation, navigation, power, water supply, irrigation, stormwater management and sanitation, has generally proceeded without consideration for freshwater biodiversity. These activities can alter hydrology and ecosystems, adversely affecting freshwater biodiversity. Massive investments in water-associated infrastructure often fail to include sufficient expenditure to protect aquatic ecosystems (Bunn, 2016), but calls to consider ecosystems as infrastructure are increasing (da Silva & Wheeler, 2017). Determining how to change or replace current infrastructure and how infrastructure and biodiversity planning can be harmonized will lead to better cost-sharing approaches (Sleight & Neeson, 2018). Also needed is a greater understanding of how urban planning, building standards, construction supply chains, recycling and reuse of construction materials, and aggregate extraction practices can take better account of ecosystem impacts to maintain the health of many freshwater ecosystems. In addition, improving engineering strategies and planning for multiuse infrastructure enables the integration of resource use and freshwater biodiversity needs (e.g. planning irrigation with both agriculture and fisheries in mind; Lynch et al., 2019).

3.4.3 | Question 18: Novel/Designed ecosystems: What is the role of novel and designed ecosystems in conservation and how can these systems be managed to benefit freshwater biodiversity?

Novel ecosystems are self-assembling and self-sustaining and inadvertently arise through human activity (e.g. new wetlands following

peat harvesting; Collier, 2014), whereas designed ecosystems, such as retention ponds or large reservoirs, result from deliberate planning for human benefit and often require intensive intervention to maintain (Higgs, 2017). The contribution of novel and designed ecosystems to biodiversity conservation is unclear. Some argue that they allow flexible management of systems unlikely to return to historical conditions (e.g. 'designer' flows; Acreman et al., 2014); others argue that adopting these ecosystems may lead to de-prioritizing restoration activities (Miller & Bestelmeyer, 2016). It remains to be seen whether these ecosystems can provide suitable habitats for native species (but see Ebner, Lintermans & Dunford, 2011). Increased research will lead to new conservation opportunities (Heger et al., 2019).

3.5 | Theme 5: Reforming policy and investments

This theme highlights the growing need to implement and enforce strong policies that benefit freshwater biodiversity while recognizing the need for increased financial investments in freshwater conservation and restoration efforts. Policy and investment are both regionally and socio-economically dependent and must be addressed at the level of implementation in a targeted manner. Questions related to this theme aim at understanding what government structures and strategies are needed to implement change, as well as determining mechanisms to scale up and improve public and private sector financial investments for implementing specific conservation efforts. Effective policy and the identification of investment models for scaling up conservation financing can promote incentives that will ultimately lead to the protection of freshwater biodiversity.

3.5.1 | Question 19: Policy and legislation: What public policy measures can most effectively promote conservation and restoration of freshwater biodiversity?

Effective policy and legislation with a focus on freshwater ecosystems are necessary for future conservation efforts (Harrison et al., 2018; van Rees et al., 2020); however, conservation policy and legislation are often designed primarily for terrestrial or oceanic environments and do not fully account for the needs of freshwater ecosystems (Castello & Macedo, 2016). For example, freshwater biodiversity was not specifically mentioned in the United Nations Sustainable Development Goal (SDG) 14: 'Life Under Water' (United Nations, 2018), although many SDGs implicitly require conservation of fresh water (Lynch et al., 2017) and recent efforts show how freshwater fish and fisheries, for example, are integral to achieving the SDGs (Lynch et al., 2020). Understanding how to better account for environmental costs and consider trade-offs that favour solutions that benefit biodiversity, people and the economy would provide

major improvements in freshwater biodiversity policy. There is also a need to explore policy options related to incentivizing conservation actions that protect freshwater biodiversity and embracing nature-based solutions.

3.5.2 | Question 20: Financial investment: How can we scale up and optimize financial investments from government, private sector and other sources such that there is a step change in funding for global freshwater conservation and restoration efforts?

Although funding for conservation and restoration programmes has increased, there is a growing concern that consistent funding may not be available to support the long-term effectiveness of conservation efforts (Huwyler et al., 2014). Conservation financing has typically been provided on a small scale and investment opportunities remain underdeveloped. Generating economic and management benefits from conservation funding programmes and describing how they might create returns similar to traditional business models could provide a way forward (Huwyler et al., 2014). Highlighting improvements in efficiency, cost reductions and supply chain stability can support a solid business case for investment in conservation efforts by major corporations and insurance companies, among others (Clark, Reed & Sunderland, 2018). By identifying methods and incentives for scaling up financial investments and capitalizing on opportunities that reduce business risk, conservation financing could create significant contributions towards sustainable development and protection of freshwater biodiversity for the future.

3.5.3 | Question 21: Environmental flows: What are the social and natural science investments needed to develop and implement environmental flows that benefit freshwater biodiversity?

Knowledge of environmental flow requirements has improved, but implementation requires the continued collaboration of a variety of stakeholders, especially considering the diversity and interdependencies of human/flow relationships (Anderson et al., 2019). Collaboration could be enhanced by investments in social initiatives to improve support and increase understanding, and investments in the natural sciences to improve knowledge of effective environmental flow regimes. Continued research on incorporating environmental flows into policy and governance (Arthington et al., 2018) and creating mechanisms for their practical application is necessary. Setting reliable environmental flows, incorporating them into water management (i.e. at what scale; Opperman, Kendy & Barrios, 2019) and adapting flow-management strategies in the face of changing hydro-ecological conditions (Capon et al., 2018) will enable further improvements in environmental flows to support the needs of freshwater biodiversity.

3.5.4 | Question 22: Ex situ conservation: What type of investments in ex situ conservation (e.g. captive breeding, reintroduction, managed relocation) are most effective for imperilled freshwater biodiversity?

Despite attempts to conserve freshwater taxa *in situ*, increasing rates of habitat loss and climate change highlight the need for investments in alternative conservation tools (Olden et al., 2011; Brütting, Hensen & Wesche, 2013). *Ex situ* conservation is the process of conserving biological diversity at the gene, population and species level, outside the environment where it evolved. This technique can raise awareness of the plight of the species, but is expensive and requires extensive investments in time, tools and research. This is especially true given the number of imperilled freshwater organisms that need species-specific *ex situ* conservation strategies and the scale at which such efforts are needed (Snyder et al., 1996; Fischer & Lindenmayer, 2000). Identifying the most appropriate and cost-effective *ex situ* methods for different freshwater species, especially those with complex life cycles and unique ecosystem and habitat requirements (e.g. the development of an extensive captive breeding and reintroduction programme for Kihansi spray toads after the loss of their unique spray wetland; Lee et al., 2006) could lead to investments in *ex situ* conservation that create positive results for freshwater biodiversity restoration and improved technical guidelines for global cooperation.

3.6 | Theme 6: Enabling transformative change

This theme features research gaps that need to be addressed to enable transformative changes in individual human behaviour, societal actions and practice. Underpinning such efforts is the need to enhance knowledge exchange and raise awareness of the present state of freshwater biodiversity through better communication among researchers, between researchers and decision-makers, and between researchers and the general public. Questions relating to this theme include identifying methods to develop and enhance management frameworks for restoring biodiversity, sharing science and communicating findings, and increasing public engagement to lead to changes in individual behaviour to help bend the curve of freshwater biodiversity loss. Promoting better research practices could lead to improved conservation initiatives and, by translating these findings into more accessible forms, will increase public support and political will for restoring freshwater biodiversity.

3.6.1 | Question 23: Management frameworks: How do we develop management frameworks and evidence bases that gain greater traction with stakeholders and managers?

Conceptual management frameworks are tools by which complex systems, interactions and research gaps can be explained. Although

more recent frameworks (Millennium Ecosystem Assessment Board, 2005; Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services [IPBES], 2019) and a growing evidence base (Schreckenberg, Mace & Poudyal, 2018) have highlighted the strong linkages among freshwater biodiversity, human well-being, ecosystem services and government systems, active engagement by stakeholders and policy makers remains low. There remains a lack of empirical and targeted guidance for processes that consider complex dynamic interactions between these linkages. Related to this, guidance must necessarily be focused on a variety of different scales (geographically, socio-economically and in terms of governance) to reflect the context in which management decisions and conservation efforts are made. Frameworks for the management of freshwater biodiversity that not only foster evidence-based action, but also embed authentic participation by stakeholders and partners, are needed to realistically design and plan for conservation intervention (Langhans et al., 2019).

3.6.2 | Question 24: Science communication: What steps should be taken to better communicate and share evidence and knowledge about the science of freshwater biodiversity among stakeholders?

One of the key requirements for improving conservation of freshwater biodiversity is the establishment of stronger partnerships across sectors (Dudley et al., 2016). Building partnerships that create meaningful freshwater biodiversity outcomes requires effective communication between researchers, conservationists, practitioners, policymakers and the public. Using methods such as collaborative alliance models (Gray & Wood, 1991) or co-design would allow the integration of researchers and stakeholders in the planning and conduct of research on complex problems. This would improve the interpretation of results and the communication and use of findings. This can further be achieved by effectively translating scientific findings into material that is comprehensive, usable and accessible to other stakeholders. Communication among disparate knowledge-users requires enhancement and long-term maintenance of data-publishing and sharing platforms (Schmidt-Kloiber et al., 2019), improvement of evidence syntheses (Cooke et al., 2017) and the general implementation and acceptance of open-access publishing (Tennant et al., 2016) to ensure the availability of high-quality evidence.

3.6.3 | Question 25: Changing mindsets: How can we increase the level of public engagement to change mindsets and build social licence and political will to bend the curve of freshwater biodiversity loss?

Awareness of the current state of freshwater biodiversity among the general public remains low (Darwall et al., 2018). Engaging the public, and local political representatives, through community science,

environmental education (Sousa et al., 2016) or unique collaborations (e.g. with public aquariums; Murchie, Knapp & McIntyre, 2018) could result in improved understanding and willingness to support freshwater biodiversity initiatives. Changing attitudes and perspectives is difficult, especially if biodiversity initiatives are perceived as detrimental to human livelihoods (e.g. turtle bycatch reduction strategies; Nguyen et al., 2013), but is not impossible (Larocque et al., 2020). Designing methods to motivate involvement (e.g. community science activities) in environmental initiatives and to foster greater understanding and support for freshwater conservation will be challenging, and is likely to require long-term efforts and collaboration across the natural and social sciences. Increased public engagement and incorporation of diverse world views into these messages can raise the profile of freshwater biodiversity, leading to necessary actions directed towards improved conservation.

4 | DISCUSSION

In many areas of freshwater biodiversity conservation there is extensive evidence to demonstrate that actions to bend the curve must not be delayed. Conservation actions will be most effective when supported by sound evidence. If addressed comprehensively, the research questions presented here will fill critical knowledge gaps to better inform conservation activities and improve the effectiveness of present and future initiatives.

4.1 | Themes and questions

The six themes presented here are broadly applicable to many initiatives in freshwater biodiversity conservation. Although specific questions submitted by participants tended to have a narrower focus (see Table 2), they were collectively generalized into broader groups that cut across boundaries. The themes included: 1, learning from successes and failures; 2, improving current practices; 3, balancing resource needs; 4, rethinking built environments; 5, reforming policy and investment; and 6, enabling transformative change. One concept that connects all six themes is the need for interdisciplinary research, communication and collaboration with those beyond the freshwater conservation community. Examples of successful research efforts that have led to positive change for freshwater biodiversity highlight the effectiveness of these efforts (Boon & Baxter, 2020). There are many social science questions that can be asked for each of the research questions posed here (e.g. understanding barriers to change; Bennett et al., 2017a) and furthering research at the intersection of the natural and social sciences will only improve conservation outcomes, especially when paired with active and adaptive management as new knowledge becomes available.

The broad questions developed during this process tended to include concepts of proactive and meaningful development of policies, tools and metrics that would enhance and prioritize the effective management of freshwater biodiversity conservation initiatives at a

variety of spatial and temporal scales. In addition, they include a focus on scaling up investment and integrating various levels of research, public engagement and policy to balance priorities and provide optimal benefits for freshwater biodiversity and human needs. The 25 essential questions in this list provide starting points for identifying future research and a loose framework within which to prioritize more specific initiatives. The many cross-cutting and foundational issues contained in these questions (e.g. spatial scale, human behaviour) highlight how interconnected solutions and policies will be necessary in the future. The answers to these questions are not solely sufficient to bend the curve of freshwater biodiversity loss (Tickner et al., 2020) and these questions should by no means constrain research in other areas. We therefore call on the freshwater conservation community to continue to add new questions to this list, and to promote and implement recommended actions resulting from current or future research.

By our definition, bending the curve questions are those where answers will lead to actions for the recovery of freshwater biodiversity. Many of the submitted questions included calls to improve understanding of understudied regions and habitats (e.g. tropical ecosystems and non-perennial streams and wetlands), under-represented taxa (including macrophytes, algae, invertebrates and microbes) and emerging threats (e.g. invasive pathogens). These would, therefore, not directly produce the knowledge needed for changing the current trajectory of freshwater biodiversity loss. Furthermore, many of the original questions submitted were very specific to location or taxa. We recognize the importance of these types of questions to inform local-scale conservation and encourage the community to continue their efforts in these areas. Questions relating to these understudied topics are included in the complete list of submitted questions (see Supporting Information).

4.2 | Limitations

The call for questions attempted to reach the broadest possible audience, but there are limitations in the methodology. Despite being largely untargeted and freely available to anyone who wished to participate, the questionnaire was distributed only in English. Distribution through the professional and social networks of the authors probably limited its reach and accessibility to English-speaking nations and individuals. Most responses were received from Canada, the United States, Australia and other high-income nations (Table 1; Figure S2). As a result, the list of research questions may better reflect the interests of nations with well-developed conservation programmes, freshwater sciences and Western science perspectives. Many nations were represented by a single participant, resulting in a list of questions that may not have been adequately representative of broad geographical and socio-economic concerns. The lack of more comprehensive representation is likely to have influenced both the questions submitted and the resulting final list. Despite recruiting a diverse team of coauthors with regional, taxonomic and disciplinary expertise, the full diversity of research needs in freshwater biodiversity conservation

may not have been captured. To help mitigate this, any missing topics considered essential by the authors could be brought forward for consideration at other phases where themes are determined and refined. The relative importance of questions in this list will necessarily vary by geography, socio-economic and political conditions, knowledge systems and cultural norms. Our list is not intended to provide a specific road map, but rather to provide a list of potential areas to consider when establishing research agendas. We believe that providing this list is important for continuing conversations surrounding future actions for bending the curve.

Although attempts were made to reach out beyond research institutions, more responses were received from researchers (43%) compared with practitioners (20.5%) and decision-makers (12%). Students/post-docs and other roles make up the remaining 24.5% (Figure 1b). No responses were received from funders (Figure 1b). Since practitioners and decision-makers are less well represented in the responses, it is possible that questions seeking directly applicable solutions may not have been submitted. However, practitioners and decision-makers represent the experts on the ground in many regions and additional effort is needed to collate their experiences and knowledge to share with the broader community. Because practitioners may tend to maintain the status quo when engaging in conservation actions (Pullin & Knight, 2003; Nguyen, Young & Cooke, 2017), concerted efforts to disrupt these norms and ensure that work is founded on best available evidence will improve conservation outcomes (Sutherland et al., 2004; Cooke et al., 2017). Several new journals (e.g. *Ecological Solutions and Evidence*, *Conservation Science & Practice*) have been developed to provide mechanisms for practitioners to share their knowledge and findings at the interface between practical experience, management and theory, allowing for increased representation in research and decision-making. We encourage the community to utilize these and other avenues for increased knowledge sharing.

4.3 | Thinking globally

The implementation and enforcement of strong policies that benefit freshwater biodiversity are necessary both regionally and globally, and must be addressed in a targeted and equitable manner. Understanding the key role of freshwater biodiversity in contributing to ecosystem services is often overlooked at the international policy level. For instance, the 2020 Aichi biodiversity targets of the Convention on Biological Diversity (CBD) had no direct linkages to bending the curve for freshwater biodiversity (Tickner et al., 2020). The post-2020 framework for biodiversity, currently under negotiation at CBD, should ensure that there is an explicit goal focused on protection of freshwater biodiversity. Direct engagement in the discussion of the United Nations plan to protect 30% of the Earth's surface by 2030 (Dinerstein et al., 2019) at upcoming CBD plenaries focused on protecting freshwater systems will be important to ensure that fresh water is not ignored in selecting criteria for siting protected areas or developing targets to measure progress towards agreed goals.

Furthermore, to ensure that freshwater biodiversity research needs are identified, engagement of experts focused on aspects of freshwater biodiversity in initiatives such as the proposed assessments on the nexus between food, water, energy and health, and transformative change by IPBES will be important to highlight the importance of freshwater biodiversity (www.ipbes.net). Engagement with the climate community, through the Intergovernmental Panel on Climate Change, can help to ensure that science assessments focused on reducing carbon emissions will not have undue impacts on freshwater biodiversity as a trade-off for increased energy development.

5 | CONCLUSION

Our aspiration is that the essential questions presented here will serve as a springboard for multidisciplinary and multisectoral collaborations that succeed in tackling the challenges of the freshwater biodiversity crisis. Bold, efficient, science-based actions are necessary to halt and reverse biodiversity loss (Mace et al., 2018), especially for freshwater biodiversity (Tickner et al., 2020). Addressing many of the research questions listed here will require the allocation of significant resources, but not all questions need to be addressed in all regions. Regional priorities need to be developed and funding strategies identified, which will require coordinated efforts from key non-governmental organizations, governments and communities (including rights- and stakeholders). The extensive focus on social sciences and policy in these questions showcases the need for collaboration and multi- and trans-disciplinary efforts that bridge the gap between research, public participation and policy. Targeted, multidisciplinary research funding will enhance urgent efforts to protect the world's freshwater biodiversity by making conservation and restoration efforts more effective and applicable at scale. In addition, global syntheses emerging from distributed empirical research will also be needed to enable evidence-based decision-making. Conservation actions will be most effective when supported by sound evidence, but we are also emphatic that action should not be delayed in the face of uncertainty (O'Riordan & Cameron, 1994; Rytwinski et al., 2021). The themes and questions presented here help to highlight current research needs in freshwater biodiversity conservation. Addressing these questions comprehensively is achievable and necessary.

ACKNOWLEDGEMENTS

This effort was inspired by the long-term goal of the Alliance for Freshwater Life (<https://allianceforfreshwaterlife.org>) to achieve a better understanding of freshwater biodiversity decline and developing solutions to reverse biodiversity loss (Darwall et al., 2018). The authors are grateful for the engagement of members of The Alliance for Freshwater Life, for helpful comments from Candace M. Hansen-Hendriks on the original essential questions and for comments from two anonymous reviewers that greatly improved this manuscript. We also thank the freshwater conservation community for their involvement in this project. Funding was provided by National Sciences and Engineering Research Council of Canada. This is a product of the

Canadian Centre for Evidence-Based Conservation. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the US Government. We dedicate this paper to friend, colleague and co-author Dr Olaf Weyl, who passed away while this paper was in review. Olaf was a champion for freshwater biodiversity and a founding member of the Alliance for Freshwater Life. We commit to continuing the work of Olaf with the same passion, enthusiasm and excellence that Olaf brought to everything he did.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

ORCID

Meagan Harper  <https://orcid.org/0000-0002-8462-2039>
 Hebah S. Mejbil  <https://orcid.org/0000-0001-5737-7608>
 Robin Abell  <https://orcid.org/0000-0001-9407-7807>
 T. Douglas Beard  <https://orcid.org/0000-0003-2632-2350>
 Joseph R. Bennett  <https://orcid.org/0000-0002-3901-9513>
 Stephanie M. Carlson  <https://orcid.org/0000-0003-3055-6483>
 William Darwall  <https://orcid.org/0000-0002-4589-5579>
 Sami Domisch  <https://orcid.org/0000-0002-8127-9335>
 David Dudgeon  <https://orcid.org/0000-0003-4632-3473>
 Jörg Freyhof  <https://orcid.org/0000-0002-7042-3127>
 Ian Harrison  <https://orcid.org/0000-0001-8686-8502>
 Sonja C. Jähnig  <https://orcid.org/0000-0002-6349-9561>
 Harmony Patricia  <https://orcid.org/0000-0002-3565-6881>
 Andrea J. Reid  <https://orcid.org/0000-0001-7251-7824>
 Astrid Schmidt-Kloiber  <https://orcid.org/0000-0001-8839-5913>
 Michele Thieme  <https://orcid.org/0000-0003-3216-9129>
 Eren Turak  <https://orcid.org/0000-0001-7383-9112>
 Steven J. Cooke  <https://orcid.org/0000-0002-5407-0659>

REFERENCES

- Acreman, M., Arthington, A.H., Colloff, M.J., Couch, C., Crossman, N.D., Dyer, F. et al. (2014). Environmental flows for natural, hybrid, and novel riverine ecosystems in a changing world. *Frontiers in Ecology and the Environment*, 12(8), 466–473. <https://doi.org/10.1890/130134>
- Acreman, M., Hughes, K.A., Arthington, A.H., Tickner, D. & Dueñas, M.A. (2019). Protected areas and freshwater biodiversity: A novel systematic review distils eight lessons for effective conservation. *Conservation Letters*, 13(1), e12684. <https://doi.org/10.1111/conl.12684>
- Anderson, E.P., Jackson, S., Tharme, R.E., Douglas, M., Flotemersch, J.E., Zwartveen, M. et al. (2019). Understanding rivers and their social relations: A critical step to advance environmental water management. *WIREs Water*, 6(6), e1381. <https://doi.org/10.1002/wat2.1381>
- Arthington, A.H., Bhaduri, A., Bunn, S.E., Jackson, S.E., Tharme, R.E., Tickner, D. et al. (2018). The Brisbane Declaration and global action agenda on environmental flows (2018). *Frontiers in Environmental Science*, 6, 45. <https://doi.org/10.3389/fenvs.2018.00045>
- Belletti, B., Garcia de Leaniz, C., Jones, J., Bizzi, S., Börger, L., Segura, G. et al. (2020). More than one million barriers fragment Europe's rivers. *Nature*, 588(7838), 436–441. <https://doi.org/10.1038/s41586-020-3005-2>
- Bennett, E.M., Solan, M., Biggs, R., McPhearson, T., Norström, A.V., Olsson, P. et al. (2016). Bright spots: Seeds of a good Anthropocene.

- Frontiers in Ecology and the Environment*, 14(8), 441–448. <https://doi.org/10.1002/fee.1309>
- Bennett, N.J., Roth, R., Klain, S.C., Chan, K., Christie, P., Clark, D.A. et al. (2017a). Conservation social science: Understanding and integrating human dimensions to improve conservation. *Biological Conservation*, 205, 93–108. <https://doi.org/10.1016/j.biocon.2016.10.006>
- Bennett, N.J., Roth, R., Klain, S.C., Chan, K.M., Clark, D.A., Cullman, G. et al. (2017b). Mainstreaming the social sciences in conservation. *Conservation Biology*, 31(1), 56–66. <https://doi.org/10.1111/cobi.12788>
- BioFresh. (2011). *Summary of responses to 'top questions facing freshwater biodiversity science, policy and conservation'*. Montserrat, Spain: BioFresh Project Meeting.
- Birk, S., Chapman, D., Carvalho, L., Spears, B.M., Andersen, H.E., Argillier, C. et al. (2020). Impacts of multiple stressors on freshwater biota across spatial scales and ecosystems. *Nature Ecology & Evolution*, 4(8), 1060–1068. <https://doi.org/10.1038/s41559-020-1216-4>
- Boon, P.J. & Baxter, J.M. (2016). Aquatic conservation: Reflections on the first 25 years. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 26(5), 809–816. <https://doi.org/10.1002/aqc.2713>
- Boon, P.J. & Baxter, J.M. (Eds.) (2020). Aquatic Conservation in action: Demonstrating the practical impact of the journal's publications. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 30(9). [Special issue].
- Bringezu, S., Potočník, J., Schandl, H., Lu, Y., Ramaswami, A., Swilling, M. et al. (2016). Multi-scale governance of sustainable natural resource use – Challenges and opportunities for monitoring and institutional development at the national and global level. *Sustainability*, 8(8), 778. <https://doi.org/10.3390/su8080778>
- Brütting, C., Hensen, I. & Wesche, K. (2013). Ex situ cultivation affects genetic structure and diversity in arable plants. *Plant Biology*, 15(3), 505–513. <https://doi.org/10.1111/j.1438-8677.2012.00655.x>
- Bunn, S.E. (2016). Grand challenge for the future of freshwater ecosystems. *Frontiers in Environmental Science*, 4, 21. <https://doi.org/10.3389/fenvs.2016.00021>
- Burkhead, N.M. (2012). Extinction rates in North American freshwater fishes, 1900–2010. *Bioscience*, 62(9), 798–808. <https://doi.org/10.1525/bio.2012.62.9.5>
- Capon, S.J., Leigh, C., Hadwen, W.L., George, A., McMahon, J.M., Linke, S. et al. (2018). Transforming environmental water management to adapt to a changing climate. *Frontiers in Environmental Science*, 6, 80. <https://doi.org/10.3389/fenvs.2018.00080>
- Carrizo, S.F., Lengyel, S., Kapusi, F., Szabolcs, M., Kasperidus, H.D., Scholz, M. et al. (2017). Critical catchments for freshwater biodiversity conservation in Europe: Identification, prioritisation and gap analysis. *Journal of Applied Ecology*, 54(4), 1209–1218. <https://doi.org/10.1111/1365-2664.12842>
- Castello, L. & Macedo, M.N. (2016). Large-scale degradation of Amazonian freshwater ecosystems. *Global Change Biology*, 22(3), 990–1007. <https://doi.org/10.1111/gcb.13173>
- Chan, F.K.S., Griffiths, J.A., Higgitt, D., Xu, S., Zhu, F., Tang, Y.T. et al. (2018). 'Sponge City' in China – A breakthrough of planning and flood risk management in the urban context. *Land Use Policy*, 76, 772–778. <https://doi.org/10.1016/j.landusepol.2018.03.005>
- Chandler, M., See, L., Copas, K., Bonde, A.M.Z., López, B.C., Danielsen, F. et al. (2017). Contribution of citizen science towards international biodiversity monitoring. *Biological Conservation*, 213(B), 280–294. <https://doi.org/10.1016/j.biocon.2016.09.004>
- Cheruvilil, K.S. & Soranno, P.A. (2018). Data-intensive ecological research is catalyzed by open science and team science. *Bioscience*, 68(10), 813–822. <https://doi.org/10.1093/biosci/biy097>
- Christensen, M.R., Graham, M.D., Vinebrooke, R.D., Findlay, D.L., Paterson, M.J. & Turner, M.A. (2006). Multiple anthropogenic stressors cause ecological surprises in boreal lakes. *Global Change Biology*, 12(12), 2316–2322. <https://doi.org/10.1111/j.1365-2486.2006.01257.x>
- Clark, R., Reed, J. & Sunderland, T. (2018). Bridging funding gaps for climate and sustainable development: Pitfalls, progress and potential of private finance. *Land Use Policy*, 71, 335–346. <https://doi.org/10.1016/j.landusepol.2017.12.013>
- Collier, M.J. (2014). Novel ecosystems and the emergence of cultural ecosystem services. *Ecosystem Services*, 9, 166–169. <https://doi.org/10.1016/j.ecoser.2014.06.002>
- Cooke, S.J., Rous, A.M., Donaldson, L.A., Taylor, J.J., Rytwinski, T., Prior, K. A. et al. (2018). Evidence-based restoration in the Anthropocene – From acting with purpose to acting for impact. *Restoration Ecology*, 26(2), 201–205. <https://doi.org/10.1111/rec.12675>
- Cooke, S.J., Wesch, S., Donaldson, L.A., Wilson, A.D.M. & Haddaway, N.R. (2017). A call for evidence-based conservation and management of fisheries and aquatic resources. *Fisheries*, 42(3), 143–149. <https://doi.org/10.1080/03632415.2017.1276343>
- Cottrell, R.S., Fleming, A., Fulton, E.A., Nash, K.L., Watson, R.A. & Blanchard, J.L. (2018). Considering land–sea interactions and trade-offs for food and biodiversity. *Global Change Biology*, 24(2), 580–596. <https://doi.org/10.1111/gcb.13873>
- Couto, T.B. & Olden, J.D. (2018). Global proliferation of small hydropower plants – Science and policy. *Frontiers in Ecology and the Environment*, 16(2), 91–100. <https://doi.org/10.1002/fee.1746>
- Craig, L.S., Olden, J.D., Arthington, A.H., Entekin, S., Hawkins, C.P., Kelly, J.J. et al. (2017). Meeting the challenge of interacting threats in freshwater ecosystems: A call to scientists and managers. *Elementa: Science of the Anthropocene*, 5, 72. <https://doi.org/10.1525/elementa.256>
- Czech, B. (2008). Prospects for reconciling the conflict between economic growth and biodiversity conservation with technological progress. *Conservation Biology*, 22(6), 1389–1398. <https://doi.org/10.1111/j.1523-1739.2008.01089.x>
- da Silva, J.M.C. & Wheeler, E. (2017). Ecosystems as infrastructure. *Perspectives in Ecology and Conservation*, 15(1), 32–35. <https://doi.org/10.1016/j.pecon.2016.11.005>
- Darwall, W., Bremerich, V., Wever, A.D., Dell, A.I., Freyhof, J., Gessner, M.O. et al. (2018). The Alliance for Freshwater Life: A global call to unite efforts for freshwater biodiversity science and conservation. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 28(4), 1015–1022. <https://doi.org/10.1002/aqc.2958>
- Dawson, T.P., Jackson, S.T., House, J.I., Prentice, I.C. & Mace, G.M. (2011). Beyond predictions: Biodiversity conservation in a changing climate. *Science*, 332(6025), 53–58. <https://doi.org/10.1126/science.1200303>
- Dinerstein, E., Vynne, C., Sala, E., Joshi, A.R., Fernando, S., Lovejoy, T.E. et al. (2019). A global deal for nature: Guiding principles, milestones, and targets. *Science Advances*, 5(4), eaaw2869. <https://doi.org/10.1126/sciadv.aaw2869>
- Dixon, W. & Chiswell, B. (1996). Review of aquatic monitoring program design. *Water Research*, 30(9), 1935–1948. [https://doi.org/10.1016/0043-1354\(96\)00087-5](https://doi.org/10.1016/0043-1354(96)00087-5)
- Dudgeon, D., Arthington, A.H., Gessner, M.O., Kawabata, Z.I., Knowler, D.J., Lévêque, C. et al. (2006). Freshwater biodiversity: Importance, threats, status and conservation challenges. *Biological Reviews*, 81(2), 163–182. <https://doi.org/10.1017/S1464793105006950>
- Dudley, N., Harrison, I.J., Kettunen, M., Madgwick, J. & Mauerhofer, V. (2016). Natural solutions for water management of the future: Freshwater protected areas at the 6th World Parks Congress. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 26(S1), 121–132. <https://doi.org/10.1002/aqc.2657>
- Ebner, B.C., Lintermans, M. & Dunford, M. (2011). A reservoir serves as refuge for adults of the endangered Macquarie perch. *Lakes & Reservoirs: Science, Policy and Management for Sustainable Use*, 16(1), 23–33. <https://doi.org/10.1111/j.1440-1770.2011.00463.x>

- Ellender, B.R., Woodford, D.J., Weyl, O.L.F. & Cowx, I.G. (2014). Managing conflicts arising from fisheries enhancements based on non-native fishes in southern Africa. *Journal of Fish Biology*, 85(6), 1890–1906. <https://doi.org/10.1111/jfb.12512>
- European Commission. (2011). Our Life Insurance, our Natural Capital: An EU Biodiversity Strategy to 2020. Available at: <https://www.eea.europa.eu/data-and-maps/indicators/plant-phenology-1/european-commission-2011-our-life> [Accessed February 16, 2020]
- Feld, C.K., Fernandes, M.R., Ferreira, M.T., Hering, D., Ormerod, S.J., Venohr, M. et al. (2018). Evaluating riparian solutions to multiple stressor problems in river ecosystems – A conceptual study. *Water Research*, 139, 381–394. <https://doi.org/10.1016/j.watres.2018.04.014>
- Fidler, F., Chee, Y.E., Wintle, B.C., Burgman, M.A., McCarthy, M.A. & Gordon, A. (2017). Metaresearch for evaluating reproducibility in ecology and evolution. *Bioscience*, 67(3), 282–289. <https://doi.org/10.1093/biosci/biw159>
- Fischer, J. & Lindenmayer, D.B. (2000). An assessment of the published results of animal relocations. *Biological Conservation*, 96(1), 1–11. [https://doi.org/10.1016/S0006-3207\(00\)00048-3](https://doi.org/10.1016/S0006-3207(00)00048-3)
- Flitcroft, R., Cooperman, M.S., Harrison, I.J., Juffe-Bignoli, D. & Boon, P.J. (2019). Theory and practice to conserve freshwater biodiversity in the Anthropocene. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 29(7), 1013–1021. <https://doi.org/10.1002/aqc.3187>
- García-Moreno, J., Harrison, I.J., Dudgeon, D., Clausnitzer, V., Darwall, W., Farrell, T. et al. (2014). Sustaining freshwater biodiversity in the Anthropocene. In: A. Bhaduri, J. Bogardi, J. Leentvaar, S. Marx (Eds.) *The global water system in the Anthropocene: Challenges for science and governance*. Cham: Springer, pp. 247–270. https://doi.org/10.1007/978-3-319-07548-8_17
- Garrard, G.E., Williams, N.S.G., Mata, L., Thomas, J. & Bekessy, S.A. (2018). Biodiversity sensitive urban design. *Conservation Letters*, 11(2), e12411. <https://doi.org/10.1111/conl.12411>
- Geschke, A., James, S., Bennett, A.F. & Nimmo, D.G. (2018). Compact cities or sprawling suburbs? Optimising the distribution of people in cities to maximise species diversity. *Journal of Applied Ecology*, 55(5), 2320–2331. <https://doi.org/10.1111/1365-2664.13183>
- Gray, B. & Wood, D.J. (1991). Collaborative alliances: Moving from practice to theory. *The Journal of Applied Behavioral Science*, 27(1), 3–22. <https://doi.org/10.1177/0021886391271001>
- Haley, H., McCawley, M., Epstein, A.C., Arrington, B. & Ferrell Bjerke, E. (2016). Adequacy of current state setbacks for directional high-volume hydraulic fracturing in the Marcellus, Barnett, and Niobrara shale plays. *Environmental Health Perspectives*, 124(9), 1323–1333. <https://doi.org/10.1289/ehp.1510547>
- Harrison, I., Abell, R., Darwall, W., Thieme, M.L., Tickner, D. & Timboe, I. (2018). The freshwater biodiversity crisis. *Science*, 362(6421), 1369–1369. <https://doi.org/10.1126/science.aav9242>
- Hassall, C. & Anderson, S. (2015). Stormwater ponds can contain comparable biodiversity to unmanaged wetlands in urban areas. *Hydrobiologia*, 745(1), 137–149. <https://doi.org/10.1007/s10750-014-2100-5>
- Heger, T., Bernard-Verdier, M., Gessler, A., Greenwood, A.D., Grossart, H.P., Hilker, M. et al. (2019). Towards an integrative, eco-evolutionary understanding of ecological novelty: Studying and communicating interlinked effects of global change. *Bioscience*, 69(11), 888–899. <https://doi.org/10.1093/biosci/biz095>
- Hermoso, V., Abell, R., Linke, S. & Boon, P. (2016). The role of protected areas for freshwater biodiversity conservation: Challenges and opportunities in a rapidly changing world. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 26(51), 3–11. <https://doi.org/10.1002/aqc.2681>
- Higgs, E. (2017). Novel and designed ecosystems. *Restoration Ecology*, 25(1), 8–13. <https://doi.org/10.1111/rec.12410>
- Huang, L., Liao, F.H., Lohse, K.A., Larson, D.M., Fragkias, M., Lybecker, D.L. et al. (2019). Land conservation can mitigate freshwater ecosystem services degradation due to climate change in a semiarid catchment: The case of the Portneuf River catchment, Idaho, USA. *Science of the Total Environment*, 651(2), 1796–1809. <https://doi.org/10.1016/j.scitotenv.2018.09.260>
- Hurford, A.P., McCartney, M.P., Harou, J.J., Dalton, J., Smith, D.M. & Odada, E. (2020). Balancing services from built and natural assets via river basin trade-off analysis. *Ecosystem Services*, 45, 101144. <https://doi.org/10.1016/j.ecoser.2020.101144>
- Huwylar, F., Kappeli, J., Serafimova, K., Swanson, E. & Tobin, J. (2014). *Conservation finance: Moving beyond donor funding toward an investor-driven approach*. Washington, DC: Credit Suisse, World Wildlife Fund and McKinsey, pp. 1–32.
- Huysman, S., Sala, S., Mancini, L., Ardente, F., Alvarenga, R.A.F., De Meester, S. et al. (2015). Toward a systematized framework for resource efficiency indicators. *Resources, Conservation and Recycling*, 95, 68–76. <https://doi.org/10.1016/j.resconrec.2014.10.014>
- Hynes, H.B.N. (1975). The stream and its valley. *Verhandlungen der Internationalen Vereinigung für theoretische und angewandte Limnologie*, 19(1), 1–15. <https://doi.org/10.1080/03680770.1974.11896033>
- Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services [IPBES]. (2019). Summary for policymakers of the global assessment report on biodiversity and ecosystem services. Zenodo. <https://doi.org/10.5281/zenodo.3553579>
- International Union for Conservation of Nature [IUCN]. (2012). *Red list categories and criteria: Version 3.1*, Second edition. Gland, Switzerland: IUCN. Available at: <https://portals.iucn.org/library/node/10315>
- International Union for Conservation of Nature [IUCN]. (2016). *A global standard for the identification of key biodiversity areas: Version 1.0*, First edition. Gland, Switzerland: IUCN. Available at: <https://portals.iucn.org/library/node/46259>
- Intriago, J.C., López-Gálvez, F., Allende, A., Vivaldi, G.A., Camposeo, S., Nicolás, E. et al. (2018). Agricultural reuse of municipal wastewater through an integral water reclamation management. *Journal of Environmental Management*, 213, 135–141. <https://doi.org/10.1016/j.jenvman.2018.02.011>
- Jackson, M.C., Loewen, C.J.G., Vinebrooke, R.D. & Chimimba, C.T. (2016). Net effects of multiple stressors in freshwater ecosystems: A meta-analysis. *Global Change Biology*, 22(1), 180–189. <https://doi.org/10.1111/gcb.13028>
- Jaeger, K.L., Olden, J.D. & Pelland, N.A. (2014). Climate change poised to threaten hydrologic connectivity and endemic fishes in dryland streams. *Proceedings of the National Academy of Sciences*, 111(38), 13894–13899. <https://doi.org/10.1073/pnas.1320890111>
- Jähnig, S., Arlinghaus, R., Becks, L., Behrmann-Godel, J., Berendonk, T., Borchardt, D. et al. (2019). Living waters: A research agenda for the biodiversity of inland coastal waters. Germany: Research Initiative for the Conservation of Biodiversity. Available at: <https://repository.publisso.de/resource/frl:6418180>
- Johnson, P.T.J. & Paull, S.H. (2011). The ecology and emergence of diseases in fresh waters. *Freshwater Biology*, 56(4), 638–657. <https://doi.org/10.1111/j.1365-2427.2010.02546.x>
- Kalinkat, G., Cabral, J.S., Darwall, W., Ficetola, G.F., Fisher, J.L., Giling, D.P. et al. (2017). Flagship umbrella species needed for the conservation of overlooked aquatic biodiversity. *Conservation Biology*, 31(2), 481–485. <https://doi.org/10.1111/cobi.12813>
- Knight, A.T. (2006). Failing but learning: Writing the wrongs after Redford and Taber. *Conservation Biology: The Journal of the Society for Conservation Biology*, 20(4), 1312–1314. <https://doi.org/10.1111/j.1523-1739.2006.00366.x>
- Koehnken, L., Rintoul, M.S., Goichot, M., Tickner, D., Loftus, A.C. & Acreman, M.C. (2020). Impacts of riverine sand mining on freshwater ecosystems: A review of the scientific evidence and guidance for

- future research. *River Research and Applications*, 36(3), 362–370. <https://doi.org/10.1002/rra.3586>
- Koning, A.A., Perales, K.M., Fluet-Chouinard, E. & McIntyre, P.B. (2020). A network of grassroots reserves protects tropical river fish diversity. *Nature*, 588(7839), 631–635. <https://doi.org/10.1038/s41586-020-2944-y>
- Krabbenhoft, T.J., Myers, B.J.E., Wong, J.P., Chu, C., Tingley, R.W., Falke, J.A. et al. (2020). FiCli, the Fish and Climate Change Database, informs climate adaptation and management for freshwater fishes. *Scientific Data*, 7(1), 1–6. <https://doi.org/10.1038/s41597-020-0465-z>
- Kuehne, L.M., Strecker, A.L. & Olden, J.D. (2020). Knowledge exchange and social capital for freshwater ecosystem assessments. *Bioscience*, 70(2), 174–183. <https://doi.org/10.1093/biosci/biz142>
- Lange, K., Meier, P., Trautwein, C., Schmid, M., Robinson, C.T., Weber, C. et al. (2018). Basin-scale effects of small hydropower on biodiversity dynamics. *Frontiers in Ecology and the Environment*, 16(7), 397–404. <https://doi.org/10.1002/fee.1823>
- Langhans, S.D., Domisch, S., Balbi, S., Delacámara, G., Hermoso, V., Kuemmerlen, M. et al. (2019). Combining eight research areas to foster the uptake of ecosystem-based management in fresh waters. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 29(7), 1161–1173. <https://doi.org/10.1002/aqc.3012>
- Lant, T.K. & Montgomery, D.B. (1987). Learning from strategic success and failure. *Journal of Business Research*, 15(6), 503–517. [https://doi.org/10.1016/0148-2963\(87\)90035-X](https://doi.org/10.1016/0148-2963(87)90035-X)
- Larocque, S.M., Lake, C., Midwood, J.D., Nguyen, V.M., Blouin-Demers, G. & Cooke, S.J. (2020). Freshwater turtle bycatch research supports science-based fisheries management. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 30(9), 1783–1790. <https://doi.org/10.1002/aqc.3404>
- Lee, S., Zippel, K., Ramos, L. & Searle, J. (2006). Captive-breeding programme for the Kihansi spray toad *Nectophrynoides asperginis* at the Wildlife Conservation Society, Bronx, New York. *International Zoo Yearbook*, 40(1), 241–253. <https://doi.org/10.1111/j.1748-1090.2006.00241.x>
- Lennox, R.J., Aarestrup, K., Cooke, S.J., Cowley, P.D., Deng, Z.D., Fisk, A.T. et al. (2017). Envisioning the future of aquatic animal tracking: Technology, science, and application. *Bioscience*, 67(10), 884–896. <https://doi.org/10.1093/biosci/bix098>
- Lennox, R.J., Paukert, C.P., Aarestrup, K., Auger-Méthé, M., Baumgartner, L., Birnie-Gauvin, K. et al. (2019). One hundred pressing questions on the future of global fish migration science, conservation, and policy. *Frontiers in Ecology and Evolution*, 7, 286. <https://doi.org/10.3389/fevo.2019.00286>
- Li, B.V. & Pimm, S.L. (2016). China's endemic vertebrates sheltering under the protective umbrella of the giant panda. *Conservation Biology*, 30(2), 329–339. <https://doi.org/10.1111/cobi.12618>
- Linke, S., Turak, E. & Nel, J. (2011). Freshwater conservation planning: The case for systematic approaches. *Freshwater Biology*, 56(1), 6–20. <https://doi.org/10.1111/j.1365-2427.2010.02456.x>
- Lintermans, M. (2013). A review of on-ground recovery actions for threatened freshwater fish in Australia. *Marine and Freshwater Research*, 64(9), 775–791. <https://doi.org/10.1071/MF12306>
- Liquete, C., Udias, A., Conte, G., Grizzetti, B. & Masi, F. (2016). Integrated valuation of a nature-based solution for water pollution control. Highlighting hidden benefits. *Ecosystem Services*, 22(B), 392–401. <https://doi.org/10.1016/j.ecoser.2016.09.011>
- Loury, E.K., Ainsley, S.M., Bower, S.D., Chuenpagdee, R., Farrell, T., Guthrie, A.G. et al. (2018). Salty stories, fresh spaces: Lessons for aquatic protected areas from marine and freshwater experiences. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 28(2), 485–500. <https://doi.org/10.1002/aqc.2868>
- Lynch, A.J., Baumgartner, L.J., Boys, C.A., Conallin, J., Cowx, I.G., Finlayson, C.M. et al. (2019). Speaking the same language: Can the sustainable development goals translate the needs of inland fisheries into irrigation decisions? *Marine and Freshwater Research*, 70(9), 1211–1228. <https://doi.org/10.1071/MF19176>
- Lynch, A.J., Cowx, I.G., Fluet-Chouinard, E., Glaser, S.M., Phang, S.C., Beard, T.D. et al. (2017). Inland fisheries – Invisible but integral to the UN Sustainable Development Agenda for ending poverty by 2030. *Global Environmental Change*, 47, 167–173. <https://doi.org/10.1016/j.gloenvcha.2017.10.005>
- Lynch, A.J., Elliott, V., Phang, S.C., Claussen, J.E., Harrison, I., Murchie, K.J. et al. (2020). Inland fish and fisheries integral to achieving the Sustainable Development Goals. *Nature Sustainability*, 3(8), 579–587. <https://doi.org/10.1038/s41893-020-0517-6>
- Mace, G.M., Barrett, M., Burgess, N.D., Cornell, S.E., Freeman, R., Grooten, M. et al. (2018). Aiming higher to bend the curve of biodiversity loss. *Nature Sustainability*, 1(9), 448–451. <https://doi.org/10.1038/s41893-018-0130-0>
- Martin, J.L., Maris, V. & Simberloff, D.S. (2016). The need to respect nature and its limits challenges society and conservation science. *Proceedings of the National Academy of Sciences*, 113(22), 6105–6112. <https://doi.org/10.1073/pnas.1525003113>
- Mascarenhas, M. (2007). Where the waters divide: First Nations, tainted water and environmental justice in Canada. *Local Environment*, 12(6), 565–577. <https://doi.org/10.1080/13549830701657265>
- McGowan, J., Beaumont, L.J., Smith, R.J., Chauvenet, A.L.M., Harcourt, R., Atkinson, S.C. et al. (2020). Conservation prioritization can resolve the flagship species conundrum. *Nature Communications*, 11(1), 1–7. <https://doi.org/10.1038/s41467-020-14554-z>
- Millennium Ecosystem Assessment Board [MA]. (2005). *Ecosystems and human well-being: Current state and trends*, Vol. 1. Washington: Island Press. Available at: <https://www.millenniumassessment.org/en/Global.html>
- Miller, J.R. & Bestelmeyer, B.T. (2016). What's wrong with novel ecosystems, really? *Restoration Ecology*, 24(5), 577–582. <https://doi.org/10.1111/rec.12378>
- Mirtl, M., Borer, E.T., Djukic, I., Forsius, M., Haubold, H., Hugo, W. et al. (2018). Genesis, goals and achievements of Long-Term Ecological Research at the global scale: A critical review ofILTER and future directions. *Science of the Total Environment*, 626, 1439–1462. <https://doi.org/10.1016/j.scitotenv.2017.12.001>
- Moran, E.F., Lopez, M.C., Moore, N., Müller, N. & Hyndman, D.W. (2018). Sustainable hydropower in the 21st century. *Proceedings of the National Academy of Sciences of the United States of America*, 115(47), 11891–11898. <https://doi.org/10.1073/pnas.1809426115>
- Motesharrei, S., Rivas, J., Kalnay, E., Asrar, G.R., Busalacci, A.J., Cahalan, R.F. et al. (2016). Modeling sustainability: Population, inequality, consumption, and bidirectional coupling of the Earth and human systems. *National Science Review*, 3(4), 470–494. <https://doi.org/10.1093/nsr/nww081>
- Murchie, K.J., Knapp, C.R. & McIntyre, P.B. (2018). Advancing freshwater biodiversity conservation by collaborating with public aquaria. *Fisheries*, 43(4), 172–178. <https://doi.org/10.1002/fsh.10056>
- Naiman, R.J., Decamps, H. & McClain, M.E. (2010). *Riparia: Ecology, conservation, and management of streamside communities*. Burlington, MA: Elsevier.
- National Research Council. (2000). *Watershed management for potable water supply: Assessing the New York City strategy*. Washington DC: The National Academies Press. <https://doi.org/10.17226/9677>
- Nel, J.L., Roux, D.J., Abell, R., Ashton, P.J., Cowling, R.M., Higgins, J.V. et al. (2008). Progress and challenges in freshwater conservation planning. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 19(4), 474–485. <https://doi.org/10.1002/aqc.1010>

- Nguyen, V.M., Larocque, S.M., Stoot, L.J., Cairns, N.A., Blouin-Demers, G. & Cooke, S.J. (2013). Perspectives of fishers on turtle bycatch and conservation strategies in a small-scale inland commercial fyke net fishery. *Endangered Species Research*, 22(1), 11–22. <https://doi.org/10.3354/esr00530>
- Nguyen, V.M., Young, N. & Cooke, S.J. (2017). A roadmap for knowledge exchange and mobilization research in conservation and natural resource management. *Conservation Biology*, 31(4), 789–798. <https://doi.org/10.1111/cobi.12857>
- Olden, J.D., Kennard, M.J., Lawler, J.J. & Poff, N.L. (2011). Challenges and opportunities in implementing managed relocation for conservation of freshwater species. *Conservation Biology*, 25(1), 40–47. <https://doi.org/10.1111/j.1523-1739.2010.01557.x>
- Olugunorisa, T.E. (2009). Strategies for mitigation of flood risk in the Niger Delta, Nigeria. *Journal of Applied Sciences and Environmental Management*, 13(2), 17–22. <https://doi.org/10.4314/jasem.v13i2.55295>
- Opperman, J., Hartmann, J., Lambrides, M., Carvallo, J.P., Chapin E., Baruch-Mordo, S. et al. (2019). Connected and flowing: A renewable future for rivers, climate and people. WWF and the nature conservancy, Washington, DC. Available at: <https://www.worldwildlife.org/publications/connected-flowing-a-renewable-future-for-rivers-climate-and-people>
- Opperman, J.J., Kendy, E. & Barrios, E. (2019). Securing environmental flows through system reoperation and management: Lessons from case studies of implementation. *Frontiers in Environmental Science*, 7, 104. <https://doi.org/10.3389/fenvs.2019.00104>
- O'Riordan, T. & Cameron, J. (1994). *Interpreting the precautionary principle*. London: Earthscan Publications Ltd.
- Ormerod, S.J., Durance, I., Terrier, A. & Swanson, A.M. (2010). Priority wetland invertebrates as conservation surrogates. *Conservation Biology*, 24(2), 573–582. <https://doi.org/10.1111/j.1523-1739.2009.01352.x>
- Parsons, E.C.M., Favaro, B., Aguirre, A.A., Bauer, A.L., Blight, L.K., Cigliano, J.A. et al. (2014). Seventy-one important questions for the conservation of marine biodiversity. *Conservation Biology*, 28(5), 1206–1214. <https://doi.org/10.1111/cobi.12303>
- Peres, C.K., Lambrecht, R.W., Tavares, D.A. & Chiba de Castro, W.A. (2018). Alien express: The threat of aquarium e-commerce introducing invasive aquatic plants in Brazil. *Perspectives in Ecology and Conservation*, 16(4), 221–227. <https://doi.org/10.1016/j.pecon.2018.10.001>
- Phang, S.C., Cooperman, M., Lynch, A.J., Steel, E.A., Elliott, V., Murchie, K. J. et al. (2019). Fishing for conservation of freshwater tropical fishes in the Anthropocene. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 29(7), 1039–1051. <https://doi.org/10.1002/aqc.3080>
- Pittock, J., Dumaresq, D. & Orr, S. (2017). The Mekong River: Trading off hydropower, fish, and food. *Regional Environmental Change*, 17(8), 2443–2453. <https://doi.org/10.1007/s10113-017-1175-8>
- Poff, N.L. & Olden, J.D. (2017). Can dams be designed for sustainability? *Science*, 358(6368), 1252–1253. <https://doi.org/10.1126/science.aaq1422>
- Pullin, A.S. & Knight, T.M. (2003). Support for decision making in conservation practice: An evidence-based approach. *Journal for Nature Conservation*, 11(2), 83–90. <https://doi.org/10.1078/1617-1381-00040>
- Pyšek, P., Hulme, P.E., Simberloff, D., Bacher, S., Blackburn, T.M., Carlton, J.T. et al. (2020). Scientists' warning on invasive alien species. *Biological Reviews*, 95(6), 1511–1534. <https://doi.org/10.1111/brv.12627>
- Rahel, F.J. & Olden, J.D. (2008). Assessing the effects of climate change on aquatic invasive species. *Conservation Biology*, 22(3), 521–533. <https://doi.org/10.1111/j.1523-1739.2008.00950.x>
- Reid, A.J., Carlson, A.K., Creed, I.F., Eliason, E.J., Gell, P.A., Johnson, P.T.J. et al. (2019). Emerging threats and persistent conservation challenges for freshwater biodiversity. *Biological Reviews*, 94(3), 849–873. <https://doi.org/10.1111/brv.12480>
- Richardson, J.S., Naiman, R.J. & Bisson, P.A. (2012). How did fixed-width buffers become standard practice for protecting freshwaters and their riparian areas from forest harvest practices? *Freshwater Science*, 31(1), 232–238. <https://doi.org/10.1899/11-031.1>
- Rytwinski, T., Cooke, S.J., Taylor, J.J., Roche, D.G., Smith, P.A., Mitchell, G.W. et al. (2021). Acting in the face of evidentiary ambiguity, bias, and absence arising from systematic reviews in applied environmental science. *Science of the Total Environment*, 775, 145122. <https://doi.org/10.1016/j.scitotenv.2021.145122>
- Rytwinski, T., Taylor, J.J., Donaldson, L.A., Britton, J.R., Browne, D.R., Gresswell, R.E. et al. (2018). The effectiveness of non-native fish removal techniques in freshwater ecosystems: A systematic review. *Environmental Reviews*, 27(1), 71–94. <https://doi.org/10.1139/er-2018-0049>
- Saunders, D.L., Meeuwig, J.J. & Vincent, A.C.J. (2002). Freshwater protected areas: Strategies for conservation. *Conservation Biology*, 16(1), 30–41. <https://doi.org/10.1046/j.1523-1739.2002.99562.x>
- Schmidt-Kloiber, A., Bremerich, V., De Wever, A., Jähniq, S.C., Martens, K., Strackbein, J. et al. (2019). The Freshwater Information Platform: A global online network providing data, tools and resources for science and policy support. *Hydrobiologia*, 838(1), 1–11. <https://doi.org/10.1007/s10750-019-03985-5>
- Schreckenberg, K., Mace, G. & Poudyal, M. (Eds.) (2018). *Ecosystem services and poverty alleviation (OPEN ACCESS) – Trade-offs and governance*, 1st edition. London: Routledge. Available at: <https://www.taylorfrancis.com/books/e/9780429507090>
- Schuster, R., Germain, R.R., Bennett, J.R., Reo, N.J. & Arcese, P. (2019). Vertebrate biodiversity on indigenous-managed lands in Australia, Brazil, and Canada equals that in protected areas. *Environmental Science & Policy*, 101, 1–6. <https://doi.org/10.1016/j.envsci.2019.07.002>
- Shao, X., Fang, Y., Jawitz, J.W., Yan, J. & Cui, B. (2019). River network connectivity and fish diversity. *Science of the Total Environment*, 689, 21–30. <https://doi.org/10.1016/j.scitotenv.2019.06.340>
- Sleight, N. & Neeson, T.M. (2018). Opportunities for collaboration between infrastructure agencies and conservation groups: Road-stream crossings in Oklahoma. *Transportation Research Part D: Transport and Environment*, 63, 622–631. <https://doi.org/10.1016/j.trd.2018.07.002>
- Snyder, N.F.R., Derrickson, S.R., Beissinger, S.R., Wiley, J.W., Smith, T.B., Toone, W.D. et al. (1996). Limitations of captive breeding in endangered species recovery. *Conservation Biology*, 10(2), 338–348. <https://doi.org/10.1046/j.1523-1739.1996.10020338.x>
- Søndergaard, M., Jeppesen, E., Lauridsen, T.L., Skov, C., Nes, E.H.V., Roijackers, R. et al. (2007). Lake restoration: Successes, failures and long-term effects. *Journal of Applied Ecology*, 44(6), 1095–1105. <https://doi.org/10.1111/j.1365-2664.2007.01363.x>
- Sousa, E., Quintino, V., Palhas, J., Rodrigues, A.M. & Teixeira, J. (2016). Can environmental education actions change public attitudes? An example using the pond habitat and associated biodiversity. *PLoS ONE*, 11(5), e0154440. <https://doi.org/10.1371/journal.pone.0154440>
- Strayer, D.L. & Dudgeon, D. (2010). Freshwater biodiversity conservation: Recent progress and future challenges. *Journal of the North American Benthological Society*, 29(1), 344–358. <https://doi.org/10.1899/08-171.1>
- Sutherland, W.J., Fleishman, E., Mascia, M.B., Pretty, J. & Rudd, M.A. (2011). Methods for collaboratively identifying research priorities and emerging issues in science and policy. *Methods in Ecology and Evolution*, 2(3), 238–247. <https://doi.org/10.1111/j.2041-210X.2010.00083.x>

- Sutherland, W.J., Pullin, A.S., Dolman, P.M. & Knight, T.M. (2004). The need for evidence-based conservation. *Trends in Ecology & Evolution*, 19(6), 305–308. <https://doi.org/10.1016/j.tree.2004.03.018>
- Tanentzap, A.J., Lamb, A., Walker, S. & Farmer, A. (2015). Resolving conflicts between agriculture and the natural environment. *PLoS Biology*, 13(9), e1002242. <https://doi.org/10.1371/journal.pbio.1002242>
- Tennant, J.P., Waldner, F., Jacques, D.C., Masuzzo, P., Collister, L.B. & Hartgerink, C.H.J. (2016). The academic, economic and societal impacts of Open Access: An evidence-based review. *F1000Research*, 5, 632. <https://doi.org/10.12688/f1000research.8460.3>
- The Food and Land Use Coalition [FOLU]. (2019). Growing better: Ten critical transitions to transform food and land use. The Global Consultation Report of the Food and Land Use Coalition.
- Thomas, S.M., Griffiths, S.W. & Ormerod, S.J. (2016). Beyond cool: Adapting upland streams for climate change using riparian woodlands. *Global Change Biology*, 22(1), 310–324. <https://doi.org/10.1111/gcb.13103>
- Tickner, D., Opperman, J.J., Abell, R., Acreman, M., Arthington, A.H., Bunn, S.E. et al. (2020). Bending the curve of global freshwater biodiversity loss: An emergency recovery plan. *Bioscience*, 70(4), 330–342. <https://doi.org/10.1093/biosci/biaa002>
- Turak, E., Bush, A., Dela-Cruz, J. & Powell, M. (2020). Freshwater reptile persistence and conservation in cities: Insights from species occurrence records. *Water*, 12(3), 651. <https://doi.org/10.3390/w12030651>
- Turak, E., Harrison, I., Dudgeon, D., Abell, R., Bush, A., Darwall, W. et al. (2017). Essential biodiversity variables for measuring change in global freshwater biodiversity. *Biological Conservation*, 213(B), 272–279. <https://doi.org/10.1016/j.biocon.2016.09.005>
- United Nations. (2018). Sustainable development knowledge platform. Available at: <https://sustainabledevelopment.un.org/>
- van Rees, C.B., Waylen, K.A., Schmidt-Kloiber, A., Thackeray, S.J., Kalinkat, G., Martens, K. et al. (2020). Safeguarding freshwater life beyond 2020: Recommendations for the New Global Biodiversity Framework from the European experience. *Conservation Letters*, 14(1), e12771. <https://doi.org/10.20944/preprints202001.0212.v1>
- Verissimo, D., Pongiluppi, T., Santos, M.C.M., Develey, P.F., Fraser, I., Smith, R.J. et al. (2014). Using a systematic approach to select flagship species for bird conservation. *Conservation Biology*, 28(1), 269–277. <https://doi.org/10.1111/cobi.12142>
- Watson, B., Archer, E., Dziba, L., Fischer, M., Karki, M., Mulongoy, K.J. et al. (2018). Key findings from the four IPBES regional assessments of biodiversity and ecosystem services. In: *Convention on Biological Diversity*. Sharm El-Sheikh, Egypt: Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, p. 20.
- Weijters, M.J., Janse, J.H., Alkemade, R. & Verhoeven, J.T.A. (2009). Quantifying the effect of catchment land use and water nutrient concentrations on freshwater river and stream biodiversity. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 19(1), 104–112. <https://doi.org/10.1002/aqc.989>
- Whitehead, A.L., Kujala, H., Ives, C.D., Gordon, A., Lentini, P.E., Wintle, B.A. et al. (2014). Integrating biological and social values when prioritizing places for biodiversity conservation. *Conservation Biology*, 28(4), 992–1003. <https://doi.org/10.1111/cobi.12257>
- World Wildlife Fund [WWF]. (2020). In: R.E.A. Almond, M. Grooten, T. Petersen (Eds.) *Living planet report 2020 – Bending the curve of biodiversity loss*. Gland, Switzerland: WWF.
- Zhou, S., Kolding, J., Garcia, S.M., Plank, M.J., Bundy, A., Charles, A. et al. (2019). Balanced harvest: Concept, policies, evidence, and management implications. *Reviews in Fish Biology and Fisheries*, 29(3), 711–733. <https://doi.org/10.1007/s11160-019-09568-w>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Harper, M., Mejbil, H.S., Longert, D., Abell, R., Beard, T.D., Bennett, J.R. et al. (2021). Twenty-five essential research questions to inform the protection and restoration of freshwater biodiversity. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 1–22. <https://doi.org/10.1002/aqc.3634>