


# Exploring relationships between oxygen consumption and bilogger-derived estimates of heart rate in two warmwater piscivores

Claire L.J. Doherty<sup>1</sup> | Aaron T. Fisk<sup>1,2</sup> | Steven J. Cooke<sup>3</sup> | Trevor E. Pitcher<sup>1,4</sup> | Graham D. Raby<sup>5</sup> 

<sup>1</sup>Great Lakes Institute for Environmental Research, University of Windsor, Windsor, Ontario, Canada

<sup>2</sup>School of the Environment, University of Windsor, Windsor, Ontario, Canada

<sup>3</sup>Department of Biology and Institute of Environmental and Interdisciplinary Science, Carleton University, Ottawa, Ontario, Canada

<sup>4</sup>Department of Integrative Biology, University of Windsor, Windsor, Ontario, Canada

<sup>5</sup>Department of Biology, Trent University, Peterborough, Ontario, Canada

## Correspondence

Graham D. Raby, Department of Biology, Trent University, Peterborough, ON, K9L 0G2, Canada.

Email: grahamraby@trentu.ca

## Funding information

Canada Research Chairs; Natural Sciences and Engineering Research Council of Canada; Canada Foundation for Innovation; Ontario Research Fund; University of Windsor

## Abstract

Estimating metabolic rate in wild, free-swimming fish is inherently challenging. Here, we explored using surgically implanted heart rate loggers to estimate metabolic rate in two warmwater piscivores, bowfin *Amia calva* (Linnaeus 1766) and largemouth bass *Micropterus salmoides* (Lacepède 1802). Fish were surgically implanted with heart rate loggers, allowed to recover for 24 h, exposed to a netting and air exposure challenge, and then placed into respirometry chambers so that oxygen consumption rate ( $\dot{M}_{O_2}$ ) could be measured in parallel to heart rate ( $f_H$ ) for a minimum of 20 h (ca. 20 estimates of  $\dot{M}_{O_2}$ ). Heart rate across the duration of the experiment (at 19°C) was significantly higher in largemouth bass (mean  $\pm$  s.d.,  $45 \pm 14$  beats  $\text{min}^{-1}$ , range 18–86) than in bowfin ( $27 \pm 9$  bpm, range 16–98). Standard metabolic rate was also higher in largemouth bass ( $1.06 \pm 0.19$  mg  $\text{O}_2$   $\text{kg}^{-1}$   $\text{min}^{-1}$ , range 0.46–1.36) than in bowfin ( $0.89 \pm 0.17$  mg  $\text{O}_2$   $\text{kg}^{-1}$   $\text{min}^{-1}$ , range 0.61–1.28). There were weak relationships between  $f_H$  and  $\dot{M}_{O_2}$ , with heart rate predicting 28% of the variation in oxygen consumption in bowfin and 23% in largemouth bass. The shape of the relationship differed somewhat between the two species, which is perhaps unsurprising given their profound differences in physiology and life history, illustrating the need to carry out species-specific validations. Both species showed some potential for a role of  $f_H$  in efforts to estimate field metabolic rates, although further validation experiments with a wider range of conditions (e.g., digestive states, swimming activity) would likely help improve the strength of the  $\dot{M}_{O_2}$ – $f_H$  relationship for use in field applications.

## KEYWORDS

bioenergetics, bilogger, cardiac output, ECG, electrocardiogram

## 1 | INTRODUCTION

A large amount of research in biology is focused on examining how variation in metabolic rate shapes and responds to ecological and evolutionary processes (Réale *et al.*, 2010; Ricklefs & Wikelski, 2002; Tomlinson *et al.*, 2014). In fishes, bioenergetics modelling provides a holistic, first-principles framework for quantifying how fish respond

to environmental dynamics (Deslauriers *et al.*, 2017; Jobling, 1994). Metabolic rate, a crucial parameter in the field of ecological energetics and in bioenergetics modelling (Winberg, 1960), is typically estimated *via* the rate of oxygen consumption ( $\dot{M}_{O_2}$ ) by confining fish in respirometry chambers in the laboratory (Nelson, 2016). However, our ability to understand metabolism in wild fish and to develop ecologically accurate bioenergetic models could be

improved by being able to quantify natural variation in the metabolic rate of fish *in situ* (Cooke *et al.*, 2016).

There has long been an interest among fish biologists in using electronic sensor tags to remotely estimate metabolic rate in free-swimming fish. Heart rate ( $f_H$ ), which can be estimated *via* electrocardiogram (ECG) data, has the potential to act as a proxy for metabolic rate because the heart is directly involved in the uptake and transport of oxygen (Butler *et al.*, 2004). Heart rate tags have been the subject of experimentation with fish for decades, having been usually custom-made (e.g., Armstrong, 1986; Kneis & Siegmund, 1976; Priede, 1974; Priede & Tytler, 1977) and difficult to work with, often being restricted to a laboratory setting or including an external battery pack (e.g., Cooke *et al.*, 2004). In recent years, researchers have begun using commercially available heart rate loggers developed by Star Oddi (<https://www.star-oddi.com/>; e.g., Prystay *et al.*, 2017, Brijis *et al.*, 2019), which are much smaller than older models and measure electrocardiograms (ECGs) through electrodes on the side of the logger and calculate  $f_H$  using pre-programmed algorithms. These loggers can be surgically implanted in wild fish and make intermittent recordings for weeks or months, overcoming limitations associated with laboratory experiments (see Cooke *et al.*, 2016), although recapture of the study animals is required for logger download; a logistical challenge for many wild fishes. In addition,  $f_H$  needs to be validated as a proxy for metabolic rate before use and validations likely need to be done on a genus- or species-specific basis (Butler *et al.*, 2004). Even in the ‘best-case scenario’,  $f_H$  is only likely to provide a relative approximation of  $\dot{M}_{O_2}$  because  $\dot{M}_{O_2}$  is modulated by physiological functions other than  $f_H$ , such as the stroke volume of the heart and the rate of oxygen diffusion at the gills (e.g., changes in ventilation rate; Farrell, 1991). However, in free-swimming fish,  $f_H$  currently represents one of our only options for attempting to indirectly estimate variation in  $\dot{M}_{O_2}$  because both  $\dot{M}_{O_2}$  and  $f_H$  typically respond similarly to swimming activity (Webber *et al.*, 1998), recovery from exercise (Prystay *et al.*, 2017), feeding and digestion (Armstrong, 1986), or variation associated with temperature (Priede, 1983).

This study assessed the use of  $f_H$  as a proxy for metabolic rate in two freshwater fishes, bowfin *Amia calva* and largemouth bass *Micropterus salmoides*. These species were selected due to their broad distribution in North American freshwaters and the fact that both grow to a size appropriate for use of electronic tags of the size we used here (13 mm diameter  $\times$  39.5 mm long). Cardiac physiology has been examined in these species a handful of times (bowfin, Porteus *et al.*, 2014a,b; largemouth bass, Cooke *et al.*, 2002, 2004). Given the heart’s direct role in modulating oxygen transport and uptake (Satchell, 1991), we hypothesized that there would be a positive relationship between  $f_H$  and  $\dot{M}_{O_2}$  in both bowfin and largemouth bass. Because of the profound differences between bowfin and largemouth bass, we predicted that the relationship between heart rate and metabolic rate would differ between these species. The largemouth bass is a warmwater piscivore with common teleost features and anatomy. Bowfin is one of few extant members of the infraclass Holostei; they differ from largemouth bass in several obvious ways,

the most notable of which is that they use a gas bladder to gulp and breathe air, an adaptation that makes them well suited to hypoxic waters (Porteus *et al.* 2014a). To facilitate the validation of the relationship between  $\dot{M}_{O_2}$  and  $f_H$ , we exposed every fish to a confinement + air exposure stressor immediately before entry into the respirometers. That step was designed to cause some level of exhaustion and physiological recovery to elicit a broader range of  $f_H$  and  $\dot{M}_{O_2}$  than would be expected in fish simply resting in respirometers. Given that relationships between  $f_H$  and  $\dot{M}_{O_2}$  are not always entirely clear (Thorarensen *et al.*, 1996), laboratory experiments are an important first step in evaluating the utility of  $f_H$  loggers for studying the ecological energetics of wild fish (Muller *et al.*, 2020; Zrini & Gamperl, 2021).

## 2 | MATERIALS AND METHODS

### 2.1 | Fish collection

Twenty-one (21) bowfin [mean mass = 1561 g, range = 467–2404 g; mean total length (TL) = 568 mm, range 395–666 mm] and 30 largemouth bass (mean 661 g, range 358–1288 g; mean TL 351 mm, range 292–444 mm) were collected by boat electrofishing from the Detroit River in the waters around Fighting Island in LaSalle, Ontario (42°13′14.1″N, 83°06′53.2″W and 42°11′26.6″N, 83°06′55.4″W) between 15 May and 11 June 2019. Once caught, fish were held in an onboard aerated tank transported to the nearby Freshwater Restoration Ecology Centre in LaSalle ON (42°14′10.4″N, 83°06′17.9″W) (total time of 15–180 min onboard). On arrival, fish were given either an anchor tag or a passive integrative transponder (PIT) tag for identification and transferred into indoor 850 l holding tanks set to either 14 or 19°C depending on tank availability as there were multiple experiments running parallel within the same facility that required the tank systems to be set to these two temperatures. All respirometry and  $f_H$  data were collected at 19°C, which was closer to the median temperature of the Detroit River during the period when the experiment was taking place. Therefore, some fish were acutely acclimated from 14 to 19°C after implantation of  $f_H$  loggers, meaning these fish were not ‘fully acclimated’ to the experimental temperature. However, both species live in shallow thermally dynamic waters and in the wild can freely move through thermally heterogeneous environments, so we considered these thermal changes to be within the scope of what is ecologically realistic. Tanks were on a recirculation system with UV sterilization, aeration and filtration of dechlorinated municipal water, with temperature regulated by thermostat-controlled chillers. Fish were held in a 7-day quarantine, as per institutional animal care requirements (for biosecurity purposes), prior to use in experiments. No fish died, and all animals appeared to be in good condition following the quarantine period. All procedures for this study were approved by the University of Windsor Animal Care Committee following guidance set by the Canadian Council on Animal Care (Animal Utilization Project Proposal #19–08).

## 2.2 | Heart rate logger implantation

Following the quarantine period, fish had Star-Oddi DST-Milli-HRT loggers (11.8 g in air, 13 mm diameter × 39.5 mm length; Star-Oddi, Gardabaer, Iceland; <https://www.star-oddi.com/>) surgically implanted into the body cavity. Loggers were programmed using the software Mercury (Star-Oddi) to turn on and record the ECG for 7.5 s at 80 Hz every 10 min, from which  $f_H$  was calculated (beats  $\text{min}^{-1}$ ) and given a quality index value using a pre-programmed algorithm. The loggers were programmed to save a raw trace of the ECG data every 6 h, which could be used to manually confirm the accuracy of the onboard  $f_H$  algorithm. In addition, the loggers recorded visceral temperature at each measurement period (data not presented here; all fish were held at a common temperature after logger implantation). After programming, loggers were fitted with a size 2 suture and sterilized along with all tools by immersion in betadine solution prior to surgery. A 20 l anaesthetic bath was prepared with a concentration of 125  $\text{mg l}^{-1}$  tricaine methanesulfonate (MS-222) buffered by 250  $\text{g l}^{-1}$  sodium bicarbonate. Individual fish were placed in the bath and left until they lost equilibrium (typically 4–5 min) and then measured (total length to nearest mm, weight to the nearest 0.5 g). Fish were continuously supplied with aerated water with a maintenance dose of 80  $\text{mg l}^{-1}$  MS-222 (160  $\text{mg l}^{-1}$  sodium bicarbonate) during surgery. Based on the differing position of the pelvic and pectoral fins (and associated cartilage) relative to the pericardium, an incision of 2 cm was made either between the pelvic fins just off the ventral midline (bowfin) or between the pelvic and pectoral fins on the left side (largemouth bass) of the fish. Loggers were inserted and oriented within the body cavity such that both electrodes were facing the pericardial cavity. Loggers were secured in place with an anchor suture and the incision closed with two sutures; all sutures were done using a 2–3–2 knot. Fish were placed in an 80 l aerated recovery tank until they had fully regained their righting reflex (*ca.* 5–10 min) and then transferred to a 19°C 850 l holding tank for 24 h of recovery from surgery. Depending on the timing of the preceding trials, the respirometers were, in a few cases, available at the time fish came out of surgery (previous fish had been removed from respirometers). In those cases, fish were transferred into respirometers immediately after surgery to begin their recovery (see below) to increase the amount of  $\dot{M}_{\text{O}_2}$  data we could compare against  $f_H$ .

## 2.3 | Respirometry

After the 24 h recovery period, fish were exposed to a handling stressor to elicit an elevated metabolic rate (or active metabolic rate, AMR) to facilitate a broader range of both  $f_H$  and  $\dot{M}_{\text{O}_2}$  values for the purpose of validation. The handling stressor involved fish being netted using a knotless nylon dipnet and then held at the water–air interface for 2 min (at least one set of gills always submerged) and then completely air exposed for 1 min. At the end of the air exposure, fish were sealed into respirometry chambers that were either 16 l (internal dimensions 42 cm long × 20.2 cm wide × 19 cm deep; 26 largemouth

bass, two bowfin) or 74.2 l (internal dimensions 80.5 cm × 30.5 cm × 30.5 cm; five largemouth bass, 20 bowfin), with the size of the chamber used for a particular fish being driven by the size of the animal. Typically, four or five respirometers were running in parallel, and fish were all exposed to the stressor and transferred to respirometers within a 5 min period. Once all fish were in respirometers, the respirometer flush pumps were manually switched off for an initial sealed measurement period to estimate AMR. Flush pumps were then switched back on when dissolved oxygen (DO) reached  $\sim 7 \text{ mg/l}$  ( $\sim 80\%$  air saturation) and thereafter controlled with timers set for 30 min flush and 15 min measurement (sealed) periods (74.2 l chambers), or with a 20 min flush and 10 min seal cycle (16 l). Respirometry trials then continued for a minimum of 20 h, and up to 45 h in some cases because of logistical constraints (availability of the research team). Mixing was maintained in each chamber through a recirculation line (Clark *et al.*, 2013) which contained an optical oxygen probe connected to a Firesting system (PyroScience, Aachen, Germany) for DO measurements (at 0.5 Hz; *e.g.*, as in Raby *et al.*, 2020; Sundin *et al.*, 2019). Each chamber was also outfitted with a standpipe for overflow of which two had temperature probes placed in them (Clark *et al.*, 2013). Background (microbial) measurements were made in each chamber before and after each trial (Sundin *et al.*, 2019). In data analyses, background  $\dot{M}_{\text{O}_2}$  was always subtracted from total respiration to calculate fish  $\dot{M}_{\text{O}_2}$  and always remained below 11.5% of fish  $\dot{M}_{\text{O}_2}$  (median = 0.89%).

## 2.4 | Data analysis and statistics

After respirometry trials, fish were euthanized using a 15  $\text{ml l}^{-1}$  clove oil bath, loggers were retrieved and their data downloaded. Loggers were re-programmed, sterilized and used in subsequent trials. The mean slope from each respirometry measurement period was calculated using a LabChart Reader (ADI Instruments, Sydney, Australia), leaving out the first and last minute of the slope to avoid any lag at the beginning of the cycle and any influence from the pumps turning back on at the end of the sealed cycle. This data was used to determine  $\dot{M}_{\text{O}_2}$  using the equation

$$\dot{M}_{\text{O}_2} = V_{\text{RE}} \times M_{\text{W}}^{-1} \times \delta\text{CO}_2\delta t^{-1}$$

in which  $V_{\text{RE}}$  is the effective chamber volume (in l),  $M_{\text{W}}$  is the mass of the fish (in kg) and  $\delta\text{CO}_2\delta t^{-1}$  is the linear slope from oxygen decrease during sealed cycles ( $\text{mg O}_2 \text{ min}^{-1}$ ) (Clark *et al.*, 2013). Standard metabolic rate (SMR) was calculated as the 15th percentile of all data points for each individual (Chabot *et al.*, 2016), routine metabolic rate (RMR) was the 50th percentile, AMR was the 100th percentile. Each of these variables used  $\dot{M}_{\text{O}_2}$  data and is presented in  $\text{mg O}_2 \text{ kg}^{-1} \text{ min}^{-1}$ .

Heart rate data were automatically assigned a quality index ranking (0, best to 3, poorest) based on how ‘clean’ the ECG is using a proprietary Star-Oddi algorithm pre-programmed into loggers, meant to indicate the reliability of the  $f_H$  estimate as the ECG data can be noisy


depending on the position of the logger within the body cavity, the strength of the electric signal from the heart and surrounding electronic noise (Brijs *et al.*, 2018; Muller *et al.*, 2020; Zrini & Gamperl, 2021). Heart rate data were filtered to include only points of quality index 0–1 (55% for bowfin, 56% for largemouth bass). Data were then inspected and filtered further. Specifically, we used a smoothing function whereby any data point that showed a greater difference than 10 bpm to the immediately preceding *and* proceeding points was removed. We also removed values  $>100$  for bpm (*ca.* 0.03% of data points) because we considered these to be physiologically unrealistic at these temperatures. However, for model fitting only points within 20–80 bpm for largemouth bass and  $\leq 50$  bpm for bowfin were kept due to lack of sufficient data beyond these ranges to produce robust model estimates (these ranges were manually chosen based on visual inspection of the distribution of all data points). Of the 21 bowfin and 30 largemouth bass that received loggers, eight bowfin and 12 largemouth bass yielded sufficient high-quality  $f_H$  data for statistical analysis for a total of 17.4% and 20.8% of data points respectively after filtering.

Filtered  $f_H$  data were then matched to corresponding  $\dot{M}_{O_2}$  data points for the same animal that had a timestamp within 8 min. The timestamp for  $f_H$  data were exact times (the measurement periods were only 7.5 s); for the longer  $\dot{M}_{O_2}$  measurement periods (10 or 15 min long, depending on respirometer size), the timestamp represented the time at the middle of the measurement period. If any  $f_H$  data point fell within 8 min of an  $\dot{M}_{O_2}$  timestamp, it was kept and matched with that  $\dot{M}_{O_2}$  data point. This meant that measurements of  $\dot{M}_{O_2}$  were typically assigned to one or two measurements of  $f_H$ . Note: We assessed the resulting model fits ( $R^2$ ) based on 6, 7, 8, 9 and 10 min windows for matching  $\dot{M}_{O_2}$  and  $f_H$  data and found subtle differences, with 8 min providing the best model fit; smaller sampling windows meant a closer temporal match between data points, but a smaller sample size for modelling (less power). Linear mixed effects models (LME) and general additive mixed models (GAMM) were applied to the relationships between  $f_H$  and  $\dot{M}_{O_2}$  for both species to determine model of best fit for each. We modelled the effect of  $f_H$  (fixed effect) on  $\dot{M}_{O_2}$  (response variable) with animal ID as a random effect ( $n = 8$  animals for the bowfin model,  $n = 12$  animals for the largemouth bass model). We assessed whether the data met the assumptions of parametric statistics by plotting model residuals against fitted values and all predictor variables by use of q-q plots, and by use of autocorrelation function plots. Models were fit using a temporal autocorrelation structure (using the 'corARMA' function in the package 'mgcv') to control for the lack of independence of data points from the same fish measured at proximate times. We compared model fits using Akaike Information Criterion (AIC) and log-likelihood tests. Model terms are reported as significant at  $P < 0.05$ . Data files and analysis code for this paper are publicly archived with figshare (<https://doi.org/10.6084/m9.figshare.15090936.v2>).


### 3 | RESULTS

There were clear differences in  $f_H$  and  $\dot{M}_{O_2}$  patterns between bowfin and largemouth bass. While bowfin demonstrated a much narrower


range of  $f_H$  when looking at the 5th and 95th percentiles (bowfin 19–41 beats  $\text{min}^{-1}$ , largemouth bass 27–69 beats  $\text{min}^{-1}$ ), the full range of  $f_H$  in bowfin (16–98 beats  $\text{min}^{-1}$ ) was comparable to largemouth bass (18–86 beats  $\text{min}^{-1}$ ) (Figure 1a). However, largemouth bass exhibited greater increases in  $f_H$  in response to the acute stressor (median increase of 32 beats  $\text{min}^{-1}$ ) than did bowfin (median of 15 beats  $\text{min}^{-1}$ ) (Figure 2). Mean  $f_H$  across the entirety of the trial was higher in largemouth bass ( $45 \pm 14$  beats  $\text{min}^{-1}$ , mean  $\pm$  s.d.) than in bowfin ( $27 \pm 9$  beats  $\text{min}^{-1}$ , Welch's *t*-test, pooling data for both species  $t_{3728.5} = -43.4$ ,  $P < 0.001$ ). A total of 986 measurements of  $\dot{M}_{O_2}$  were collected among 21 bowfin and 1872 among 30 largemouth bass that could be used to estimate SMR, RMR and AMR. SMR and AMR were higher in largemouth bass than in bowfin while RMR was



**FIGURE 1** Physiological states exhibited by bowfin and largemouth bass held under controlled laboratory conditions at *ca.* 19°C. (a) shows heart rate ( $f_H$ ) with the following sample sizes:  $n = 1626$  data points from eight bowfin, 2092 data points from 12 largemouth bass. For the oxygen consumption parameters (b) standard metabolic rate (SMR), (c) routine metabolic rate (RMR), and active metabolic rate (AMR), there were 30 largemouth bass and 21 bowfin with one estimate per fish. For boxplots, the thick middle line is the median, the boxes extend to the 25th and 75th percentiles of the data, and the whiskers extend to the most extreme data points or  $1.5\times$  the interquartile range, whichever was less



**FIGURE 2** Examples of (a) largemouth bass and (b) bowfin heart rate ( $f_H$ ) data from two fish, highlighting the between-species differences in cardiac response to stressors applied in a controlled environment (time in hh:mm on x-axis). The blue vertical lines indicate the time when surgery took place; the yellow vertical lines indicate the handling and air exposure stressor. The matching oxygen consumption (= metabolic rate) data for the same individuals is shown by blue triangles (right axis)



**FIGURE 3** Scatterplot comparison of heart rate ( $f_H$ ) and oxygen consumption rate ( $\dot{M}_{O_2}$ ) in a controlled environment overlaid by a general additive mixed model (largemouth bass, right panel) and a linear mixed effects model (bowfin, left panel), yellow data points (translucent) represent quality index zero (best) and blue quality index one (good) that passed our filtering steps and were used for statistical modelling ( $n = 599$  data points from eight bowfin, 1068 data points from 12 largemouth bass)



lower ( $t$ -tests,  $F_{46,4} = -3.26$  (SMR),  $F_{48,9} = -1.42$  (AMR),  $F_{28,1} = 3.01$  (RMR), all  $P < 0.001$ ).

The strength of the relationship between  $f_H$  and  $\dot{M}_{O_2}$  was weak in both species, with differences in the shape of the relationship. A linear mixed-effects model was the best fit for the bowfin data [marginal  $R^2$  (fixed effects only) = 0.28, conditional  $R^2$  (fixed + random effects) = 0.49], whereby  $\dot{M}_{O_2}$  increased by 0.058 mg O<sub>2</sub> kg<sup>-1</sup> min<sup>-1</sup> for each 1 bpm increase in  $f_H$  ( $P < 0.001$ ). For largemouth bass, a GAMM (nonlinear model) was the best fit for the data (adjusted  $R^2 = 0.23$ ,  $P < 0.001$  for the main effect of  $f_H$ ). A nonlinear GAMM smoothing function fit best to the largemouth bass data because there appeared to be a linear relationship up to a  $f_H$  of ca. 66, at which point there was an inflection whereafter  $\dot{M}_{O_2}$  increased at a higher rate with further increases in  $f_H$ . However, there were few data above that 66 bpm inflection point, leading to wider confidence intervals (Figure 3).

## 4 | DISCUSSION

In this experimental study using wild-caught fishes, we found weak relationships between heart rate ( $f_H$ ) and rate of oxygen consumption ( $\dot{M}_{O_2}$ ) for bowfin and largemouth bass using heart rate loggers and respirometry. These experiments were carried out under a constrained range of conditions, whereby all fish were in the laboratory, fasted, recovering from a moderate handling stressor and confined in static respirometry chambers at a constant temperature. We had expected to see some variation in  $f_H$  and  $\dot{M}_{O_2}$  from spontaneous activity within the respirometry chambers, though both species mostly appeared to remain relatively 'calm' throughout. Perhaps partly because of that lack of activity-driven variation,  $f_H$  predicted only 28% and 23% of the variation in  $\dot{M}_{O_2}$  for bowfin and largemouth bass, respectively. One possibility is that our measurements of both parameters were too infrequent to detect brief bouts of spontaneous activity that were not accompanied by excess post-exercise oxygen consumption (the latter would mean a sustained increase in  $f_H$  and  $\dot{M}_{O_2}$  even after activity ceases). Regardless, heart rate has long been thought of as an imperfect proxy for  $\dot{M}_{O_2}$  in fish – as it was in the present study – for a variety of reasons, including the fact that  $f_H$  is one of several ways fish modulate their oxygen uptake and transport (stroke volume plays an important role in cardiac output; Thorarensen *et al.*, 1996; Priede & Tytler, 1977). However, there is growing evidence (relative to early syntheses by Farrell, 1991) that many fish are frequency modulators (including largemouth bass; Cooke *et al.*, 2004) and thus heart rate may indeed be a useful proxy. This proof-of-concept experiment suggests that while heart rate loggers may have some role in the 'toolbox' to help quantify metabolic rate in wild fishes (Cooke *et al.*, 2016), validation is needed and their utility may be limited in some species. While  $f_H$  on its own appears to be a poor choice for precisely estimating energy expenditure in these species, it could be useful for probabilistically assigning different physiological states (e.g., stressed vs. unstressed, resting vs. active, digesting vs. fasted). In any case, our findings underline the need for species-

specific validation experiments before  $f_H$  can be of use for studying the ecophysiology of fish in the wild.

Heart rate and its relationship with metabolic rate in untethered animals (i.e., using loggers or transmitters) has been studied in several fish species that have produced comparable results to the species used here. Armstrong (1986) found a positive linear relationship between these variables up to 55 beats min<sup>-1</sup> in northern pike (*Esox lucius*), after which the linear relationship dissolved due to larger increases in metabolic rate relative to  $f_H$ , similar to what we found in largemouth bass. Atlantic salmon (*Salmo salar*) also showed a strong linear relationship between  $f_H$  and metabolic rate on a logarithmic scale, one that extends past the point at which both pike (Lucas, 1994) and largemouth bass appear to dissolve. Other species, however, show weaker relationships between  $f_H$  and metabolic rate, such as Atlantic cod (*Gadus morhua*), which demonstrated a large range of  $\dot{M}_{O_2}$  values per  $f_H$  value (Priede & Tytler, 1977). Strong intra-specific variation may also explain weak relationships between  $f_H$  and  $\dot{M}_{O_2}$  in fish (Farrell, 1991; Lucas, 1994; Scharold & Gruber, 1991). Indeed, in our study the random effect term significantly improved model fits for both species (different intercepts for each individual animal), confirming that there are individual differences in the precise relationship between  $\dot{M}_{O_2}$  and  $f_H$ . In addition, trials longer than the 24–48 h timespan used here could facilitate recovery from surgery and better  $f_H$ – $\dot{M}_{O_2}$  relationships; Webber *et al.* (1998) found that the relationship between these variables improved over seven days in Atlantic cod.

Aspects of the 'lifestyle' of the two species studied here may be relevant to consider when interpreting the weak relationships between  $f_H$  and  $\dot{M}_{O_2}$  we observed. Bowfin and largemouth bass are both ambush predators that live primarily in shallow, warm habitats with submerged vegetation. Our test of the  $f_H$ – $\dot{M}_{O_2}$  relationship was quite conservative in the sense that we exposed fish to a narrow set of conditions: one temperature, confinement in a respirometer in the laboratory, with all fish being post-absorptive. Considering this context casts a more positive light on the somewhat weak relationship we did observe in the two species. Given that both species are sit-and-wait predators, there would likely be a benefit to including post-prandial metabolic responses in future  $f_H$  validation experiments. For example, in the invasive lionfish (*Pterois* spp.), another sit-and-wait predator, the rise in metabolic rate during digestion eclipses that from exhaustive exercise (Stell *et al.*, 2019). Indeed, the relative advantage of  $f_H$  over accelerometry as a means of remotely quantifying energy use is that it can detect the postprandial metabolic response (i.e., specific dynamic action; Secor, 2009). In contrast, for more 'active' species like salmonids, it could be that  $f_H$  has stronger relationships with other variables (Clark *et al.*, 2010) than the sit-and-wait predators like those used here.

Despite the loggers used here being commercially available and user-friendly, there are technical challenges and limitations in their use, not unlike previously used custom-made heart rate loggers (e.g., Campbell *et al.*, 2005; Kojima *et al.*, 2003). For example, only 20.8% of the data points from 30 largemouth bass and 17.4% of data points from 21 bowfin were of reliable quality to be used in analysis.

That shortcoming highlights the need for careful pilot experiments to determine optimal logger settings and placement in the body cavity (i.e., surgical methods, steps we were unable to take with the present study). For example, in this study loggers were placed perpendicular to the incision with the electrodes facing the pericardial cavity; another option to explore would be to place them parallel to the incision with the electrodes facing the abdominal muscles (Brijs *et al.*, 2018). Placement of the logger relative to the heart also likely plays a role in its efficiency, the further away the weaker the ECG signal, and to optimize results the loggers should be kept within 20 mm of the heart (Brijs *et al.*, 2018). Exact logger placement depends on the anatomy of the species (Muller *et al.*, 2020). For bowfin and largemouth bass, the position of the pelvic and pectoral fins and surrounding cartilage dictated where the incision could be made safely (without damaging important structures like cartilage). Loggers in our study were approximately 5–20 mm from the pericardium, with some variation related to body size, although it was difficult to verify the exact positioning of the tag once it was inside the animal (in part because it could have moved slightly, despite being sutured to the body wall).

Heart rate loggers offer the potential of being able to estimate metabolic rate in free-swimming fish in the wild. In this study, we piloted the use of heart rate loggers as a proxy for metabolic rate in laboratory-held, wild-caught largemouth bass and bowfin. While the relationships between  $f_H$  and  $\dot{M}_{O_2}$  were not strong, our experiment involved a narrow range of conditions. Therefore, we suggest that the relationships we found represent sufficient evidence to consider further experimentation that includes a broader range of conditions in order to elicit more within-individual variability, including dynamic changes in temperature, oxygen level, activity levels (e.g., using a swim tunnel respirometer) and digestive states. Nevertheless, it may simply be that in these species  $f_H$  on its own is not particularly useful for estimating metabolic rate. Accelerometry has also been shown to provide a good estimate of metabolic rate when used in conjunction with  $f_H$  (Clark *et al.*, 2010); this combination is something to consider for future efforts to remotely measure metabolic rate in free-swimming fish (a logger that measures acceleration,  $f_H$  and temperature is available from the same manufacturer; /www.star-oddi.com). The differences between the two species illustrates the need for species-specific validation of this relationship and the need for pilot studies to experiment with logger placement in relation to the heart.

## ACKNOWLEDGEMENTS

The authors thank M. Charron for his help with fish collection and laboratory experiments; J. Landry, S. Laroque, J. McAndrews, A. Weinz, M. Guzzo and N. Smith for help with fish collection; K. Johnson for help with laboratory experiments and logistics; and T. Fendler for help with organizing and carrying out sample processing. C.L.J.D. was supported in part by an Ontario Graduate Scholarship and funds from the University of Windsor. G.D.R. was supported by a Natural Sciences and Engineering Research Council of Canada (NSERC) Postdoctoral Fellowship. This work was supported by an NSERC Discovery Grant and the Canada Research Chairs Program (A.T.F.), and by

Canadian Foundation Innovation funds via the Real Time Aquatic Ecosystem Observation Network (RAEON; A.T.F., S.J.C., and T.E.P.).

## AUTHOR CONTRIBUTIONS

G.D.R. and A.T.F. conceived and designed the experiment. T.E.P. and S.J.C. provided critical equipment and logistical support. C.L.J.D. and G.D.R. collected and analysed the data. C.L.J.D. wrote the manuscript with input from all co-authors.

## ORCID

Graham D. Raby  <https://orcid.org/0000-0002-0549-4017>

## REFERENCES

- Armstrong, J. D. (1986). Heart rate as an indicator of activity, metabolic, food intake, and digestion in pike, *Esox Lucius*. *Journal of Fish Biology*, 29, 207–221.
- Brijs, J., Sandblom, E., Axelsson, M., Sundell, K., Sundh, H., Huyben, D., ... Gräns, A. (2018). The final countdown: continuous physiological welfare evaluation of farmed fish during common aquaculture practices before and during harvest. *Aquaculture*, 495, 903–911.
- Brijs, J., Sandblom, E., Rosengren, M., Sundell, K., Berg, C., Axelsson, M., & Grans, A. (2019). Prospects and pitfalls of using heart rate bio-loggers to assess the welfare of rainbow trout (*Oncorhynchus mykiss*) in aquaculture. *Aquaculture*, 509, 188–197.
- Butler, P. J., Green, J. A., Boyd, I. L., & Speakman, J. R. (2004). Measuring metabolic rate in the field: the pros and cons of the doubly labelled water and heart rate methods. *Functional Ecology*, 18, 168–183.
- Campbell, H. A., Bishop, C. M., Davies, D. A., & Egginton, S. (2005). Recording long-term heart rate in *Paranotothenia angustata* using an electronic datalogger. *Journal of Fish Biology*, 67, 1150–1156.
- Chabot, D., Steffensen, J. F., & Farrell, A. P. (2016). The determination of standard metabolic rate in fishes. *Journal of Fish Biology*, 88, 81–121.
- Clark, T. D., Sandblom, E., Hinch, S. G., Patterson, D. A., Frappell, P. B., & Farrell, A. P. (2010). Simultaneous biologging of heart rate and acceleration and their relationship with energy expenditure in free-swimming sockeye salmon (*Oncorhynchus nerka*). *Journal of Comparative Physiology B*, 180, 673–684.
- Clark, T. D., Sandblom, E., & Jutfelt, F. (2013). Aerobic scope measurements of fishes in an era of climate change: respirometry, relevance and recommendations. *Journal of Experimental Biology*, 216, 2771–2782.
- Cooke, S. J., Brownscombe, J. W., Raby, G. D., Broell, F., Hinch, S. G., Clark, T. D., & Semmens, J. M. (2016). Remote bioenergetics measurements in wild fish: opportunities and challenges. *Comparative Biochemistry and Physiology Part A*, 202, 23–37.
- Cooke, S. J., Brunt, C. M., Ostrand, K. G., Philipp, D. P., & Wahl, D. H. (2004). Angling-induced cardiac disturbances of free-swimming largemouth bass (*Micropterus salmoides*) monitored with heart rate telemetry. *Journal of Applied Ichthyology*, 20, 28–36.
- Cooke, S. J., Brunt, C. M., Schreer, J. F., & Philipp, D. P. (2002). Attachment, validation, and preliminary deployment of ultrasonic heart rate transmitters on largemouth bass, *Micropterus salmoides*. *Aquatic Living Resources*, 15, 155–162.
- Deslauriers, D., Chipps, S. R., Breck, J. E., Rice, J. A., & Madenjian, C. P. (2017). Fish bioenergetics 4.0: an R-based modeling application. *Fisheries*, 42, 586–596.
- Farrell, A. P. (1991). From hagfish to tuna: a perspective on cardiac function in fish. *Physiological Zoology*, 64, 1137–1164.
- Jobling, M. (1994). *Fish bioenergetics* (p. 310). Dordrecht: Springer.

- Kneis, P., & Siegmund, R. (1976). Heart rate and locomotor activity in fish – correlation and circadian and circannual differences in *Cyprinus carpio* L. *Experientia*, 32, 474–476.
- Kojima, T., Kawabe, R., Shirasu, K., & Naito, Y. (2003). Preliminary study on heartbeats and swimming behaviour of free-ranging fish, red sea bream *Pagrus major*, measured with newly developed micro data-logger. *Polar Bioscience*, 16, 104–111.
- Lucas, M. C. (1994). Heart rate as an indicator of metabolic rate and activity in adult Atlantic salmon, *Salmo salar*. *Journal of Fish Biology*, 44, 889–903.
- Muller, C., Childs, A.-R., Duncan, M. I., Skeeles, M. R., James, N. C., van der Walt, K. A., ... Potts, W. M. (2020). Implantation, orientation and validation of a commercially produced heart-rate logger for use in a perciform teleost fish. *Conservation Physiology*, 8, coaa035.
- Nelson, J. A. (2016). Oxygen consumption rate v. rate of energy utilization of fishes: a comparison and brief history of the two measurements. *Journal of Fish Biology*, 88, 10–25.
- Porteus, C. S., Wright, P. A., & Milsom, W. K. (2014a). The effect of sustained hypoxia on the cardio-respiratory response of bowfin *Amia calva*: implications for changes in the oxygen transport system. *Journal of Fish Biology*, 84, 827–843.
- Porteus, C. S., Wright, P. A., & Milsom, W. K. (2014b). Time domains of the hypoxic-cardio response in bowfin (*Amia calva*). *Respiratory Physiology & Neurobiology*, 192, 118–127.
- Priede, I. G. (1974). The effect of swimming activity and section of the vagus nerves on heart rate in rainbow trout. *Journal of Experimental Biology*, 60, 305–319.
- Priede, I. G. (1983). Heart rate telemetry from fish in the natural environment. *Comparative Biochemistry and Physiology Part A*, 76, 515–524.
- Priede, I. G., & Tytler, P. (1977). Heart rate as a measure of metabolic rate in teleost fishes; *Salmo gairdneri*, *Salmo trutta* and *Gadus morhua*. *Journal of Fish Biology*, 10, 231–242.
- Prystay, T. S., Eliason, E. J., Lawrence, M. J., Dick, M., Brownscombe, J. W., Patterson, D. A., ... Cooke, S. J. (2017). The influence of water temperature on sockeye salmon heart rate recovery following simulated fisheries interactions. *Conservation Physiology*, 5, 1–12.
- Raby, G. D., Doherty, C. L. J., Mokdad, A., Pitcher, T. E., & Fisk, A. T. (2020). Post-exercise respirometry underestimates maximum metabolic rate in juvenile salmon. *Conservation Physiology*, 8, coaa063.
- Reale, D., Garant, D., Humphries, M. M., Bergeron, P., Careau, V., & Montiglio, P.-O. (2010). Personality and the emergence of the pace-of-life syndrome at the population level. *Philosophical Transactions of the Royal Society B*, 365, 4051–4063.
- Ricklefs, R. E., & Wikelski, M. (2002). The physiology/life-history nexus. *Trends in Ecology & Evolution*, 17, 462–468.
- Satchell, G. H. (1991). *Physiology and form of fish circulation*. Cambridge, UK: Cambridge University Press.
- Scharold, J., & Gruber, S. H. (1991). Telemetered heart rate as a measure of metabolic rate in the lemon shark, *Negaprion brevirostris*. *Copeia*, 1991, 942–953.
- Secor, S. (2009). Specific dynamic action: a review of the postprandial metabolic response. *Journal of Comparative Physiology B*, 179, 1–56.
- Steele, S. C., Van Leeuwen, T. E., Brownscombe, J. W., Cooke, S. J., & Eliason, E. J. (2019). An appetite for invasion: digestive physiology, thermal performance and food intake in lionfish (*Pterois* spp.). *Journal of Experimental Biology*, 222, jeb209437.
- Sundin, J., Amcoff, M., Mateos-González, F., Raby, G. D., & Clark, T. D. (2019). Long-term acclimation to near-future ocean acidification has negligible effects on a range of energetic attributes in a juvenile coral reef fish. *Oecologia, Oecologia*, 689–702.
- Thorarensen, H., Gallagher, P. E., & Farrell, A. P. (1996). The limitations of heart rate as a predictor of metabolic rate in fish. *Journal of Fish Biology*, 49, 226–236.
- Tomlinson, S., Arnall, S. G., Munn, A., Bradshaw, S. D., Maloney, S. K., Dixon, K. W., & Didham, R. K. (2014). Applications and implications of ecological energetics. *Trends in Ecology & Evolution*, 29, 280–290.
- Webber, D. M., Boutilier, R. G., & Kerr, S. R. (1998). Cardiac output as a predictor of metabolic rate in cod *Gadus morhua*. *Journal of Experimental Biology*, 201, 2779–2789.
- Winberg, G. G. 1960. Rate of metabolism and food requirements of fishes. Fisheries Research Board of Canada Translation Series No. 194. 285 pp.
- Zrini, Z. A., & Gamperl, A. K. (2021). Validating star-Oddi heart rate and acceleration data storage tags for use in Atlantic salmon (*Salmo salar*). *Animal Biotelemetry*, 9, 12.

**How to cite this article:** Doherty, C. L. J., Fisk, A. T., Cooke, S. J., Pitcher, T. E., & Raby, G. D. (2021). Exploring relationships between oxygen consumption and biologger-derived estimates of heart rate in two warmwater piscivores. *Journal of Fish Biology*, 1–8. <https://doi.org/10.1111/jfb.14923>