

# Seasonal effects on the acceleration of largemouth bass and Northern Pike in Toronto harbour

Received: 12 May 2025

Accepted: 2 January 2026

Published online: 17 January 2026

Cite this article as: Hlina B.L., Rous A.M., Piczak M.L. *et al.* Seasonal effects on the acceleration of largemouth bass and Northern Pike in Toronto harbour. *Anim Biotelemetry* (2026). <https://doi.org/10.1186/s40317-026-00444-6>

Benjamin L. Hlina, Andrew M. Rous, Morgan L. Piczak, Jonathan D. Midwood, Jacob W. Brownscombe, Rick J. Portiss, Thomas F. Sciscione, Mathew G. Wells, Susan E. Doka & Steven J. Cooke

We are providing an unedited version of this manuscript to give early access to its findings. Before final publication, the manuscript will undergo further editing. Please note there may be errors present which affect the content, and all legal disclaimers apply.

If this paper is publishing under a Transparent Peer Review model then Peer Review reports will publish with the final article.

# Seasonal Effects on the Acceleration of Largemouth Bass and Northern Pike in Toronto

## Harbour

### Animal Biotelemetry

Benjamin L. Hlina<sup>a1\*</sup>, Andrew M. Rous<sup>a1</sup>, Morgan L. Piczak<sup>a</sup>, Jonathan. D. Midwood<sup>b</sup>, Jacob. W. Brownscombe<sup>b</sup>, Rick J. Portiss<sup>c</sup>, Thomas F. Sciscione<sup>c</sup>, Mathew. G. Wells<sup>d</sup>, Susan. E. Doka<sup>b</sup>, and Steven J. Cooke<sup>a</sup>

<sup>a</sup>Fish Ecology and Conservation Physiology Laboratory, Department of Biology and Environmental Studies Institute, Carleton University, Ottawa, ON K1S 5B6, Canada

<sup>b</sup>Fisheries and Oceans Canada, Great Lakes Laboratory for Fisheries and Aquatic Science, 867 Lakeshore Road, Burlington, ON, Canada

<sup>c</sup>Toronto and Region Conservation Authority, 101 Exchange Avenue Vaughan, Ontario L4K 5R6, Canada

<sup>d</sup>Physical and Environmental Sciences, University of Toronto Scarborough, 1265 Military Trail, Toronto, ON M1C 1A4, Canada.

<sup>1</sup>Co-first Authors.

\*Corresponding author: benjamin.hlina@gmail.com; +1 (873) 354-2354

**Keywords:** Bioenergetics, Movement Ecology, Urbanization

## Abstract

**Background:** Habitats are heterogeneously distributed across landscapes, and animals move through them to access different resources, resulting in energetic consequences. In the Laurentian Great Lakes, nearshore areas provide important habitat for the majority of fish species; however, differences in the distribution and composition of physical (e.g., vegetation, depth, fetch, and substrate type/size and temperature) habitat attributes create habitat heterogeneity. We used high-resolution acoustic telemetry data combined with acceleration sensors to test how spatial and temporal factors relate to acceleration of two ecologically and economically important fish species, Largemouth Bass (*Micropterus nigricans*) and Northern Pike (*Esox lucius*), within a system of coastal embayments in Toronto Harbour, Lake Ontario.

**Results:** For both species, we found that sites characterized by higher vegetation cover, shallower depths, and decreased fetch (i.e., coastal vegetated and wetland sites) were associated with lower acceleration values. We also identified several areas within Toronto Harbour that likely serve as important movement corridors between higher suitability sites. Temporal variation of acceleration was significant for both species and was influenced by season and diel period. Both Largemouth Bass and Northern Pike exhibited increased acceleration during spring and summer. Both species exhibited peak acceleration at dawn and dusk.

**Conclusion:** The use of accelerometers was instrumental in identifying important habitats, movement corridors, and elucidating temporal trends across seasons and diel periods. Our results highlight the importance of incorporating habitat connectivity between habitat patches used for different purposes into conservation and restoration planning.

## Introduction

Locating suitable habitat is one of the most critical external factors influencing animal movement. Movement decisions result from the interaction between an individual's internal state (i.e., motivation and readiness to move), navigation and motion capacity, and external environmental factors [1,2]. Theoretically, animals move to acquire energy at a greater rate than they expend it, allowing for a surplus that can be allocated to fitness-enhancing processes such as growth or reproduction [3–5]. Environmental heterogeneity, including variation in food availability, temperature, predation risk, and movement costs, can influence animal behavior, including acceleration [6–9]. For instance, in lakes with smaller or less accessible prey, predators exhibited reduced resting periods and higher overall acceleration [10]. Consequently, animals adjust their acceleration in response to landscape variation, optimizing their movement strategies accordingly [7]. Animals inhabiting small, fragmented, or suboptimal habitats often face increased movement costs due to the necessity of traveling longer distances between quality habitats, whether on a daily timescale (e.g., during crepuscular periods) or seasonally [11]. These increased movement costs can contribute to the spatial and temporal structuring of populations and communities [7]. Movements can range from daily foraging routines within an area of core use to less frequent, longer-distance travel between territories over several days [12].

Regardless of temporal scale, (i.e., seconds, hours, or days), acceleration represents a substantial portion of the energy budget for an animal [13–15]. Standard metabolism (SMR) is considered a primary function and basic maintenance for ectotherms (e.g., fish) and typically requires a minimal amount of energy [13,16,17]. Ectotherms rely on temperature to regulate SMR, with warm temperatures resulting in an increase in the rate of physiological processes. This increase results in a greater demand on fundamental systems (e.g., cardio-respiratory systems)

causing an exponential increase in SMR causing less available energy for active and maximum metabolism (e.g., energy used for locomotion). Considering anthropogenic impacts to habitat and a warming climate, active metabolism (i.e., measured through acceleration) is likely to be altered, which can influence survival, growth, and reproduction. As movement patterns are shaped by the need to access suitable habitats that keep SMR low, understanding how species use different environments is crucial for predicting their spatial distribution and ecological interactions. Because acceleration reflects the energetic costs of movement, analyzing these patterns across space and time can highlight habitats that either elevate or reduce energetic demand, providing insight into how fish interact with their environment [18]. Such spatial patterns can help outline ‘energy landscapes’ that reveal where individuals forage, rest, or seek refuge [7], offering a management-relevant view of habitat function. These insights support habitat prioritization and conservation planning in systems facing ongoing environmental change.

The coastal wetlands in the Laurentian Great Lakes support great aquatic biodiversity and provide spawning, refugia, nursery, overwintering, and foraging habitat to many native fishes including Largemouth Bass (*Micropterus nigricans*) and Northern Pike (*Esox lucius*) [19]. Largemouth Bass are classified as a warmwater species throughout North America. Adult Largemouth Bass typically prefer shallow (less than 6m deep) areas that support submerged aquatic vegetation (SAV) or other submerged cover (i.e., logs, brush piles, and root wads). Largemouth Bass often require deeper areas (up to 15m) for overwintering habitat [20,21]. During summer, they are most abundant in areas with 40-60% vegetation cover, as higher vegetation densities can limit prey capture [21,22]. Adult Largemouth Bass are primarily piscivorous [23] and they use a variety of foraging strategies, including actively searching for prey [24,25] as well as sit-and-wait “ambush” style predation [26]. Northern Pike are considered a mesothermal or

‘coolwater’, apex predator throughout the circumpolar range of the species. The specific habitat requirements of Northern Pike are well documented with adults generally found in relatively shallow water in summer, usually  $\leq 4$  m with some occupancy at depths  $\leq 12$  m in relatively clear, cool, and well oxygenated areas [27–29]. In winter, Northern Pike tend to move to deeper areas with the onset of inshore ice cover (e.g., up to 8m deep in Lake Ontario) [30–32]. Throughout the year they are generally abundant at intermediate (30-80% cover) vegetation densities with smaller individuals occupying greater vegetation densities than larger individuals [22,27]. Northern Pike use aquatic macrophytes for multiple reasons (e.g., refuge from predation and/or forage fish of all sizes) at different life stages [28]. Similar to Largemouth Bass, Northern Pike are also ambush predators [33] where they are often associated with structurally complex habitat [34].

In addition to using biotelemetry to study habitat use, acceleration sensors within acoustic transmitters have become a valuable tool to examine differences across fish species and diverse aquatic environments [18,35–37]. Despite these advancements, few telemetry studies have tracked acceleration of fish in northern temperate lakes over an entire annual cycle [38] but see [35,39,40]. To address this gap, we captured near-continuous telemetry data with associated acceleration to gain insight into the timing, nature, and spatial distribution of fish behavior across a range of habitat types in an urban embayment of Lake Ontario. Our study provides novel information on acceleration of ecologically and economically important Largemouth Bass and Northern Pike in Toronto Harbour, Lake Ontario—a system with a diverse array of habitat types but also pressures [41,42]. We generated predictions of both species’ acceleration across space and time within Toronto Harbour. By exploring when and where these behaviors occur, we provide a deeper understanding of how these species interact with and navigate their environment. Broadly, this study aims to advance our understanding of acceleration in fish and serves as a case study

demonstrating how biotelemetry data can reveal patterns of potential fish-habitat interactions and behaviour at different scales.

## **Methods**

### ***Study Site***

Toronto Harbour is a large (~15 km<sup>2</sup>) and heterogenous aquatic system connected to Lake Ontario that contains exposed shorelines, protected embayments, coastal wetlands, rivermouths and port facilities (i.e., industrial and recreational boat slips and associated traffic). It is situated directly adjacent to the downtown core of Toronto, ON, Canada (~5 million people) and consists of habitats with varying degrees of anthropogenic degradation. Toronto Harbour is a heavily urbanized area comprised of four zones: Inner Harbour, Toronto Islands, Outer Harbour, and Tommy Thompson Park (TTP). Two zones dominate the Inner Harbour, the city waterfront (urban and industrial landscape with hardened shorelines), which is connected to the Toronto Islands (a series of channels and islands with more naturalized shorelines). The Outer Harbour is bounded by industrial and recreational uses, and TTP, which is an interconnected series of largely, renaturalized embayments. The eastern gap (a channel) joins the Inner and Outer harbours, and both Harbours are directly connected to Lake Ontario proper: one lake connection is via the western gap channel connecting the Inner Harbour to Humber Bay, the confluence between the Humber River and Lake Ontario. While the Outer Harbour is an embayment directly open to the Lake to the southwest. TTP is a man-made peninsula that was constructed in the early 1970's by using ecological restoration techniques to improve the quality of aquatic habitat, with current restoration permitted through infilling. The peninsula is made from clean fill materials and has been modified to naturalize portions of Toronto Harbour, and restore lost coastal features, as well as house contained disposal facilities until they are capped and enhanced naturally. This large

aquatic and terrestrial park extends 5 km into Lake Ontario covering a total surface area of over 250 ha [43].

### *Acoustic Telemetry Array*

An acoustic receiver array was first deployed in Toronto Harbour in 2010 as part of a broader telemetry effort to monitor fish and restoration activity. The present study represents a focused component of that larger initiative and was conducted between August 2014 and August 2016 (see [44]). For this study, 45 acoustic telemetry receivers (VR2W, 69 kHz, Innovasea, Halifax, Nova Scotia) were strategically positioned throughout the harbour to cover a variety of habitat types and environmental gradients, as well as key movement corridors, across 22 different groups (Figure 1; see Additional File 1). In shallow areas (<5 m), acoustic receivers were attached to a rope approximately 1 m above a steel or concrete anchor with a Castro float at the top to keep the receiver positioned vertically. Anchors were tethered to the nearest attachment point on shore by submerged steel cable. In deeper water (up to 10 m), the anchor was connected by floating rope to an additional weight approximately 20 m away from the primary anchor weight. Receivers were retrieved every 6 months to download data and to remove any accumulated biofouling and check receiver condition. Receivers were then redeployed in the same locations. Range testing (see Kessel et al. 2014) was conducted at a subset of receivers in different habitat types and in different seasons to inform receiver placement; ranges varied from 400 to 1500 m [45].

### *Fish Tagging*

All fish in this study were captured via boat electrofishing (SR-18EH, 6.0–7.0 A, 60 Hz, 340V DC, Smith-Root, Inc., Vancouver, WA) between August and September 2014 and 2015. After capture, each fish was held in a live well until the boat could be safely docked and set up for transmitter implantation. All individuals were tagged with Vemco (now Innovasea) V13A

transmitters [69kHz, 90 s transmission delay,  $\pm 3.43$  g acceleration range, 5 Hz sampling frequency, activity algorithm (root mean square of acceleration =  $\sqrt{(x^2+y^2+z^2)}$  per transmitter specification sheet, see below for more information, Vemco Ltd., Bedford, Nova Scotia). Fish were immobilized throughout the entire duration of the surgery by holding each individual in a wetted mesh net between the boat electrofisher electrodes (Smith-Root, Inc., Vancouver, WA) – a method that has previously been used to immobilize fish for surgeries (akin to electrostunning; [46]). For transmitter implantation, fish were moved from the live well with a wetted net onto a padded surgical table with the fish in a supine position. Each individual was measured for total length (mm). Prior to implanting an acoustic transmitter into an individual, the transmitter and all surgical tools were disinfected in an iodine solution and rinsed. An incision ( $< 20$  mm) was made with a scalpel on the ventral surface of the fish. Curved forceps were used to lift the skin and body wall to avoid any internal injury while making the incision. The transmitter was inserted into the coelomic cavity of the fish. The incision was closed using two simple, interrupted sutures (Ethicon PDS II, 3/0, FSL needle). After surgery, fish were returned to the livewell for recovery, and once recovered, fish were released at their original capture location. Over the course of the study, 37 individual fish were tagged: 2014, 5 Largemouth Bass and 5 Northern Pike and, and in 2015, 12 Largemouth Bass and 15 Northern Pike (see Additional File 1). Fish handling and surgical procedures were approved and followed a Canadian Council on Animal Care protocol administered by Carleton University (Certificate CU 110723).

### ***Data Processing***

Fish detections were filtered to remove any potential false detections (i.e., detections that occurred from the same tag at the same receiver within a period of less than the minimum tag delay; [47]). Furthermore, a swim-speed filter was applied to the detection data to remove detections that

consisted of improbable distances among detections (e.g., two detections that were 5 mins apart on two receivers that are 2.0 kms apart from each other). Detections of individual tags were plotted over space and time to visually examine stationary tags (i.e., those not tracking live fish due to fish mortality or tag shedding) and subsequently removed from the dataset. Transmitters measure dynamic acceleration (i.e., acceleration caused by the movement of an animal) and not total acceleration. The transmitter has an onboard processor that first filters out the static component (i.e., gravity and orientation) from the raw acceleration data, leaving only dynamic acceleration. The transmitter then calculates the root mean square [i.e.,  $\sqrt{(x^2+y^2+z^2)} = \text{RMS}$ ] of this dynamic acceleration over the sampling time period that is then transmitted as the “activity” output. This value dynamic acceleration is transmitted to receivers in a raw unitless value (1-255; ADC) that corresponds to a calibration model provided by Innovasea. These values were transformed into  $\text{m s}^{-2}$  using the following equation,  $y = 0 + 0.013588x$ , where ADC values were substituted for  $x$ . For example, if the transmitter is completely still (i.e., no motion, no vibration), the raw accelerometer sensor experiences 1 g ( $9.8 \text{ m/s}^2$ ) from gravity but that static acceleration is removed by the onboard filtering process before the RMS acceleration is calculated. Because the tag filters out slow, low-frequency changes caused by gravity and orientation, the remaining dynamic acceleration signal is essentially zero when the transmitter is motionless [48].

### ***Data analysis***

Receivers were either treated as a unique station ( $N = 14$ ) or the individual receivers were combined into groups ( $N = 9$ ) that represented locally homogeneous areas with respect to habitat features and had overlapping fields of detections (Figure 1). For example, at the connection point between the Outer Harbour and Lake Ontario, there were seven receivers deployed as a gate to track fish that may exit the telemetry array for the open lake. These receivers covered a similar

habitat type and were in close proximity. Data from all of these receivers were therefore integrated into a single receiver group to represent a “curtain” of receivers (Figure 1). Herein, both the unique stations and groups of receivers are collectively referred to as “receiver groupings” (see Additional File 1 for details). Importantly, combining receivers into “groups” also increased the total detection area relative to a single receiver (Figure 1).

### *Habitat Types*

Habitats were classified by integrating spatial habitat layers (see [44] for full details). Briefly, Fisheries and Oceans Canada provided a digital elevation model (DEM) for Toronto Harbour. From this DEM, the elevation gradient (slope) was calculated in ArcMap 10.2 (Environmental Systems Research Institute, Redlands, CA, USA; [49]) Mean exposure (i.e., effective fetch) was calculated for the entire harbour at a 10 m<sup>2</sup> grid cell size using a wind-fetch model developed by the United States Geological Survey [50]. An estimate of the percent cover of SAV (calculated using DEM, slope and mean exposure) was generated from an equation developed for Hamilton Harbour, an urban harbour located approximately 40 km southwest of Toronto Harbour on Lake Ontario [51]. Finally, mean benthic water temperature during stratification (June to September) was used for the temperature variable [52]. Only this period was selected since there was considerable spatial variability in temperatures whereas during the pre- and post-stratification periods, the water had uniform temperatures, with minimal variation with depth or spatially throughout the harbour [52]. These habitat layers were then used to generate five habitat categories derived within Toronto Harbour by [44]: 1) wetlands (shallow, low fetch, warm, vegetated), 2) coastal vegetated (moderate depth, moderate fetch, cool-warm, vegetated), 3) shallow un-vegetated (shallow, low fetch, cool-warm, no SAV), 4) deep un-vegetated (deep, moderate fetch, cool, no SAV), and 5) exposed (moderate-deep, high fetch, cool-warm, no SAV; see Additional

File 2 for receiver array details). Seasons were defined as spring (1 March until 31 May), summer (1 June until 31 August), fall (1 September until 30 November), and winter (1 December – 28 February).

### *Acceleration*

All statistical analyses were conducted using R version 4.3.2 (R Core Team 2023). The full detection database was grouped into 20-minute time bins to consider fine-scale movement among receivers for each individual and the mean acceleration and location (longitude, latitude) were calculated to create centers of activity (COA; Additional File 3). Considering the detection range of receivers did not overlap, we did not involve the first-detection heard at a receiver in our analysis as we were unsure where that acceleration was measured. Subsequent detections (e.g., 10 detections at a given receiver) were retained and used, because we were certain transmissions occurred near the receiver. To estimate spatial differences in acceleration, we modeled the mean acceleration derived from the COAs using a Generalized Additive Mixed-effects Model (GAMM; using the package {mgcv}; [53]) Prior to modelling, the mean acceleration was transformed using a power transformation to normalize the data (Largemouth Bass – skewness = 2.25, kurtosis = 9.76; Northern Pike – skewness = 6.88, kurtosis = 87.14). The model for Largemouth Bass used a Gamm error distribution with a link function set to log with the model structure being the following:

$$TA_i \sim \text{Gamma}(\mu_i, \phi)$$

$$\log(\mu_i) = \beta_0 + \beta_{\text{season}[i]} + f_{\text{spatial}}(\text{longitude}_i, \text{latitude}_i) + b_{\text{tag}[i]} + \varepsilon_i$$

With transformed acceleration ( $TA_i$ ) as the response, season as a fixed effect (i.e., non-ordered

factor;  $\beta_{\text{season}[i]}$ ), longitude and latitude (derived from COAs) as a spatial smoother [thin-plate regression spline;  $f_{\text{spatial}}(\text{longitude}_i, \text{latitude}_i)$ ], fish-tag number as a random effect [ $b_{\text{tag}[i]} b_{\text{tag}} \sim \mathcal{N}(0, \sigma_{\text{tag}}^2)$ ], and an autocorrelation structure defined below ( $\varepsilon_i$ ). The models for Northern Pike used a Gaussian error distribution with a link function set to log with the model structure being the following:

$$TA_i \sim \mathcal{N}(\mu_i, \phi)$$

$$\log(\mu_i) = \beta_0 + \beta_{\text{season}[i]} + f_{\text{spatial}}(\text{longitude}_i, \text{latitude}_i) + b_{\text{tag}[i]} + \varepsilon_i$$

The smooth terms were defined with an appropriate number of basis functions to balance model flexibility with interpretability, reduce wiggleness and prevent overfitting (as per [54]). Models included an autocorrelation structure ( $\varepsilon_i$ ) that was developed using the package `{itsadug}` and supplied the correlation matrices for rho [55] using the following:

$$\eta_i \sim \mathcal{N}(0, \sigma^2)$$

$$\varepsilon_i = \rho\varepsilon_{i-1} + \eta_i$$

Model validation was conducted using the package `{gratia}` for GAMMs [56]. The goal of the model section was to assess whether statistical models that included season (i.e., presumably more complex) indeed were better predictors of acceleration values for Largemouth Bass and Northern Pike.

To determine whether season and diel period influenced largemouth bass acceleration on a more discrete timescale (i.e., season and diel period), we used generalized linear mixed model (GLMM) with a Gamma distribution with the link function set to inverse, while acceleration of

northern pike on the discrete timescales was analyzed using a lognormal distribution with a link function set to log. Diel periods were defined using the package {suncalc} into four different time periods dawn, day, dusk, and night, with the duration of each period shifting throughout each season (e.g., day length during summer is greater than during the winter; [57]). We created four separate models to assess whether discrete temporal scales (i.e., season or diel period) influenced acceleration for these two species. These models had the following structure of acceleration as the response variable and season or diel period as categorical predictors. We created separate models for each species considering general behaviour and life-history strategies for each species are different (see [58]). An autocorrelation structure and random effect of individual fish number were included to account for autocorrelation and individual variation. GLMMs were fit using the R package {glmmTMB} [59]. Models were assessed to fit the data using {DHARMA} [60].

## Results

Northern Pike represented approximately 84% of the total detections (858,610) and Largemouth Bass 16% (164,047). Individuals were tracked for an average of 68% of the days in the study period (range = 38 – 374 days; Additional File 2). The number of detections for each tagged individual varied from 122 – 101,161 detections (mean = 27,639). On average, tagged individuals throughout the study period were detected on seven receivers: from a minimum of one and a maximum of 24 receivers. Basic seasonal movement can be inferred by evaluating the number of detections at each receiver and the number of unique individuals that were detected at each receiver which are displayed in Figure 2.

### *Largemouth Bass*

We found that mean acceleration of Largemouth Bass was significantly influenced by spatial

differences among seasons (GAMM spatial smoother using latitude and longitude by the fixed effect of season— $F = 0.5\text{--}2596.7$ ,  $\text{edf} = 0.3\text{--}9.8$ ,  $\text{ref. df} = 10$ ,  $p \leq 0.001$ ; Figure 3A). Spatially, there was minimal variation in acceleration across Toronto Harbour during winter and spring, with bass exhibiting relatively low mean values of  $\sim 0.3 \text{ m s}^{-2}$  in winter and  $\sim 0.6 \text{ m s}^{-2}$  in spring, regardless of habitat type. In contrast, summer showed much greater spatial variation, with the highest acceleration observed in deep, unvegetated areas throughout both the Inner and Outer Harbour (exceeding  $1.5 \text{ m s}^{-2}$ ). Three major areas of high acceleration occurred in Toronto Harbour for Largemouth Bass with two of them being located at the Eastern and Western Gaps which are both deep and unvegetated, where mean acceleration exceeded  $1.0 \text{ m s}^{-2}$ . The last area that resulted in Largemouth Bass exhibiting high acceleration occurred along the northeast developed coastline (unvegetated) of the Inner Harbour (IH). In comparison, sites within TTP and the Toronto Islands, characterized by shallow, vegetated habitats and reduced wind exposure, had lower acceleration values. During the fall, the highest accelerations ( $\sim 0.6 \text{ m s}^{-2}$ ) occurred in the Outer Harbour, suggesting increased acceleration in these deeper, more exposed areas.

Largemouth bass acceleration varied across seasons ( $\chi^2_{1,3} = 13840$ ,  $p \leq 0.001$ ; Table 1; Figure 4a; see Additional File 3 for centers-of-activity) and diel periods ( $\chi^2_{1,3} = 2606$ ,  $p \leq 0.001$ ; Table 1; Figure 4c). Acceleration levels for largemouth bass followed general seasonal trends with acceleration lower throughout the winter ( $0.172 \pm 0.001 \text{ m s}^{-2}$ ) and fall ( $0.277 \pm 0.002 \text{ m s}^{-2}$ ) compared to spring and summer ( $0.572 \pm 0.013$ ;  $0.642 \pm 0.004$ , respectively; Figure 3b). Largemouth bass acceleration peaked during dawn ( $0.521 \pm 0.012 \text{ m s}^{-2}$ ) and slowly decreased throughout the day with acceleration being the lowest at night ( $0.278 \pm 0.002 \text{ m s}^{-2}$ ; Figure 3c).

### *Northern Pike*

We found that mean acceleration varied across space by seasons for northern pike (GAMM spatial smoother by season— $F = 0.06-352.47$ ,  $\text{edf} = 0.42 - 49.35$ ,  $\text{ref. df} = 18.0 - 99.00$ ,  $p \leq 0.001$ ; Figure 3B). During the winter, there was minimal variation in acceleration, where values remained below  $0.2 \text{ m s}^{-2}$  throughout most of Toronto Harbour. We observed higher acceleration values for Northern Pike in the Outer Harbour during winter, particularly near the confluence with Lake Ontario, where fish moved faster in exposed habitats with minimal vegetation (Figure 3B). In spring, acceleration remained low across most of Toronto Harbour, except in the Outer Harbour, where values reached  $\sim 0.3 \text{ m s}^{-2}$  in deep, unvegetated areas. Similar to Largemouth Bass, Northern Pike exhibited the greatest spatial variability in acceleration during summer, with the highest values occurring in the Outer Harbour near the Eastern Gap. In contrast, acceleration values were lower throughout the Toronto Islands and TTP, which were characterized by shallow depths, high vegetation cover, higher temperatures, and reduced wind exposure (i.e., wetlands and coastal vegetated habitats). During the fall, acceleration values were largely uniform across the Harbour, generally remaining below  $0.2 \text{ m s}^{-2}$ . As in other seasons, the highest fall values were again found in deep, unvegetated habitats of the Outer Harbour (Figure 2B).

Northern Pike acceleration varied across seasons ( $\chi^2_{1,3} = 1937.80$ ,  $p \leq 0.001$ ; Table 1; Figure 4b; see Additional File 3 for multiple comparisons) and diel periods ( $\chi^2_{1,3} = 2644.90$ ,  $p \leq 0.001$ ; Figure 4d; Table 1). Seasonal trends in acceleration levels for northern pike were generally lower throughout the winter ( $0.114 \pm 0.001 \text{ m s}^{-2}$ ) and fall ( $0.140 \pm 0.001 \text{ m s}^{-2}$ ) compared to spring and summer ( $0.176 \pm 0.001$  and  $0.175 \pm 0.001$ , respectively; Figure 4b). Northern pike acceleration peaked at dusk ( $0.212 \pm 0.003 \text{ m s}^{-2}$ ) and dawn ( $0.178 \pm 0.003 \text{ m s}^{-2}$ ) compared to during the day ( $0.153 \pm 0.001 \text{ m s}^{-2}$ ) and night ( $0.121 \pm 0.001 \text{ m s}^{-2}$ ; Figure 4d).

## Discussion

Using acoustic telemetry with accelerometer sensors, we examined how acceleration varied across space and time for both Largemouth Bass and Northern Pike within a heterogeneous system of coastal embayments. We found that acceleration was highest at sites throughout Toronto Harbour that were deep and unvegetated, with elevated values particularly evident in the Outer Harbour across all seasons. For both species, acceleration was influenced by season and diel period, with values increasing during spring and summer, and peaking around dawn and dusk (crepuscular). Acceleration patterns for Largemouth Bass and Northern Pike, suggested that exposed, deep, and unvegetated areas may function as movement corridors connecting patches of preferred habitat. This interpretation aligns with previous research showing low residency and occupancy at exposed sites lacking vegetation [44] and with the observed spatial separation between high-quality vegetated habitats across Toronto Harbour [42]. Several high-acceleration areas also corresponded with regions of intense water exchange and current velocity (e.g., upwellings; [61]), such as the Eastern and Western Gaps, indicating that both species may increase swimming effort and therefore likely energy expenditure in these dynamic environments to maintain position or move between preferred habitat patches [61]. Although both species are generally associated with vegetated habitats, elevated acceleration was also observed at some vegetated coastal sites, particularly for Largemouth Bass, which may serve as transitional zones between areas of low acceleration. For example, vegetated sites north of the Toronto Islands lie between sheltered wetland habitats and the more exposed Inner Harbour, potentially requiring more active movement. It is important to note that acceleration values were averaged across 20-minute intervals, which captures broader movement effort but may not detect short bursts of high activity associated with behaviors such as foraging. As such, the high-acceleration areas identified

here are more likely to reflect sustained swimming or transit between habitats rather than short-burst predation events. Together, these findings emphasize the importance of transitional zones and suggest that both species use a combination of habitat types, including high-acceleration corridors, to navigate the heterogeneous landscape. To better understand the energetic consequences of these high acceleration movements in areas of high flow, future work should calibrate acceleration values to metabolic cost at varying thermal conditions using swim tunnel experiments (e.g., [62]). These measurements could help facilitate estimating yearly energetic budgets that rely on relating standard and active metabolism to food consumption and thus providing an index of available energy for survival, growth, and reproduction [3,63]. This would be especially valuable in an urbanized system like Toronto Harbour, where habitat heterogeneity and degradation may impose increased energetic demands on resident fish [64].

Acceleration values for both species remained relatively low throughout the year in the Toronto Islands and TTP, areas characterized by vegetated habitats such as wetlands and coastal vegetation. These findings align with previous research in Toronto Harbour showing high occupancy of both Largemouth Bass and Northern Pike in these locations, indicating consistent habitat use [22]. Specifically, Largemouth Bass exhibited the highest occupancy within the Toronto Islands and TTP, while occupancy remained low across the Inner and Outer Harbour [22]. This pattern is consistent with known habitat preferences for Largemouth Bass, which favour areas with SAV, shallow depths, and low exposure [65,66]. Similarly, Northern Pike were also found to have high occupancy in these same areas, consistent with their known association with vegetated, shallow habitats, typically less than 12 m in depth [27,32]. In addition, Northern Pike showed high occupancy along the northwest shoreline of the Inner Harbour, including boat slips that have undergone ecological restoration specifically for this species [67]. Our results are consistent with

these prior studies, as acceleration values were predicted to remain low across seasons in these vegetated areas. Taken together, the observed low sustained acceleration values, combined with previously documented high occupancy, underscore the ecological importance of the Toronto Islands and TTP as key habitat for both Largemouth Bass and Northern Pike in this highly urbanized harbour.

We determined that acceleration for both species varied across seasons, and that acceleration was generally higher in spring and summer for both species. Increased acceleration in spring may be in preparation for reproductive activities of both Largemouth Bass and Northern Pike. During this time fish will actively seek out food and warmer water temperatures to enable gonad development needed for spawning [68–70]. Additionally, as SAV can be sparse during spring, Northern Pike and Largemouth Bass could be quite active as they search for suitable areas to forage, and stage for spawning activities, after ice-out. Acceleration continued to increase in spring and into the summer. Fish may show increased acceleration during late summer and early fall as they intensify foraging activity to build overwinter energy reserves [71,72] or to locate overwintering regions with suitable water quality conditions (e.g., [73]). We did, however, determine that acceleration of both species decreased substantially in fall as fish resumed low acceleration as water temperatures became cold and isothermal. Each species also exhibited decreased acceleration during the winter, particularly Largemouth Bass, although neither species were completely quiescent. Average acceleration was slightly higher ( $0.17 \pm 0.0013 \text{ m s}^{-2}$ ) for Largemouth Bass compared to Northern Pike during winter ( $0.11 \pm 0.0005 \text{ m s}^{-2}$ ); however, Northern Pike did exhibit a greater number of isolated, high-acceleration events, consistent with their overall acceleration patterns throughout the rest of the year (Additional File 3). Overall, our findings highlight that although both species reduced acceleration during colder months, seasonal

and species-specific differences in acceleration were evident, reflecting underlying life history traits and habitat use strategies.

Largemouth Bass and Northern Pike acceleration were also influenced by diel periods. Largemouth Bass had the highest acceleration levels at dawn, followed by day and dusk, with the lowest values at night, although there were movements undertaken at all times of the day. These observations are similar to a previous study that determined that Largemouth Bass acceleration levels were higher during the day compared to night [74]. Higher foraging success is likely driving increased activities for Largemouth Bass during periods of more intense daylight as previous research has found that feeding drastically decreased during night [75]. Northern Pike were also found to have increased acceleration levels during dawn and dusk across all seasons, but were more pronounced during the summer, similar to other studies [40,76]. Higher acceleration by Northern Pike during low-light conditions is likely due to increased foraging because reaction time of prey is diminished with decreased light, conditions that are favourable for ambush predators (see [77]) These patterns suggest that diel variation influences both foraging activity and broader movements between foraging areas, with peaks in acceleration likely reflecting periods when individuals optimize foraging efficiency while minimizing energetic costs. Both species are piscivorous, consuming small-bodied fish such as Yellow Perch (*Perca flavescens*) [23,78]; future studies could further examine foraging movements on a fine scale for both predators and prey (e.g., with YAPS; [79]) particularly given that technological advances in acoustic telemetry are now making this possible [80].

Our study had several limitations, which should serve as sources for future research. First, while acceleration is useful for determining speed of animals in relation to their habitat, it does not distinguish among specific behaviors such as predator avoidance, or reproduction [81]. Future

studies integrating acceleration with additional environmental or physiological data could help refine interpretations of movement behavior. Next, the number of detections was much lower for Largemouth Bass relative to Northern Pike, which could have influenced our analysis and interpretations. Future research would benefit from increased sample sizes and longer monitoring periods to maximize the comprehensiveness of assessing acceleration over space and time. The habitat types were assigned at individual receiver stations; however, it would have been beneficial to have higher temporal resolution for the vegetation, depth, and exposure values to ensure adequate coverage for the fish movements. While all values for vegetation, depth, and exposure were static, ideally there would be vegetation estimates across seasons, to capture variability both within year and interannually (see [82]). Additionally, the acoustic receiver array was designed to capture specific habitats and features, as well as navigation needs within Toronto Harbour; however, examination of accelerations across space would be more conducive to a grid design where receivers are spread uniformly across space (see [83]), to minimize bias introduced by the locations of the receivers [84]. Environmental variables beyond habitat type, such as water exchange (e.g., upwelling events) and prey availability, may also contribute to spatial and seasonal variation in acceleration but were not directly measured in this study.

### **Management Implications and Conclusion**

Our findings of differing acceleration patterns in both Largemouth Bass and Northern Pike have implications for habitat management and conservation. We have built upon previous research [18], demonstrating the utility of incorporating acceleration sensors into acoustic telemetry studies. Here, accelerometers complemented standard movement information to elucidate which habitats within Toronto Harbour were used as movement corridors between preferred habitats, as demonstrated by high accelerometer values. As previously noted, aquatic habitat is often

heterogeneously distributed in space, and this is true for Toronto Harbour. Since both species exhibited increased acceleration in exposed sites, these areas may represent energetically costly corridors rather than preferred habitats, yet they likely remain important for connectivity and transit. Consequently, restoration planning should consider the spatial configuration and distance between high-quality habitats, as movement through less suitable areas still contributes to overall connectivity across the energy landscape. For example, if wetlands are restored within a system but are disparately spaced with extensive exposed sites in between, these sites could remain fragmented due to the increased energy expenditure associated with movements between wetlands. Our results suggest that areas associated with high acceleration may act as movement corridors, allowing fish to travel between essential habitats. Therefore, habitat restoration should not only focus on creating or enhancing habitat patches but also consider how fish move between these sites, including the energetic costs of these movements across different habitat types (see [85]). Additionally, given that both species had lower acceleration in coastal vegetated and wetland habitats with higher SAV cover, these habitats should be prioritized during conservation planning. Specifically, our findings regarding acceleration, in combination with previous research [22], demonstrate the importance of these habitats to both species, particularly in urbanized areas where high-quality habitat may be scarce [64]. Therefore, wetlands and coastal vegetated habitats should be prioritized for both protection and ecological restoration for Largemouth Bass and Northern Pike (see [86,87]). Conserving a diversity of habitat types and maintaining functional connectivity through both high-use corridors and refuge areas will be critical for sustaining fish populations within increasingly fragmented urban environments unless active restoration is included in planning [88,89].

### **Acknowledgements**

Thank you to the staff at the Toronto and Region Conservation Authority for assisting with receiver deployment and maintenance. We would also like to thank the two anonymous reviewers for the contributions in refining this work.

### **Funding**

MLP is supported by Natural Sciences and Engineering Research Council of Canada and Liber Ero.

### **Author Contributions:**

All authors contributed to the design of the study. Field work and sampling were conducted by A.M.R. Data analyses and figures were done by A.M.R., M.L.P, and B.L.H. The manuscript was written by A.M.R., M.L.P, and B.L.H, with all authors contributing to revisions.

### **Data availability**

Telemetry data are available at the following [GitHub repository](#).

### **Ethics, Consent to Participate, and Consent to Publish Declarations**

Not applicable

### **Competing Interest Declaration**

All authors declare that they have no competing interests.

### **References**

1. Nathan R, Getz WM, Revilla E, Holyoak M, Kadmon R, Saltz D, et al. A movement ecology paradigm for unifying organismal movement research. *Proceedings of the National Academy of Sciences*. 2008;105:19052–9. <https://doi.org/10.1073/pnas.0800375105>

2. Cooke SJ, Bergman JN, Twardek WM, Piczak ML, Casselberry GA, Lutek K, et al. The movement ecology of fishes. *Journal of Fish Biology*. 2022;101:756–79. <https://doi.org/10.1111/jfb.15153>
3. Kitchell JF. Energetics. In: Webb PW, Weihs D, editors. *Fish Biomechanics*. Praeger Publishers; 1983. p. 321–38.
4. Chipps SR, Wahl DH. Bioenergetics Modeling in the 21st Century: Reviewing New insights and revisiting old constraints. *Transactions of the American Fisheries Society*. 2008;137:298–313. <https://doi.org/10.1577/t05-236.1>
5. Sokolova I. Bioenergetics in environmental adaptation and stress tolerance of aquatic ectotherms: linking physiology and ecology in a multi-stressor landscape. *Journal of Experimental Biology*. 2021;224:1–18. <https://doi.org/10.1242/jeb.236802>
6. Lima SL, Zollner PA. Towards a behavioral ecology of ecological landscapes. *Trends in Ecology & Evolution*. Elsevier Current Trends; 1996;11:131–5.
7. Shepard ELC, Wilson RP, Rees WG, Grundy E, Lambertucci SA, Vosper SB. Energy landscapes shape animal movement ecology. *American Naturalist*. 2013;182:298–312. <https://doi.org/10.1086/671257>
8. Gallagher AJ, Creel S, Wilson RP, Cooke SJ. Energy Landscapes and the Landscape of Fear. *Trends in Ecology and Evolution*. 2017;32:88–96. <https://doi.org/10.1016/j.tree.2016.10.010>
9. Wilson RP, Quintana F, Hobson VJ. Construction of energy landscapes can clarify the movement and distribution of foraging animals. *Proceedings of the Royal Society B: Biological Sciences*. Royal Society; 2011;279:975–80. <https://doi.org/10.1098/rspb.2011.1544>
10. Cruz-Font L, Shuter BJ, Blanchfield PJ, Minns CK, Rennie MD. Life at the top: Lake ecotype influences the foraging pattern, metabolic costs and life history of an apex fish predator. *Journal of Animal Ecology*. 2019;88:702–16. <https://doi.org/10.1111/1365-2656.12956>
11. Ellis RD, McWhorter TJ, Maron M. Integrating landscape ecology and conservation physiology. *Landscape Ecol*. 2012;27:1–12. <https://doi.org/10.1007/s10980-011-9671-6>
12. Dingle H, Drake VA. What is migration? *BioScience*. 2007;57:113–21. <https://doi.org/10.1641/B570206>
13. Boisclair D, Leggett WC. Importance of activity in bioenergetics mode actively foraging fishes<sup>1</sup>. *Canadian Journal of Fisheries & Aquatic Sciences*. 1989;46:1859–67.
14. Cooke SJ, Thorstad EB, Hinch SG. Activity and energetics of free-swimming fish: Insights from electromyogram telemetry. *Fish and Fisheries*. 2004;5:21–52. <https://doi.org/10.1111/j.1467-2960.2004.00136.x>
15. Christian KA, Baudinette RV, Pamula Y. Energetic costs of activity by lizards in the field. *Functional Ecology*. 1997;11:392–7. <https://doi.org/10.1046/j.1365-2435.1997.00099.x>
16. Brett JR, Groves TDD. Physiological energetics. In: Hoar WS, Randall DJ, Brett JR, editors. *Fish Physiology: Volume VIII*. New York, New York: Academic Press, Inc.; 1979. p. 279–352.
17. Chabot D, Steffensen JF, Farrell AP. The determination of standard metabolic rate in fishes. 2016;81–121. <https://doi.org/10.1111/jfb.12845>
18. Lennox RJ, Eldøy SH, Dahlmo LS, Matley JK, Vollset KW. Acoustic accelerometer transmitters and their growing relevance to aquatic science. *Movement Ecology*. 2023;11:45. <https://doi.org/10.1186/s40462-023-00403-3>
19. Jude DJ, Pappas J. Fish utilization of Great Lakes coastal wetlands. *Journal of Great Lakes Research*. 1992;18:651–72. [https://doi.org/10.1016/S0380-1330\(92\)71328-8](https://doi.org/10.1016/S0380-1330(92)71328-8)
20. Suski C, Ridgway M. Winter biology of centrarchid fishes. *Centrarchid fishes: Diversity, biology, and conservation*. Wiley Online Library; 2009;264–92.
21. Stuber RJ, Gebhart G, Maughan OE. Habitat suitability index models: largemouth bass. *Western Energy and Land Use Team, Office of Biological Services, Fish and ...*; 1982.
22. Brownscombe JW, Midwood JD, Doka SE, Cooke SJ. Telemetry-based spatial–temporal fish habitat models for fishes in an urban freshwater harbour. *Hydrobiologia*. 2023;850:1779–800. <https://doi.org/10.1007/s10750-023-05180-z>

23. Brown T, Runciman B, Pollard S, Grant A. Biological synopsis of largemouth bass (*Micropterus salmoides*). Canadian manuscript report of fisheries and aquatic sciences. Fisheries and Oceans Canada Nanaimo, BC, Canada; 2009;2884:1–26.
24. Savino JF, Stein RA. Behavior of fish predators and their prey: habitat choice between open water and dense vegetation. *Environ Biol Fish*. 1989;24:287–93. <https://doi.org/10.1007/BF00001402>
25. Ahrenstorff TD, Sass GG, Helmus MR. The influence of littoral zone coarse woody habitat on home range size, spatial distribution, and feeding ecology of largemouth bass (*Micropterus salmoides*). *Hydrobiologia*. 2009;623:223–33. <https://doi.org/10.1007/s10750-008-9660-1>
26. Savino JF, Stein RA. Predator-prey interaction between largemouth bass and bluegills as influenced by simulated, submersed vegetation. *Transactions of the American Fisheries Society*. AFS Website; 1982;111:255–66. [https://doi.org/10.1577/1548-8659\(1982\)111%253C255:PIBLBA%253E2.0.CO;2](https://doi.org/10.1577/1548-8659(1982)111%253C255:PIBLBA%253E2.0.CO;2)
27. Casselman J. Age, growth and environmental requirements of pike. *Pike: biology and exploitation*. Springer; 1996. p. 69–101.
28. Inskip PD. Habitat suitability index models: northern pike. Western Energy and Land Use Team, Office of Biological Services, Fish and ...; 1982.
29. Headrick MR, Carline RF. Restricted summer habitat and growth of northern pike in two southern Ohio impoundments. *Transactions of the American Fisheries Society*. Taylor & Francis; 1993;122:228–36.
30. Chu C, Mandrak NE, Minns CK. Potential impacts of climate change on the distributions of several common and rare freshwater fishes in Canada. *Diversity and Distributions*. 2005;11:299–310. <https://doi.org/10.1111/j.1366-9516.2005.00153.x>
31. Minns CK, Randall RG, Moore JE, Cairns VW. A model simulating the impact of habitat supply limits on northern pike, *Esox lucius*, in Hamilton Harbour, Lake Ontario. *Canadian Journal of Fisheries and Aquatic Sciences*. NRC Research Press Ottawa, Canada; 1996;53:20–34.
32. Minns CK, Doka SE, Bakelaar CN, Brunette PC, Schertzer WM. Identifying Habitats essential for pike *Esox lucius* L. in the Long Point region of Lake Erie: a suitable supply approach. *Fish habitat: essential fish habitat and rehabilitation American Fisheries Society, Symposium*. 1999. p. 363–82.
33. Harper DG, Blake RW. Prey capture and the fast-start performance of northern pike *Esox lucius*. *Journal of experimental Biology*. The Company of Biologists Ltd; 1991;155:175–92.
34. Říha M, Rabaneda-Bueno R, Jarić I, Souza AT, Vejřík L, Draščík V, et al. Seasonal habitat use of three predatory fishes in a freshwater ecosystem. *Hydrobiologia*. 2022;849:3351–71. <https://doi.org/10.1007/s10750-022-04938-1>
35. Hlina BL, Glassman DM, Lédée EJI, Nowell LB, Claussen JE, Philipp DP, et al. Habitat-dependent metabolic costs for a wild cold-water fish. *Aquat Sci*. 2024;86:36. <https://doi.org/10.1007/s00027-024-01052-3>
36. Landsman SJ, Martins EG, Gutowsky LFG, Suski CD, Arlinghaus R, Cooke SJ. Locomotor activity patterns of muskellunge (*Esox masquinongy*) assessed using tri-axial acceleration sensing acoustic transmitters. *Environ Biol Fish*. 2015;98:2109–21. <https://doi.org/10.1007/s10641-015-0433-1>
37. O’Toole AC, Murchie KJ, Pullen C, Hanson KC, Suski CD, Danylchuk AJ, et al. Locomotory activity and depth distribution of adult great barracuda (*Sphyrnaea barracuda*) in Bahamian coastal habitats determined using acceleration and pressure biotelemetry transmitters. *Mar Freshwater Res*. 2010;61:1446. <https://doi.org/10.1071/MF10046>
38. Marra PP, Cohen EB, Loss SR, Rutter JE, Tonra CM. A call for full annual cycle research in animal ecology. *Biology Letters*. Royal Society; 2015;11:20150552. <https://doi.org/10.1098/rsbl.2015.0552>
39. Hanson KC, Cooke SJ, Suski CD, Niezgodá G, Phelan FJS, Tinline R, et al. Assessment of largemouth bass (*Micropterus salmoides*) behaviour and activity at multiple spatial and temporal scales utilizing a whole-lake telemetry array. *Hydrobiologia*. 2007;582:243–56. <https://doi.org/10.1007/s10750-006-0549-6>
40. Baktoft H, Aarestrup K, Berg S, Boel M, Jacobsen L, Jepsen N, et al. Seasonal and diel effects on the activity of northern pike studied by high-resolution positional telemetry. *Ecology of Freshwater Fish*. 2012;21:386–94. <https://doi.org/10.1111/j.1600-0633.2012.00558.x>

41. Doka SE, Munawar M, Midwood JD, Wells MG. The state of Toronto and Region's ecosystem: Synthesis and highlights. *Aquatic Ecosystem Health & Management*. 2018;21:362–7. <https://doi.org/10.1080/14634988.2018.1505355>
42. Leisti K, Gardner Costa J, Maceachern J, Gertzen E, Doka S. Toronto Shoreline & Nearshore Survey, 2012. 2020.
43. TRCA. Tommy Thompson Park – public urban wilderness: habitat creation and enhancement projects 1995–2000., 2000 p. 1–46.
44. Midwood JD, Rous AM, Doka SE, Cooke SJ. Acoustic telemetry in Toronto Harbour: Assessing residency, habitat selection, and within-harbour movements of fishes over a five-year period. Fisheries and Oceans Canada, Great Lakes Laboratory for Fisheries and Aquatics; 2019.
45. Veilleux M. Spatial ecology of fish in Toronto Harbour in Response to aquatic habitat enhancement [Master of Science]. [Ottawa, Ontario]: Carleton University; 2014 [cited 2025 Nov 7]. <https://doi.org/10.22215/etd/2014-10589>
46. Abrams AE, Rous AM, Brooks JL, Lawrence MJ, Midwood JD, Doka SE, et al. Comparing immobilization, recovery, and stress indicators associated with electric fish handling gloves and a portable electrosedation system. *Transactions of the American Fisheries Society*. 2018;147:390–9. <https://doi.org/10.1002/tafs.10034>
47. Pincock DG. False detections: What they are and how to remove them from detection data. Amirix Document DOC-004691 Version 03. 2012; Available: -<http://vemco.com/wp-content/uploads/2012/08/False-Detections-What-they-are-and-how-to-remove-them-from-detection-data.pdf>. <https://doi.org/10.1109/TNSRE.2002.802877>
48. Innovasea. Beyond Pings: Getting the most out of your accelerometer tags: Function, configuration and biological insights. Innovasea; 2025 p. 1–23. Report No.: DOC-9224-01.
49. Dosen JM, Remillard CYL, MacEachern J, Doolittle A, Murphy S, Doka SE. Fish habitat geodatabase and gap analysis for the Toronto and Region Area of Concern. *Can. Tech. Rep. Fish. Aquat. Sci*; 2025. Report No.: xxx: xiii +.
50. Rohweder JJ, Rogala JT, Johnson BL, Anderson D, Clark S, Chamberlin F, et al. Application of wind fetch and wave models for habitat rehabilitation and enhancement projects. Geological Survey (US); 2008.
51. Doolittle A, Bakelaar CN, Doka SE. Spatial framework for storage and analyses of fish habitat data in Great Lakes' Areas of Concern: Hamilton Harbour geodatabase case study. Great Lakes Laboratory for Fisheries and Aquatic Sciences, Central & Arctic ...; 2010.
52. Hlevca B, Cooke SJ, Midwood JD, Doka SE, Portiss R, Wells MG. Characterization of water temperature variability within a harbour connected to a large lake. *Journal of Great Lakes Research*. 2015;41:1010–23. <https://doi.org/10.1016/j.jglr.2015.07.013>
53. Wood SN. *Generalized Additive Models: An Introduction with R*. 2nd ed. Chapman and Hall/CRC; 2017.
54. Pedersen EJ, Miller DL, Simpson GL, Ross N. Hierarchical generalized additive models in ecology: An introduction with mgcv. *PeerJ*. 2019;7:e6876. <https://doi.org/10.7717/peerj.6876>
55. Rij J van, Wieling M, Baayen RH, Rijn H van. *itsadug: Interpreting Time Series and Autocorrelated Data Using GAMMs*. 2022.
56. Simpson GL. *gratia: An R package for exploring generalized additive models*. *Journal of Open Source Software*. 2024;9:6962. <https://doi.org/10.21105/joss.06962>
57. Thieurmel B, Elmarhraoui A. *suncalc: Compute Sun Position, Sunlight Phases, Moon Position and Lunar Phase [Internet]*. 2022. <https://doi.org/10.32614/CRAN.package.suncalc>
58. Peat TB, Gutowsky LFG, Doka SE, Midwood JD, Lapointe NWR, Hlevca B, et al. Comparative thermal biology and depth distribution of largemouth bass (*Micropterus salmoides*) and northern pike (*Esox lucius*) in an urban harbour of the Laurentian Great Lakes. *Can J Zool*. 2016;94:767–76. <https://doi.org/10.1139/cjz-2016-0053>
59. Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, et al. *glmmTMB: Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling*. *The R Journal*. 2017;9:378–400. <https://doi.org/10.32614/RJ-2017-066>


60. Hartig F. DHARMA: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models [Internet]. 2024. <https://doi.org/10.32614/CRAN.package.DHARMA>
61. Hlevca B, Wells MG, Cruz Font L, Doka SE, Portiss R, St. John M, et al. Water circulation in Toronto Harbour. *Aquatic Ecosystem Health & Management*. Taylor & Francis; 2018;21:234–44. <https://doi.org/10.1080/14634988.2018.1500059>
62. Wright S, Metcalfe J, Hetherington S, Wilson R. Estimating activity-specific energy expenditure in a teleost fish, using accelerometer loggers. *Mar Ecol Prog Ser*. 2014;496:19–32. <https://doi.org/10.3354/meps10528>
63. Hansen MJ, Boisclair D, Brandt SB, Hewett SW, Kitchell JF, Lucas MC, et al. Applications of bioenergetics models to fish ecology and management: Where do we go from here? *Transactions of the American Fisheries Society*. 1993;122:1019–30. [https://doi.org/10.1577/1548-8659\(1993\)122%253C1019:aobmtf%253E2.3.co;2](https://doi.org/10.1577/1548-8659(1993)122%253C1019:aobmtf%253E2.3.co;2)
64. Jeffrey JD, Hasler CT, Chapman JM, Cooke SJ, Suski CD. Linking landscape-scale disturbances to stress and condition of fish: Implications for restoration and conservation. *Integr Comp Biol*. 2015;55:618–30. <https://doi.org/10.1093/icb/icv022>
65. Bunt CM, Jacobson B, Fernandes T, Ridgway L, McMeans B. Site fidelity and seasonal habitat preferences of largemouth bass (*Micropterus salmoides*) in a temperate regulated reservoir. *Hydrobiologia*. 2021;848:2595–609. <https://doi.org/10.1007/s10750-021-04582-1>
66. Driscoll MT, Norman JD, Bennett DL, Metz BK, Smith DR. Movement, home range, and structural habitat use of the Largemouth Bass complex in two large Texas reservoirs. *North American Journal of Fisheries Management*. 2024;44:572–89. <https://doi.org/10.1002/nafm.10996>
67. Veilleux MAN, Midwood JD, Boston CM, Lapointe NWR, Portiss R, Wells M, et al. Assessing occupancy of freshwater fishes in urban boat slips of Toronto Harbour. *Aquatic Ecosystem Health & Management*. Taylor & Francis; 2018;21:331–41. <https://doi.org/10.1080/14634988.2018.1507530>
68. Diana JS, Mackay WC. Timing and Magnitude of Energy Deposition and Loss in the Body, Liver, and Gonads of Northern Pike (*Esox lucius*). *J Fish Res Bd Can*. 1979;36:481–7. <https://doi.org/10.1139/f79-071>
69. Lucas M. Spawning activity of male and female pike, *Esox lucius* L., determined by acoustic tracking. *Can J Zool*. 1992;70:191–6. <https://doi.org/10.1139/z92-029>
70. Lucas MC, Johnstone ADF, Priede IG. Use of physiological telemetry as a method of estimating metabolism of fish in the natural environment. *Transactions of the American Fisheries Society*. 1993;122:822–33. [https://doi.org/10.1577/1548-8659\(1993\)122%253C0822:UOPTAA%253E2.3.CO;2](https://doi.org/10.1577/1548-8659(1993)122%253C0822:UOPTAA%253E2.3.CO;2)
71. Miranda LE, Hubbard WD. Length-dependent winter survival and lipid composition of age-0 largemouth bass in Bay Springs Reservoir, Mississippi. *Transactions of the American Fisheries Society*. 1994;123:80–7. [https://doi.org/10.1577/1548-8659\(1994\)123%253C0080:LDWSAL%253E2.3.CO;2](https://doi.org/10.1577/1548-8659(1994)123%253C0080:LDWSAL%253E2.3.CO;2)
72. Mackereth RW, Noakes DLG, Ridgway MS. Size-based variation in somatic energy reserves and parental expenditure by male smallmouth bass, *Micropterus dolomieu*. *Developments in Environmental Biology of Fishes*. 1999:263–75.
73. Raibley PT, Irons KS, O'Hara TM, Blodgett KD, Sparks RE. Winter habitats used by largemouth bass in the Illinois River, a large river-floodplain ecosystem. *North American Journal of Fisheries Management*. Taylor & Francis; 1997;17:401–12.
74. Binder TR, Nannini MA, Wahl DH, Arlinghaus R, Klefoth T, Philipp DP, et al. Largemouth bass selected for differential vulnerability to angling exhibit similar routine locomotory activity in experimental ponds. *Trans Am Fish Soc*. 2012;141:1252–9. <https://doi.org/10.1080/00028487.2012.688919>
75. McMahan TE, Holanov SH. Foraging success of largemouth bass at different light intensities: Implications for time and depth of feeding. *Journal of Fish Biology*. 1995;46:759–67. <https://doi.org/10.1111/j.1095-8649.1995.tb01599.x>
76. Masters JEG, Hodder K, Beaumont W, Gozlan R, Pinder AC, Kenward R, et al. Spatial behaviour of pike *Esox lucius* L. in the River Frome, UK. *Proceedings of the Fifth Conference on Fish Telemetry held in Europe*. 2005.

77. Pitcher TJ, Turner JR. Danger at dawn: Experimental support for the twilight hypothesis in shoaling minnows. *Journal of Fish Biology*. 1986;29:59–70. <https://doi.org/10.1111/j.1095-8649.1986.tb04999.x>
78. Harvey B. A biological synopsis of northern pike (*Esox lucius*). Fisheries and Oceans Canada, Science Branch, Pacific Region, Pacific ...; 2009;
79. Baktoft H, Gjelland KØ, Økland F, Thygesen UH. Positioning of aquatic animals based on time-of-arrival and random walk models using YAPS (Yet Another Positioning Solver). *Scientific Reports*. 2017;7. <https://doi.org/10.1038/s41598-017-14278-z>
80. Lennox RJ, Kambestad M, Berhe S, Birnie-Gauvin K, Cooke SJ, Dahlmo LS, et al. The role of habitat in predator–prey dynamics with applications to restoration. *Restoration Ecology*. 2025;33:e14354. <https://doi.org/10.1111/rec.14354>
81. Gleiss AC, Wilson RP, Shepard ELC. Making overall dynamic body acceleration work: On the theory of acceleration as a proxy for energy expenditure. *Methods in Ecology and Evolution*. 2011;2:23–33. <https://doi.org/10.1111/j.2041-210X.2010.00057.x>
82. Tang RWK, Doka SE, Midwood JD, M GCJ. Development and spatial application of a submerged aquatic vegetation model for Cootes Paradise Marsh, Ontario, Canada. *Aquatic Sciences [Internet]*. Basel, Netherlands: Springer Nature B.V.; 2021 [cited 2025 Nov 7];83. <https://doi.org/10.1007/s00027-020-00760-w>
83. Kraus RT, Holbrook CM, Vandergoot CS, Stewart TR, Faust MD, Watkinson DA, et al. Evaluation of acoustic telemetry grids for determining aquatic animal movement and survival. *Methods in Ecology and Evolution*. 2018;2018:1489–502. <https://doi.org/10.1111/2041-210X.12996>
84. Legendre P, Dale MRT, Fortin M-J, Gurevitch J, Hohn M, Myers D. The consequences of spatial structure for the design and analysis of ecological field surveys. *Ecography*. 2002;25:601–15. <https://doi.org/10.1034/j.1600-0587.2002.250508.x>
85. Simberloff D. Flagships, umbrellas, and keystones: Is single-species management passe in the landscape era? *Biological Conservation*. 1998;83:247–57. [https://doi.org/10.1016/S0006-3207\(97\)00081-5](https://doi.org/10.1016/S0006-3207(97)00081-5)
86. Piczak ML, Perry D, Cooke SJ, Harrison I, Benitez S, Koning A, et al. Protecting and restoring habitats to benefit freshwater biodiversity. *Environ Rev*. 2024;32:438–56. <https://doi.org/10.1139/er-2023-0034>
87. Piczak ML, Theis S, Portiss R, Ruppert JLW, Midwood JD, Cooke SJ. Evaluating the efficacy of ecological restoration of fish habitat in coastal waters of Lake Ontario. *Science of The Total Environment*. 2024;953:176088. <https://doi.org/10.1016/j.scitotenv.2024.176088>
88. Prime L, Lake C, Matos L, Morrison B, Morton E, Portiss R, et al. The Toronto Waterfront Aquatic Habitat Restoration Strategy: A unique collaborative approach to streamlining approvals and restoring aquatic habitat. *Plan Canada*. 2013;53:26–31.
89. Doka SE, Minns CK, Valere BG, Cooke SJ, Portiss RJ, Sciscione TF, et al. An ecological accounting system for integrated aquatic planning and habitat banking with case study on the Toronto waterfront, Ontario, Canada. *Environmental management*. Springer; 2022;69:952–71.


## Tables

**Table 1.** Significant predictors of activity estimated with a generalized linear mixed effects model. Transmitter ID was included as a random effect.


Species	Predictor	$\chi^2$	df	p
<b>Largemouth Bass</b>	Season	13840.00	3	$\leq 0.0001$
	Diel Period	2606.00	3	$\leq 0.0001$
<b>Northern Pike</b>	Season	1937.80	3	$\leq 0.0001$
	Diel Period	2644.90	3	$\leq 0.0001$




**Figure 1.** Map of Lake Ontario and locations of acoustic receivers groups in Toronto Harbour points represent individual receivers in different habitat types. The following regions of Toronto Harbour are defined as the Inner Harbour (IH), Toronto Islands (TI), Outer Harbour (OH), and Tommy Thompson Park (TTP)



**Figure 2.** Maps of detection history for each season for Largemouth Bass (A) and Northern Pike (B) in Toronto Harbour. Colour denotes the number of detections for that receiver within a season while the size of the point is the number of unique individuals that were detected at that receiver.



**Figure 3.** Predicted mean acceleration values derived from Generalized Additive Mixed-effect Models for A) for Largemouth Bass and B) Northern Pike in Toronto Harbour across habitat types denoted by different symbols.



**Figure 4.** Mean acceleration ( $\text{m s}^{-2}$ ) for Largemouth Bass and Northern Pike in Toronto Harbour delineated by season (top panels a and b) and diel period (bottom panels c and d). Letters in all panels represent statistical significance among groups with alpha set to 0.05.